

[Help](#)
[Logout](#)
[Main Menu](#)
[Search Form](#)
[Posting Counts](#)
[Show S Numbers](#)
[Edit S Numbers](#)

Search Results - Record(s) 1 through 11 of 11 returned.

1. Document ID: US 5849992 A

Entry 1 of 11

File: USPT

Dec 15, 1998

US-PAT-NO: 5849992

DOCUMENT-IDENTIFIER: US 5849992 A

TITLE: Transgenic production of antibodies in milk

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KWIC	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	------	-------

2. Document ID: US 5843705 A

Entry 2 of 11

File: USPT

Dec 1, 1998

US-PAT-NO: 5843705

DOCUMENT-IDENTIFIER: US 5843705 A

TITLE: Transgenically produced antithrombin III

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KWIC	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	------	-------

3. Document ID: US 5827690 A

Entry 3 of 11

File: USPT

Oct 27, 1998

US-PAT-NO: 5827690

DOCUMENT-IDENTIFIER: US 5827690 A

TITLE: Transgenic production of antibodies in milk

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KWIC	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	------	-------

4. Document ID: US 5750172 A

Entry 4 of 11

File: USPT

May 12, 1998

US-PAT-NO: 5750172

DOCUMENT-IDENTIFIER: US 5750172 A

TITLE: Transgenic non human mammal milk

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KWIC	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	------	-------

5. Document ID: US 4873316 A

Entry 5 of 11

File: USPT

Oct 10, 1989

US-PAT-NO: 4873316

DOCUMENT-IDENTIFIER: US 4873316 A

TITLE: Isolation of exogenous recombinant proteins from the milk of transgenic mammals

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KWIC	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	------	-------

6. Document ID: US 5750172 A
Entry 6 of 11

File: EPAB

May 12, 19

PUB-NO: US005750172A
DOCUMENT-IDENTIFIER: US 5750172 A
TITLE: Transgenic non human mammal milk

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KMC	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	-------

7. Document ID: WO 9837224 A1
Entry 7 of 11

File: EPAB

Aug 27, 1998

PUB-NO: WO009837224A1
DOCUMENT-IDENTIFIER: WO 9837224 A1
TITLE: TRANSGENICALLY PRODUCED NON-SECRETED PROTEINS

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KMC	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	-------

8. Document ID: WO 9626268 A1
Entry 8 of 11

File: EPAB

Aug 29, 1996

PUB-NO: WO009626268A1
DOCUMENT-IDENTIFIER: WO 9626268 A1
TITLE: TRANSGENICALLY PRODUCED ANTITHROMBIN III

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KMC	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	-------

9. Document ID: WO 9517085 A1
Entry 9 of 11

File: EPAB

Jun 29, 1995

PUB-NO: WO009517085A1
DOCUMENT-IDENTIFIER: WO 9517085 A1
TITLE: TRANSGENIC PRODUCTION OF ANTIBODIES IN MILK

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KMC	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	-------

10. Document ID: WO 9113151 A1
Entry 10 of 11

File: EPAB

Sep 5, 1991

PUB-NO: WO009113151A1
DOCUMENT-IDENTIFIER: WO 9113151 A1
TITLE: IMPROVED EXPRESSION OF POLYPEPTIDES

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KMC	Clip Img	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	----------	-------

11. Document ID: US 4873316 A
Entry 11 of 11

File: EPAB

Oct 10, 1989

PUB-NO: US004873316A
DOCUMENT-IDENTIFIER: US 4873316 A
TITLE: Isolation of exogenous recombinant proteins from the milk of transgenic mammals

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KMC	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	-------

Terms	Documents
meade-harry.in.	11

TI Lactogenic immunity in transgenic mice producing recombinant antibodies neutralizing coronavirus.

AU Castilla J, Sola I, Pintado B, Sanchez-Morgado J M, Enjuanes L
CS Department of Molecular and Cell Biology, Centro Nacional de Biotecnología, CSIC, Madrid, Spain.

SO ADVANCES IN EXPERIMENTAL MEDICINE AND BIOLOGY, (1998) 440 675-86.
Journal code: 2LU. ISSN: 0065-2598.

CY United States
DT Journal Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199903
EW 19990303
AB Protection against coronavirus infections can be provided by the oral administration of virus neutralizing antibodies. To provide lactogenic immunity, eighteen lines of transgenic mice secreting a recombinant IgG1 monoclonal antibody (rIgG1) and ten lines of transgenic mice secreting recombinant IgA monoclonal antibodies (rIgA) neutralizing gastroenteritis coronavirus (TGEV) into the milk were generated. Genes encoding the light and heavy chains of monoclonal antibody (MAb) 6A.C3 were expressed under the control of regulatory sequences derived from the mouse genomic DNA encoding the ***whcy*** ***acidic*** ***protein*** (WAP) and beta-lactoglobulin (BLG), which are highly abundant milk proteins. The MAb 6A.C3 binds to a highly conserved epitope present in coronaviruses of several species. This MAb does not allow the selection of neutralization escaping virus mutants. The antibody was expressed in the milk of transgenic mice with titers of one million as determined by RIA, and neutralized TGEV infectivity by one million fold corresponding to ***immunoglobulin*** concentrations of 5 to 6 mg per ml. Matrix attachment regions (MAR) sequences were not essential for rIgG1 transgene expression, but co-microinjection of MAR and antibody genes led to a twenty to ten thousand-fold increase in the antibody titer in 50% of the rIgG1 transgenic animals generated. Co-microinjection of the genomic BLG gene with rIgA light and heavy chain genes led to the generation of

transgenic mice carrying the three transgenes. The highest antibody titers were produced by transgenic mice that had integrated the antibody and BLG genes, although the number of transgenic animals generated does not allow a definitive conclusion on the enhancing effect of BLG co-integration.

Antibody expression levels were transgene copy number independent and integration site dependent. The generation of transgenic animals producing virus neutralizing antibodies in the milk could be a general approach to provide protection against neonatal infections of the enteric tract.

L1 ANSWER 2 OF 4 MEDLINE
AN 1998040670 MEDLINE
DN 98040670

TI Production of active anti-CD6 mouse/human chimeric antibodies in the milk of transgenic mice.

AU Linonta J; Pedraza A; Rodriguez A; Freyre F M; Barral A M; Leonart R; Gracia C A; Gavilondo J V; de la Fuente J
CS Mammalian Cell Genetics Division, Center for Genetic Engineering and Biotechnology, Havana, Cuba.

SO IMMUNOTECHNOLOGY, (1995 Aug) 1 (2) 107-13.
Journal code: CR0. ISSN: 1380-2933.

CY Netherlands
DT Journal Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199803
EW 19980302
AB The expression of chimeric genes in the mammary gland of transgenic farm animals has become an alternative for the large-scale production of recombinant proteins and for the modification of milk composition. In this paper, we show that a mouse/human chimeric antibody against the human CD6 leukocyte antigen can be assembled and correctly folded by the mammary gland, and secreted to milk, where it maintains its specificity. The base sequences encoding for the heavy and light chain variable regions of the anti-CD6 mouse monoclonal antibody IOR-T1 were cloned by the polymerase chain reaction from hybridoma cDNA, coupled to human heavy and light chain constant region genes, and inserted in a vector containing the 5' regulatory region of the rabbit ***whcy*** ***acidic*** ***protein*** gene. Transgenic mice were produced by conventional

*****STN Columbus*****
**

FILE 'HOME' ENTERED AT 16:55:03 ON 04 AUG 1999

=> file medline

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
ENTRY	SESSION	
0.15	0.15	

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 16:55:09 ON 04 AUG 1999

FILE LAST UPDATED: 30 JUL 1999 (19990730/UP). FILE COVERS 1960 TO DATE.

MEDLINE has been reloaded to reflect the annual MeSH changes made by the National Library of Medicine for 1999. Enter HELP RLOAD for details.

OLDMEDLINE, data from 1960 through 1965 from the Cumulated Index Medicus (CUM), has been added to MEDLINE. See HELP CONTENT for details.

Left, right, and simultaneous left and right truncation are available in the Basic Index. See HELP SFIELDS for details.

THIS FILE CONTAINS CAS REGISTRY NUMBERS FOR EASY AND ACCURATE SUBSTANCE IDENTIFICATION.

=> s immunoglobulin and whey acidic protein/ab,bi

'AB' IS NOT A VALID FIELD CODE

66111 IMMUNOGLOBULIN

0 WHEY ACIDIC PROTEIN/AB

1812 WHEY/BI

32864 ACIDIC/BI

829012 PROTEIN/BI

145 WHEY ACIDIC PROTEIN/BI

((WHEY(W)ACIDIC(W)PROTEIN(BI))

L1 4 IMMUNOGLOBULIN AND WHEY ACIDIC PROTEIN/AB,BI

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 4 ANSWERS - CONTINUE? Y(N)?y

L1 ANSWER 1 OF 4 MEDLINE
AN 1998455664 MEDLINE
DN 98455664

pronuclei microinjection techniques. Integration and transgene copy number were determined by Southern blot. Assembled human ***immunoglobulin*** was detected in milk using a sandwich ELISA. Expression levels of chimeric antibodies in milk were determined to be around 400 micrograms/ml by Western blot, using CHO-derived chimeric IOR-T1 antibodies as reference. The chimeric antibodies produced in milk recognized human peripheral blood T lymphocytes by indirect immunofluorescence, with the classical patch-like pattern of IOR-T1.

L1 ANSWER 3 OF 4 MEDLINE
AN 95358828 MEDLINE
DN 95358828

T1 The effect of various introns and transcription terminators on the efficiency of expression vectors in various cultured cell lines and in the mammary gland of transgenic mice.
AU Petitclerc D; Attal J; Theron M C; Bearzotti M; Bolifraud P; Kann G; Sinnakre M G; Pointu H; Puissant C; Houdebine L M
CS Agriculture et Agro-Alimentaire Canada, Est Lennoxville, Quebec.
SO JOURNAL OF BIOTECHNOLOGY, (1995 Jun 21) 40 (3) 169-78.
Journal code: AL6. ISSN: 0168-1656.

CY Netherlands
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; B
EM 199511
AB Various combinations of promoters, introns and transcription terminators were used to drive the expression of bovine growth hormone (bGH) cDNA in different cell types. In constructs containing the human cytomegalovirus (hCMV) promoter and the SV40 late genes terminator, the intron from SV40 genes (VP1) was much more efficient, than the intron from the early genes (i). The synthetic intron SIS generated by the association of an adenovirus splice donor and an ***immunoglobulin*** G splice acceptor showed the highest activity. The respective potency of these introns was similar in several mammalian (CHO, HC11 and COS) and fish (TO2 and EPC) cells. The rabbit ****whye*** ***acidic*** ***protein*** (WAP) gene promoter was highly efficient to drive the expression of bGH gene in the HC11 mammary cell lines. In contrast, the bGH cDNA under

the control of the same promoter was much less efficiently expressed when the SV40 VP1 intron and transcription terminator were used. The rabbit WAP gene and the human GH gene terminators did not or only moderately enhanced the expression of the construct WAP bGH cDNA. Introduction of a promoter sequence from the mouse mammary tumor virus (MMTV) LTR in the VP1 intron increased very significantly the expression of the WAP bGH cDNA. Although several of these vectors showed high potency when expressed stably in HC11 cells, all of them were only moderately efficient in transgenic mice. These data indicate that the VP1 and the SIS introns may be used to express foreign cDNAs with good efficiency in different cell types. The addition of an enhancer within an intron may still reinforce its efficiency. However, transfection experiments, even when stable expression is carried out, are poorly predictive of the potential efficiency of a vector in transgenic animals.

L1 ANSWER 4 OF 4 MEDLINE
AN 93173503 MEDLINE
DN 93173503

T1 T1, an ***immunoglobulin*** superfamily member, is expressed in H-ras-dependent epithelial tumours of mammary cells.
AU Rossler U; Andres A C; Reichmann E; Schmah W; Werenskiold A K
CS Department of Cell Chemistry, GSF-Forschungszentrum fur Umwelt und Gesundheit, Neuherberg, Germany.
SO ONCOGENE, (1993 Mar) 8 (3) 609-17.
Journal code: QNC. ISSN: 0950-9232.

CY ENGLAND; United Kingdom
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; Cancer Journals
EM 199305
AB T1 is a glycosylated protein in the carcinoembryonic antigen (CEA) family of tumour marker molecules. It was originally identified by virtue of its transient induction after the expression of p21H-ras in NIH3T3 fibroblasts. Here we show that the T1 gene is activated in mammary adenocarcinomas of transgenic mice harbouring an H-ras transgene under the control of the mammary-specific ****whye*** ***acidic*** ***protein*** (WAP) promoter. By contrast, T1 mRNA was not, or only faintly, detectable in mammary carcinomas of transgenic mice bearing a

WAP-myc transgene. Thus, T1 overexpression does not appear to be a general tumour-specific phenomenon. A dependence of T1 gene expression on the action of p21H-ras is suggested by the observation of T1 mRNA in nude mouse tumours generated from H-ras-transformed cultured mammary epithelial cells. Interestingly, activation of the T1 gene is also found during the maturation of the mammary gland (3-4 weeks after birth), whereas it is absent during its terminal differentiation in pregnancy and lactation. This expression pattern suggests a role for the secreted T1 glycoprotein in the phase of epithelial proliferation of the mammary gland. It appears that p21H-ras-induced transformation of mammary epithelial cells mimics the situation occurring in puberty. In both developmental stages the T1 glycoprotein might affect cell interactions of the proliferating epithelial cells with the surrounding stroma. It might thus promote outgrowth in gland maturation as well as invasive growth of p21H-ras-transformed mammary epithelial cells.

=> s immunoglobulin# and whey acidic protein/ab,bi

'AB' IS NOT A VALID FIELD CODE
108494 IMMUNOGLOBULIN#
0 WHEY ACIDIC PROTEIN/AB
1812 WHEY/BI
32864 ACIDIC/BI
829012 PROTEIN/BI
145 WHEY ACIDIC PROTEIN/BI
(WHEY(W)ACIDIC(W)PROTEIN(BI))
L2 4 IMMUNOGLOBULIN# AND WHEY ACIDIC PROTEIN/AB,BI

=> s immunoglobulin# and beta-lactoglobulin promoter/ab,bi

'AB' IS NOT A VALID FIELD CODE
108494 IMMUNOGLOBULIN#
0 BETA-LACTOGLOBULIN PROMOTER/AB
350029 BETA/BI
1402 LACTOGLOBULIN/BI
61727 PROMOTER/BI
22 BETA-LACTOGLOBULIN PROMOTER/BI
(BETA(W)LACTOGLOBULIN(W)PROMOTER(BI))
L3 0 IMMUNOGLOBULIN# AND BETA-LACTOGLOBULIN PROMOTER/AB,BI

=> s immunoglobulin# and casein promoter/ab,bi

'AB' IS NOT A VALID FIELD CODE
108494 IMMUNOGLOBULIN#

EW 19990503
 AB Phosphorylation sites for ***casein*** kinase I were introduced into chimeric monoclonal antibody CC49 (MAB-chCC49) by inserting a synthetic fragment (CK1) encoding two ***casein*** kinase I phosphorylation sites into an expression ***vector***. The phosphorylation sites were created by incorporating the predicted consensus sequences for phosphorylation by the ***casein*** kinase I at the carboxyl terminus of the heavy-chain constant region of the MAB-chCC49. The resultant modified MAB-chCC49 (MAB-chCC49CK1) was expressed and purified. The MAB-chCC49CK1 protein can be phosphorylated by the ***casein*** kinase I with [gamma-32P]ATP to high radiospecific activity. The 32P-labeled MAB-chCC49CK1 protein binds to cells expressing TAG-72 antigens. The introduction of phosphorylation sites into MAB provides new reagents for the diagnosis and treatment of cancer. This demonstrates that, as was described for the cAMP-dependent protein kinase site, the ***casein*** kinase I recognition site can also be used to introduce phosphorylation sites into proteins. Copyright 1999 Academic Press.

L12 ANSWER 2 OF 8 MEDLINE
 AN 97138303 MEDLINE
 DN 97138303
 TI Varicella-zoster virus Fc receptor gE glycoprotein: serine/threonine and tyrosine phosphorylation of monomeric and dimeric forms.
 AU Olson J K; Bishop G A; Grose C
 CS Department of Microbiology and Immunology Program, University of Iowa
 College of Medicine, Iowa City 52242, USA.
 NC A127795 (NIAID)
 A128847 (NIAID)
 SO JOURNAL OF VIROLOGY, (1997 Jan) 71 (1) 110-9.
 Journal code: KCV. ISSN: 0022-538X.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals; Cancer Journals
 EM 199704
 EW 19970402
 AB Varicella-zoster virus (VZV) glycoprotein gE is the predominant viral cell surface molecule; it behaves as an Fc receptor for ***immunoglobulin***
 G, but its central function may be more closely related to viral

0 CONSTRUCT#/AB
 26298 CONSTRUCT#/BI
 0 VECTOR#/AB
 59832 VECTOR#/BI
 L9 0 L8 AND (CONSTRUCT# OR VECTOR#)/AB,BI
 => s l8 and transgen?/ab,bi
 'AB' IS NOT A VALID FIELD CODE
 0 TRANSGEN?/AB
 22862 TRANSGEN?/BI
 L10 0 L8 AND TRANSGEN?/AB,BI
 => s immunoglobulin# and casein/ab,bi
 'AB' IS NOT A VALID FIELD CODE
 108494 IMMUNOGLOBULIN#
 0 CASEIN/AB
 11523 CASEIN/BI
 L11 190 IMMUNOGLOBULIN# AND CASEIN/AB,BI
 => s l11 and (construct# or vector#)/ab,bi
 'AB' IS NOT A VALID FIELD CODE
 0 CONSTRUCT#/AB
 26298 CONSTRUCT#/BI
 0 VECTOR#/AB
 59832 VECTOR#/BI
 L12 8 L11 AND (CONSTRUCT# OR VECTOR#)/AB,BI
 => d l1-bib ab
 YOU HAVE REQUESTED DATA FROM 8 ANSWERS - CONTINUE? Y(N)?

L12 ANSWER 1 OF 8 MEDLINE
 AN 1999150486 MEDLINE
 DN 99150486
 TI Construction of phosphorylatable chimeric monoclonal antibody CC49 with a ***casein*** kinase I recognition site.
 AU Lin L; Gillies S D; Schlom J; Pestka S
 CS University of Medicine and Dentistry of New Jersey-Robert Wood Johnson
 Medical School, 675 Hoes Lane, Piscataway, New Jersey, 08854-5635, USA.
 NC R01 CA46465 (NCI)
 R01 CA52363 (NCI)
 SO PROTEIN EXPRESSION AND PURIFICATION, (1999 Feb) 15 (1) 83-91.
 Journal code: BIV. ISSN: 1046-5928.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199905

0 CASEIN PROMOTER/AB
 11523 CASEIN/BI
 61727 PROMOTER/BI
 46 CASEIN PROMOTER/BI
 ((CASEIN(W)PROMOTER)/BI)
 L4 0 IMMUNOGLOBULIN# AND CASEIN PROMOTER/AB,BI
 => s immunoglobulin# and beta-casein promoter/ab,bi
 'AB' IS NOT A VALID FIELD CODE
 108494 IMMUNOGLOBULIN#
 0 BETA-CASEIN PROMOTER/AB
 350029 BETA/BI
 11523 CASEIN/BI
 61727 PROMOTER/BI
 44 BETA-CASEIN PROMOTER/BI
 ((BETA(W)CASEIN(W)PROMOTER)/BI)
 L5 0 IMMUNOGLOBULIN# AND BETA-CASEIN PROMOTER/AB,BI
 => s immunoglobulin# and kappa-casein promoter/ab,bi
 'AB' IS NOT A VALID FIELD CODE
 108494 IMMUNOGLOBULIN#
 0 KAPPA-CASEIN PROMOTER/AB
 21261 KAPPA/BI
 11523 CASEIN/BI
 61727 PROMOTER/BI
 0 KAPPA-CASEIN PROMOTER/BI
 ((KAPPA(W)CASEIN(W)PROMOTER)/BI)
 L6 0 IMMUNOGLOBULIN# AND KAPPA-CASEIN PROMOTER/AB,BI
 => s immunoglobulin# and lactalbumin promoter/ab,bi
 'AB' IS NOT A VALID FIELD CODE
 108494 IMMUNOGLOBULIN#
 0 LACTALBUMIN PROMOTER/AB
 2108 LACTALBUMIN/BI
 61727 PROMOTER/BI
 3 LACTALBUMIN PROMOTER/BI
 ((LACTALBUMIN(W)PROMOTER)/BI)
 L7 0 IMMUNOGLOBULIN# AND LACTALBUMIN PROMOTER/AB,BI
 => s immunoglobulin# and lactalbumin/ab,bi
 'AB' IS NOT A VALID FIELD CODE
 108494 IMMUNOGLOBULIN#
 0 LACTALBUMIN/AB
 2108 LACTALBUMIN/BI
 96 IMMUNOGLOBULIN# AND LACTALBUMIN/AB,BI
 => s l8 and (construct# or vector#)/ab,bi
 'AB' IS NOT A VALID FIELD CODE

egress and cell-to-cell spread. To further analyze the receptor properties of VZV gE, the gE gene (also called open reading frame 68) was expressed by a baculovirus ***vector*** in insect cells. The recombinant baculovirus gE product had a molecular mass of 64 kDa, smaller than the previously documented 98 kDa of mature gE expressed in mammalian cells. The major reason for the lowered molecular mass was diminished glycosylation. In addition to the 64-kDa form, a larger (130-kDa) form was observed in insect cells and represented dimerized 64-kDa molecules. Both the monomeric and dimeric gE forms were highly phosphorylated in insect cells. Protein kinase assays conducted in vitro with [gamma-32P]ATP and [gamma-32P]GTP indicated that endogenous ***casein*** kinase II was phosphorylating monomeric gE, while the dimeric gE form was phosphorylated by another kinase which did not utilize [gamma-32P]GTP. When immobilized recombinant gE molecules were probed with a monoclonal antibody which specifically recognizes a phosphotyrosine linkage, the gE dimer was found to be tyrosine phosphorylated whereas the monomer was not similarly modified. When recombinant gE produced in HeLa cells was probed with the same antiphosphotyrosine antibody, a dimeric gE form at 130 kDa was detected on the cell surface. These results suggested that VZV gE closely resembled other cell surface receptors, being modified on its various forms by both serine/threonine and tyrosine protein kinases. In this case, tyrosine phosphorylation occurred on a previously unrecognized and underglycosylated VZV gE dimeric product.

L12 ANSWER 3 OF 8 MEDLINE
AN 96375171 MEDLINE
DN 96375171
TI Efficient vasoactive intestinal polypeptide hydrolyzing autoantibody light chains selected by phage display.
AU Tyutyulkova S; Gao Q S; Thompson A; Remard S; Paul S
CS Department of Anesthesiology, University of Nebraska Medical Center,
Omaha, USA.
NC HL44126 (NHLBI)

A31268 (NIAID)
SO BIOCHIMICA ET BIOPHYSICA ACTA, (1996 Aug 23) 1316 (3) 217-23.
Journal code: AOW. ISSN: 0006-3002.
CY Netherlands
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; Cancer Journals
OS GENBANK-L43498; GENBANK-L43499
EM 199612
AB An ***immunoglobulin*** light chain (L chain) library derived from the peripheral blood lymphocytes of a patient with asthma was cloned into a phagemid ***vector***. Phage particles displaying L chains capable of binding vasoactive intestinal polypeptide (VIP) were isolated by affinity chromatography. Two VIP binding L chains were expressed in *Escherichia coli* in soluble form and purified to electrophoretic homogeneity by metal chelating and protein L affinity chromatography. Both L chains catalyzed the hydrolysis of [Iyr10-125]VIP substrate. The catalytic activity eluted at the molecular mass of the monomer form of the L chain (28 kDa) from a gel filtration column. The activity was bound by immobilized anti-kappa-chain antibody. A control recombinant L chain displayed no catalytic activity. Hydrolysis of VIP by the catalytic L chains was saturable and consistent with Michaelis-Menten kinetics. The turnover of the L chains was moderate (0.22 and 2.21/min) and their Km values indicated comparatively high affinity recognition of VIP[111 and 202 nM]. producing catalytic efficiencies comparable to or greater than trypsin. Unlike trypsin, the L chains did not display detectable cleavage of ***casein***, suggesting a catalytic activity specialized for VIP. Comparisons of the nucleotide sequences of the L chain cDNA with their putative germ-line counterparts suggested the presence of several replacement mutations in the complementarity determining regions (CDRs). These observations suggest: (a) Retention or acquisition of catalytic activity by the L chains is compatible with affinity maturation of antibodies; and (b) The autoimmune L chain repertoire can serve as a source of substrate-specific and efficient catalysts.

L12 ANSWER 4 OF 8 MEDLINE
AN 93234096 MEDLINE
DN 93234096
TI Expression of Porphyromonas gingivalis proteolytic activity in *Escherichia*

coli.
AU Madden T E; Thompson T M; Clark V L
CS Department of Dental Research, University of Rochester, New York.
NC 5R01 DE08512 (NIDR)
5K16 DE00159 (NIDR)
SO ORAL MICROBIOLOGY AND IMMUNOLOGY, (1992 Dec) 7 (6) 349-56.
Journal code: ORA. ISSN: 0902-0055.
CY Denmark
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Dental Journals; Dental
EM 199307
AB Porphyromonas gingivalis (formerly Bacteroides gingivalis) degrades numerous protein substrates including collagen, fibrinogen, fibronectin, gelatin, ***casein***, ***immunoglobulins*** and complement components. In order to clone one or more of these protease genes, a genomic library was constructed with Sau3A1 restriction fragments of chromosomal DNA from *P. gingivalis* ATCC 33277 ligated into the temperature-regulated ***vector*** pCQV2, and expressed in *Escherichia coli* DH5 alpha mcr. The electro-transformants (3 x 10(4)) were screened for general protease activity on Luria broth agar containing ampicillin (50 mg/l) and sodium caseinate (2%). One ***casein***-hydrolyzing clone was detected and subcultured, and the activity of the cell extracts was characterized. We were able to show that the protease-positive clone, (pTEM1), had broad substrate specificity. Colorimetric assays indicated the hydrolysis of azocoll, azocasein, collagen, elastin-congo red and artificial substrates. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis was used to confirm that collagen, ***casein***, fibrinogen and fibronectin were degraded by the clone.

L12 ANSWER 5 OF 8 MEDLINE
AN 93206099 MEDLINE
DN 93206099
TI Effect of PU.1 phosphorylation on interaction with NF-EM5 and transcriptional activation.
AU Pongubala J M; Van Beveren C; Nagulapalli S; Klemsz M J; McKercher S R;
Maki R A; Atchison M L
CS Department of Animal Biology, University of Pennsylvania, School of Veterinary Medicine, Philadelphia 19104.
NC GM 42415 (NIGMS)
AI 30656 (NIAID)

CA 42909 (NCI)
SO SCIENCE, (1993 Mar 12) 259 (5101) 1622-5.
Journal code: U17. ISSN: 0036-8075.
CY United States
DT Journal: Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; Cancer Journals
EM 199306
AB PU.1 recruits the binding of a second B cell-restricted nuclear factor, NF-EM5, to a DNA site in the ***immunoglobulin*** kappa 3' enhancer.
DNA binding by NF-EM5 requires a protein-protein interaction with PU.1 and specific DNA contacts. Dephosphorylated PU.1 bound to DNA but did not interact with NF-EM5. Analysis of serine-to-alanine mutations in PU.1 indicated that serine 148 (Ser148) is required for protein-protein interaction. PU.1 produced in bacteria did not interact with NF-EM5.
Phosphorylation of bacterially produced PU.1 by purified ***casein*** kinase II modified it to a form that interacted with NF-EM5 and that recruited NF-EM5 to bind to DNA. Phosphopeptide analysis of bacterially produced PU.1 suggested that Ser148 is phosphorylated by ***casein*** kinase II. This site is also phosphorylated in vivo. Expression of wild-type PU.1 increased expression of a reporter ***construct*** containing the PU.1 and NF-EM5 binding sites nearly sixfold, whereas the Ser148 mutant form only weakly activated transcription. These results demonstrate that phosphorylation of PU.1 at Ser148 is necessary for interaction with NF-EM5 and suggest that this phosphorylation can regulate transcriptional activity.

L12 ANSWER 6 OF 8 MEDLINE
AN 92260636 MEDLINE
DN 92260636
TI Receptor properties of two varicella-zoster virus glycoproteins, gpI and gpIV, homologous to herpes simplex virus gE and gI.
AU Litwin V; Jackson W; Grose C
CS Department of Microbiology, University of Iowa College of Medicine, Iowa City 52242.
NC A122795 (NIAID)
SO JOURNAL OF VIROLOGY, (1992 Jun) 66 (6) 3643-51.
Journal code: KCV. ISSN: 0022-538X.
CY United States
DT Journal: Article; (JOURNAL ARTICLE)
LA English

FS Priority Journals; Cancer Journals
EM 199208
AB The varicella-zoster virus (VZV) genome contains 70 reading frames (ORF), 5 of which encode the glycoproteins gpI, gpII, gpIII, gpIV, and gpV. ORF 67 and 68 lie adjacent to each other in the unique short region of the VZV genome and code for gpIV and gpI, respectively. These two genes, which are contained within the HindIII C fragment of the VZV genome, were subcloned in the correct orientation downstream from the promoter regions of the eukaryotic expression ***vectors*** pCMV5 and pBJ. After transfection, 5 to 20% of the Cos cells bound antibody specific for the given glycoprotein. In this study, it was shown that only the cells transfected with the gpI ***construct*** bound to the Fc fragment of human ***immunoglobulin*** G. Neither the transfected gpIV gene product nor the ***vector*** only bound to the Fc fragment. Thus, VZV gpI is confirmed to be the VZV-encoded Fc-binding glycoprotein. Like the wild-type form of gpI expressed in VZV-infected cells, gpI precipitated from transfected cells contained both N-linked and O-linked glycans and was heavily sialated. In addition, the transfected gpI gene product phosphorylated both in cell culture and in protein kinase assays by mammalian ***casein*** kinases I and II. Extensive computer-assisted analyses of the VZV gpI sequence, as well as those of alpha herpesviral homolog glycoproteins, disclosed properties similar to those of other cell surface receptors; these included (i) exocytoplasmic regions rich in cysteine residues, (ii) membrane-proximal regions with potential O-linked glycosylation sites, and (iii) cytoplasmic domains with consensus phosphorylation sites.

L12 ANSWER 7 OF 8 MEDLINE
AN 91060583 MEDLINE
DN 91060583
TI Primary structure of the target of calcium ***vector*** protein of amphioxus.
AU Takagi T; Cox J A
CS Biological Institute, Faculty of Science, Tohoku University, Sendai, Japan.
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1990 Nov 15)

265 (32) 19721-7.
Journal code: HIV. ISSN: 0021-9258.
CY United States
DT Journal: Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; Cancer Journals
EM 199103
AB CaVPT, a target protein of Ca2(+)- ***vector*** from amphioxus muscle, was purified from its complex with CaVP after dissociation by 6 M urea and chromatographies on DEAE-cellulose and calmodulin-Sepharose. The amino acid sequence of CaVPT has been determined. The protein is composed of 243 residues and possesses an unblocked N terminus. Its molecular weight is 26,621, distinctly lower than the apparent molecular weight deduced from electrophoresis on sodium dodecyl sulfate-containing gels. CaVPT contains a potential Asn-linked glycosylation site, four potential protein kinase C phosphorylation sites, and two ***casein*** kinase II phosphorylation sites. From the sequence the following three particular domains can be inferred: a collagen-like N-terminal segment, rich in Pro and Ala, that resembles the N-terminal segment of skeletal muscle myosin light chain kinase; next to it (from residues 33 to 50) is located a strongly amphiphilic and basic alpha-helical segment which likely binds the calcium ***vector*** protein since a proteolytic cut after Arg50, occurring occasionally during the purification of CaVPT, impairs the binding to immobilized calmodulin. This segment is followed by two ***immunoglobulin*** folds. The two ***immunoglobulin*** folds typically belong to the C2 subclass and particularly resemble those present in the neural cell surface adhesion molecules NCAM, L1, F11, MAG, TAG-1, fasciclin II, and amalgam. Recently, the presence of ***immunoglobulin*** folds of this type has been reported in some intracellular muscular proteins, namely in smooth muscle myosin light chain kinase, striated muscle C protein and titin, as well as in the nematode 600-kDa protein twitchin. From this structural study we can formulate the working hypothesis that CaVPT acts on the structure of the thick filament in muscle or regulates, perhaps via other ***immunoglobulin*** fold-containing proteins.

L12 ANSWER 8 OF 8 MEDLINE
 AN 83164366 MEDLINE
 DN 83164366
 TI Site-directed point mutation in the src gene of Rous sarcoma virus results in an inactive src gene product.
 AU Bryant D; Parsons J T
 NC CA29243 (NCI)
 CA27578 (NCI)
 SO JOURNAL OF VIROLOGY, (1983 Mar) 45 (3) 1211-6.
 Journal code: KCV. ISSN: 0022-538X.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 OS GENBANK-J02351
 EM 198307
 AB Site-directed mutagenesis techniques were used to defined point mutations within the src gene of the Prague A strain of Rous sarcoma virus. Bisulfite mutagenesis at a Bg/I restriction site in the src gene yielded three mutations which contained the same single base change, a guanine-to-adenine transition. The resulting genomes encoded an src protein containing a substitution of threonine for alanine at amino position 433. Transfection of chicken cells with mutagenized DNA did not result in cellular transformation even though the cells produced a pp60src. Immune complexes containing mutant pp60src did not phosphorylate
 immunoglobulin G heavy chain or ***casein***
 => d his

(FILE 'HOME' ENTERED AT 16:55:03 ON 04 AUG 1999)

FILE 'MEDLINE' ENTERED AT 16:55:09 ON 04 AUG 1999
 L1 4 S IMMUNOGLOBULIN AND WHEY ACIDIC
 L2 4 S IMMUNOGLOBULIN AND WHEY ACIDIC
 L3 0 S IMMUNOGLOBULIN# AND
 L4 0 S IMMUNOGLOBULIN PROMOTER/AB,BI
 L5 0 S IMMUNOGLOBULIN# AND CASEIN
 L6 0 S IMMUNOGLOBULIN# AND BETA-CASEIN
 L7 0 S IMMUNOGLOBULIN# AND KAPPA-CASEIN
 L8 96 S IMMUNOGLOBULIN# AND

LACTALBUMIN/AB,BI
 L9 0 S L8 AND (CONSTRUCT# OR VECTOR#)/AB,BI
 L10 0 S L8 AND TRANSGEN#/AB,BI
 L11 190 S IMMUNOGLOBULIN# AND CASEIN/AB,BI
 L12 8 S L11 AND (CONSTRUCT# OR VECTOR#)/AB,BI
 => file medline embase biosis inpadoc caplus
 COST IN U.S. DOLLARS ENTRY SESSION TOTAL
 FULL ESTIMATED COST 7.14 7.29
 FILE 'MEDLINE' ENTERED AT 17:05:15 ON 04 AUG 1999
 FILE 'EMBASE' ENTERED AT 17:05:15 ON 04 AUG 1999
 COPYRIGHT (C) 1999 Elsevier Science B.V. All rights reserved.
 FILE 'BIOSIS' ENTERED AT 17:05:15 ON 04 AUG 1999
 COPYRIGHT (C) 1999 BIOSIS(R)
 FILE 'INPADOC' ENTERED AT 17:05:15 ON 04 AUG 1999
 COPYRIGHT (C) 1999 European Patent Office, Vienna (EPO)
 FILE 'CAPLUS' ENTERED AT 17:05:15 ON 04 AUG 1999
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 1999 AMERICAN CHEMICAL SOCIETY (ACS)
 => s II

'AB' IS NOT A VALID FIELD CODE
 'AB' IS NOT A VALID FIELD CODE
 'AB' IS NOT A VALID FIELD CODE
 'AB' IS NOT A VALID FIELD CODE
 L13 13 L1
 => dup rem l13
 PROCESSING COMPLETED FOR L13
 L14 6 DUP REM L13 (7 DUPLICATES REMOVED)
 => d l-bib ab
 YOU HAVE REQUESTED DATA FROM 6 ANSWERS -
 CONTINUE? Y(N)?

L14 ANSWER 1 OF 6 MEDLINE DUPLICATE 1
 AN 1998455664 MEDLINE
 DN 98455664
 TI Lactogenic immunity in transgenic mice producing recombinant antibodies neutralizing coronavirus.
 AU Castilla J; Sola I; Pintado B; Sanchez-Morgado J M; Enjuanes L
 CS Department of Molecular and Cell Biology, Centro Nacional de Biotecnologia, CSIC, Madrid, Spain.

SO ADVANCES IN EXPERIMENTAL MEDICINE AND BIOLOGY, (1998) 440 675-86.
 Journal code: 2LU. ISSN: 0065-2598.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199903
 EW 199903
 AB Protection against coronavirus infections can be provided by the oral administration of virus neutralizing antibodies. To provide lactogenic immunity, eighteen lines of transgenic mice secreting a recombinant IgG1 monoclonal antibody (rIgG1) and ten lines of transgenic mice secreting recombinant IgA monoclonal antibodies (rIgA) neutralizing gastroenteritis coronavirus (TGEV) into the milk were generated. Genes encoding the light and heavy chains of monoclonal antibody (MAb) 6A.C3 were expressed under the control of regulatory sequences derived from the mouse genomic DNA encoding the ***whey*** ***acidic*** ***protein*** (WAP) and beta-lactoglobulin (BLG), which are highly abundant milk proteins. The MAb 6A.C3 binds to a highly conserved epitope present in coronaviruses of several species. This MAb does not allow the selection of neutralization escaping virus mutants. The antibody was expressed in the milk of transgenic mice with titers of one million as determined by RIA, and neutralized TGEV infectivity by one million fold corresponding to ***immunoglobulin*** concentrations of 5 to 6 mg per ml. Matrix attachment regions (MAR) sequences were not essential for rIgG1 transgene expression, but co-microinjection of MAR and antibody genes led to a twenty to ten thousand-fold increase in the antibody titer in the rIgG1 transgenic animals generated. Co-microinjection of the BLG gene with rIgA light and heavy chain genes led to the generation of transgenic mice carrying the three transgenes. The highest antibody titers were produced by transgenic mice that had integrated the antibody and BLG genes, although the number of transgenic animals generated does not allow

a definitive conclusion on the enhancing effect of BLG co-integration.
 Antibody expression levels were transgene copy number independent and integration site dependent. The generation of transgenic animals producing virus neutralizing antibodies in the milk could be a general approach to provide protection against neonatal infections of the enteric tract.

L14 ANSWER 2 OF 6 EMBASE COPYRIGHT 1999 ELSEVIER SCI B V
 AN 1998119750 EMBASE
 TI Engineering passive immunity in transgenic mice secreting virus neutralizing antibodies in milk.
 AU Castilla J.; Pintado B.; Sola I.; Sanchez-Morgado J.M.; Enjuanes L.
 CS L. Enjuanes, Department of Molecular/Cell Biology, Centro Nacional de Biotecnología, Consejo Superior Invest. Científicas, Cantoblanco, 28049 Madrid, Spain. L.Enjuanes@cnb.uam.es
 SO Nature Biotechnology, (1998) 16/4 (349-354).
 Refs: 42
 ISSN: 1087-0156 CODEN: NABIF
 CY United States
 DT Journal: Article
 FS 004 Microbiology
 007 Pediatrics and Pediatric Surgery
 026 Immunology, Serology and Transplantation
 LA English
 SL English
 AB Protection against enteric infections can be provided by the oral administration of pathogen-neutralizing antibodies. To provide passive immunity, 18 lines of transgenic mice secreting a recombinant mono-clonal antibody (Mab) neutralizing transmissible gastroenteritis coronavirus (TGEV) into the milk were generated. The genes encoding a chimeric Mab with the variable modules of the murine TGEV-specific Mab 6A.C3 and the constant modules of a human IgG, isotype Mab were expressed under the control of regulatory sequences derived from the ***whey***acidic*** ***protein***, which is an abundant milk protein. The Mab 6A.C3 binds to a highly conserved epitope present in coronaviruses of several species, which does not allow the selection of neutralization escape mutants. Antibody expression titers of 104 were obtained in the milk of transgenic mice that reduced TGEV infectivity 104-fold. The antibody was synthesized at high levels throughout lactation. Integration

of matrix attachment region sequences with the antibody genes led to a 20- to 10,000-fold increase in the antibody titer in 50% of the transgenic animals. Antibody expression levels were transgene copy number independent and related to the site of integration. The generation of transgenic animals producing virus neutralizing antibodies in milk could provide an approach to protection against neonatal infections of the enteric tract.

L14 ANSWER 3 OF 6 MEDLINE DUPLICATE 2
 AN 95358828 MEDLINE
 DN 95358828
 TI The effect of various introns and transcription terminators on the efficiency of expression vectors in various cultured cell lines and in the mammary gland of transgenic mice.
 AU Peticlerc D; Attal J; Theron M C; Bearzotti M; Bolifraud P; Kann G; Stinnakre M G; Pointu H; Puissant C; Houdebine L M
 CS Agriculture et Agro-Alimentaire Canada, Est Lennoxville, Quebec.
 SO JOURNAL OF BIOTECHNOLOGY, (1995 Jun 21) 40 (3) 169-78.
 Journal code: AL6 ISSN: 0168-1656.
 CY Netherlands
 DT Journal: Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals; B
 EM 199511
 AB Various combinations of promoters, introns and transcription terminators were used to drive the expression of bovine growth hormone (bGH) cDNA in different cell types. In constructs containing the human cytomegalovirus (hCMV) promoter and the SV40 late genes terminator, the intron from SV40 genes (VP1) was much more efficient, than the intron from the early genes (i). The synthetic intron SIS generated by the association of an adenovirus splice donor and an ***immunoglobulin*** G splice acceptor showed the highest activity. The respective potency of these introns was similar in several mammalian (CHO, HC11 and COS) and fish (TO2 and EPC) cells. The rabbit ***whey*** ***acidic*** ***protein*** (WAP) gene promoter was highly efficient to drive the expression of bGH in the HC11 mammary cell lines. In contrast, the bGH cDNA under the control of the same promoter was much less efficiently expressed when the SV40 VP1

intron and transcription terminator were used. The rabbit WAP gene and the human GH gene terminators did not or only moderately enhanced the expression of the construct WAP bGH cDNA. Introduction of a promoter sequence from the mouse mammary tumor virus (MMTV) LTR in the VP1 intron increased very significantly the expression of the WAP bGH cDNA. Although several of these vectors showed high potency when expressed stably in HC11 cells, all of them were only moderately efficient in transgenic mice. These data indicate that the VP1 and the SIS introns may be used to express foreign cDNAs with good efficiency in different cell types. The addition of an enhancer within an intron may still reinforce its efficiency. However, transfection experiments, even when stable expression is carried out, are poorly predictive of the potential efficiency of a vector in transgenic animals.

L14 ANSWER 4 OF 6 MEDLINE DUPLICATE 3
 AN 1998040670 MEDLINE
 DN 98040670
 TI Production of active anti-CD6 mouse/human chimeric antibodies in the milk of transgenic mice.
 AU Limonta J; Pedraza A; Rodriguez A; Freyre F M; Barral A M; Castro F O; Leonart R; Gracia C A; Gavilondo J V; de la Fuente J
 CS Mammalian Cell Genetics Division, Center for Genetic Engineering and Biotechnology, Havana, Cuba.
 SO IMMUNOTECHNOLOGY, (1995 Aug) 1 (2) 107-13.
 Journal code: CR0 ISSN: 1380-2933.
 CY Netherlands
 DT Journal: Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199803
 EW 19980302
 AB The expression of chimeric genes in the mammary gland of transgenic farm animals has become an alternative for the large-scale production of recombinant proteins and for the modification of milk composition. In this paper, we show that a mouse/human chimeric antibody against the human CD6 leukocyte antigen can be assembled and correctly folded by the mammary gland, and secreted to milk, where it maintains its specificity. The base sequences encoding for the heavy and light chain variable regions of the anti-CD6 mouse monoclonal antibody IOR-T1 were cloned by the

polymerase chain reaction from hybridoma cDNA, coupled to human heavy and light chain constant region genes, and inserted in a vector containing the 5' regulatory region of the rabbit ***whey*** ***acidic*** ***protein*** gene. Transgenic mice were produced by conventional pronuclei microinjection techniques. Integration and transgene copy number were determined by Southern blot. Assembled human ***immunoglobulin*** was detected in milk using a sandwich ELISA. Expression levels of chimeric antibodies in milk were determined to be around 400 micrograms/ml by Western blot, using CHO-derived chimeric IOR-T1 antibodies as reference. The chimeric antibodies produced in milk recognized human peripheral blood T lymphocytes by indirect immunofluorescence, with the classical patch-like pattern of IOR-T1.

L14 ANSWER 5 OF 6 BIOSIS COPYRIGHT 1999 BIOSIS
AN 1993:342690 BIOSIS
DN PRE199396039690
T1 A novel gene product associated with mu chains in immature B cells.
AU Shirasawa, Takui; Ohnishi, Kazuo; Hagiwara, Shinji; Shigemoto, Kazuhiro; Takebe, Yutaka; Rajewsky, Klaus; Takemori, Toshitada (1)
CS (1) Dep. Immunol., NIH, Tokyo Japan
SO EMBO (European Molecular Biology Organization) Journal, (1993) Vol. 12, No. 5, pp. 1827-1834.
ISSN: 0261-4189.
DT Article
LA English
AB A previously unreported B cell specific gene, which we have named 8HS-20, was isolated from the cDNA library of a pre-B cell clone by subtraction and differential hybridization. This gene is selectively expressed as a 0.75 kb transcript in pre-B and bone marrow-derived B cell lines; a transcript of the same size is also found in bone marrow and, albeit at low levels, in spleen. The deduced amino acid sequence of the 8HS-20 cDNA displayed homology to a B cell specific gene, VpreB-1, and to members of the ***immunoglobulin*** supergene family including V-lambda, V-kappa, V-H, TCRV-alpha, V-beta and CD8. Biochemical analysis using purified antiserum against 8HS-20 oligopeptides indicates that the gene encodes proteins with mol. wts of 13.5, 14, 15.5 and 16 kDa, which

associate with mu chains in pre-B cell lines, and that these molecules are expressed concomitantly with VpreB-1 and lambda-5 gene products in the same cell lines.

L14 ANSWER 6 OF 6 MEDLINE DUPLICATE 4
AN 93173503 MEDLINE
DN 93173503
T1 T1, an ***immunoglobulin*** superfamily member, is expressed in H-ras-dependent epithelial tumours of mammary cells.
AU Rossler U; Andres A C; Reichmann E; Schmal W; Werenkiold A K
CS Department of Cell Chemistry, GSF-Forschungszentrum für Umwelt und Gesundheit, Neuherberg, Germany.
SO ONCOGENE; (1993 Mar) 8 (3) 609-17.
Journal code: ONC. ISSN: 0950-9232.
CY ENGLAND; United Kingdom
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; Cancer Journals
EM 199305
AB T1 is a glycosylated protein in the carcinoembryonic antigen (CEA) family of its tumour marker molecules. It was originally identified by virtue of transient induction after the expression of p21H-ras in NIH3T3 fibroblasts. Here we show that the T1 gene is activated in mammary adenocarcinomas of transgenic mice harbouring an H-ras transgene under the control of the mammary-specific ***whey*** ***acidic*** ***protein*** (WAP) promoter. By contrast, T1 mRNA was not, or only faintly, detectable in mammary carcinomas of transgenic mice bearing a WAP-myc transgene. Thus, T1 overexpression does not appear to be a general tumour-specific phenomenon. A dependence of T1 gene expression on the action of p21H-ras is suggested by the observation of T1 mRNA in nude mouse tumours generated from H-ras-transformed cultured mammary epithelial cells. Interestingly, activation of the T1 gene is also found during the maturation of the mammary gland (3-4 weeks after birth), whereas it is absent during its terminal differentiation in pregnancy and lactation. This expression pattern suggests a role for the secreted T1 glycoprotein in the phase of epithelial proliferation of the mammary gland. It appears that p21H-ras-induced transformation of mammary epithelial cells mimics

the situation occurring in puberty. In both developmental stages the glycoprotein might affect cell interactions of the proliferating epithelial cells with the surrounding stroma. It might thus promote ductal outgrowth in gland maturation as well as invasive growth of p21H-ras-transformed mammary epithelial cells.

=> s 12
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L15 23 L2
=> s 12 not 11
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L16 10 L2 NOT L1
=> dup rem 116
PROCESSING COMPLETED FOR L16
L17 10 DUP REM L16 (0 DUPLICATES REMOVED)
=> d 1- bib ab
YOU HAVE REQUESTED DATA FROM 10 ANSWERS - CONTINUE? Y(N)?
L17 ANSWER 1 OF 10 CAPLUS COPYRIGHT 1999 ACS
AN 1998:696460 CAPLUS
DN 130:108924
T1 Lactogenic immunity in transgenic mice producing recombinant antibodies neutralizing coronavirus
AU Catilla, J.; Sola, I.; Pintado, B.; Sanchez-Morgado, J. M.; Enjuanes, L.
CS Department of Molecular and Cell Biology Centro Nacional de Biotecnologia, CSIC Campus Universidad Autonoma, Madrid, 28049, Spain
SO Adv. Exp. Med. Biol. (1998), 440(Coronaviruses and Arteriviruses), 675-686
CODEN: AEMBAP; ISSN: 0065-2598
PB Plenum Publishing Corp.
DT Journal
LA English
AB Protection against coronavirus infections can be provided by the oral administration of virus neutralizing antibodies. To provide lactogenic immunity, eighteen lines of transgenic mice secreting a recombinant IgG1

monoclonal antibody (rlgG1) and ten lines of transgenic mice secreting recombinant IgA monoclonal antibodies (rlgA) neutralizing transmissible gastroenteritis coronavirus (TGEV) into the milk were generated. Genes encoding the light and heavy chains of monoclonal antibody (MAb) 6A.C3 were expressed under the control of regulatory sequences derived from the mouse genomic DNA encoding the ***whey*** ***acidic*** ***protein*** (WAP) and beta-lactoglobulin (BLG), which are highly abundant milk proteins. The MAb 6A.C3 binds to a highly conserved epitope present in coronaviruses of several species. This MAb does not allow the selection of neutralization escaping virus mutants. The antibody was expressed in the milk of transgenic mice with titers of one million as detd. by RIA, and neutralized TGEV infectivity by one million fold corresponding to Ig concns. of 5 to 6 mg per mL. Matrix attachment regions (MAR) sequences were not essential for rlgG1 transgene expression, but co-microinjection of MAR and antibody genes led to a twenty to ten thousand-fold increase in the antibody titer in 50% of the rlgG1 transgenic animals generated. Co-microinjection of the genomic BLG gene with rlgA light and heavy chain genes led to the generation of transgenic mice carrying the three transgenes. The highest antibody titers were produced by transgenic mice that had integrated the antibody and BLG genes, although the no. of transgenic animals generated does not allow a definitive conclusion on the enhancing effect of BLG co-integration. Antibody expression levels were transgene copy no. independent and integration site dependent. The generation of transgenic animals producing virus neutralizing antibodies in the milk could be a general approach to provide protection against neonatal infections of the enteric tract.

L17 ANSWER 2 OF 10 CAPLUS COPYRIGHT 1999 ACS
AN 1997:650467 CAPLUS
DN 127:315589
TI Cytochrome P450 encoding retroviral vectors and their use as antitumor agents

IN Gunzburg, Walter H.; Karle, Peter; Saller, Robert Michael
PA Bavarian Nordic Research Institute A/S, Den.;
GSF-Forschungszentrum Fur Umwelt Und Gesundheit; Gunzburg, Walter H.; Karle, Peter;
Saller, Robert Michael
SO PCT Int. Appl., 25 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN/CNT 1
PATENT NO. KIND DATE APPLICATION NO.
DATE
PI WO 9735994 A2 19971002 WO 1997-EP1585
19970327
WO 9735994 A3 19971120
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN,
CU, CZ, DE,
DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP,
KR, KZ,
LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW,
MX, NO, NZ, PL,
PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG,
US, UZ,
VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK,
ES, FI, FR, GB,
GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM,
GA, GN,
ML, MR, NE, SN, TD, TG
AU 9723827 A1 19971017 AU 1997-23827 19970327
EP 892852 A2 19990127 EP 1997-919307 19970327
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,
MC, PT,
IE, SI, LT, LV, FI, RO
NO 9804540 A 19980928 NO 1998-4540 19980928
PRAI DK 1996-352 19960327
WO 1997-EP1585 19970327
AB A replication-defective retroviral vector carrying a cytochrome P
450 gene
under transcriptional control of target cell specific regulatory
elements
or promoters, or X-ray inducible promoters is disclosed.

L17 ANSWER 3 OF 10 CAPLUS COPYRIGHT 1999 ACS
AN 1997:18384 CAPLUS
DN 126:43610
TI Animal gene therapy expression cassettes and DNA constructs for
treatment
of infectious diseases
IN Gagne, Marc
PA Immunova, Can.; Gagne, Marc
SO PCT Int. Appl., 55 pp.
CODEN: PIXXD2
DT Patent
LA English

FAN/CNT 1
PATENT NO. KIND DATE APPLICATION NO.
DATE
PI WO 9635793 A1 19961114 WO 1996-CA297
19960510
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ,
DE, DK, EE,
ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR,
LS, LT,
LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
RO, RU, SD, SE,
SG, SI
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI,
FR, GB, GR,
IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA,
GN
CA 2220472 AA 19961114 CA 1996-2220472 19960510
AU 9656416 A1 19961129 AU 1996-56416 19960510
EP 828839 A1 19980318 EP 1996-913403 19960510
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,
MC, PT,
IE, FI
JP 11505113 T2 19990518 JP 1996-533627 19960510
PRAI GB 1995-9461 19950510
WO 1996-CA297 19960510
AB The present invention relates to DNA sequences, expression
cassettes and
DNA constructs for use in therapy, specifically in gene therapy for
the
treatment of infectious diseases such as mastitis. Also included are
pharmaceutical and veterinary compns. contg. the constructs, and
cells
which have been transformed with the DNA and which are suitable
for
implantation into a host mammal. The gene therapy of infectious
diseases
can be effected in situ in targeted tissue or systemically.

L17 ANSWER 4 OF 10 CAPLUS COPYRIGHT 1999 ACS
AN 1997:6067 CAPLUS
DN 126:27673
TI Transgenic multicellular eukaryotes expressing genes for enzymes
of
post-translational modification of proteins
IN Lubon, Henryk; Drohan, William N.; Paleyanda, Rekha K.
PA American Red Cross, USA
SO PCT Int. Appl., 59 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN/CNT 1
PATENT NO. KIND DATE APPLICATION NO.
DATE
PI WO 9634966 A2 19961107 WO 1996-US6121
19960506

W: AU, CA, JP, MX

RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE

CA 2220109 AA 19961107 CA 1996-2220109 19960506
AU 9663474 AI 19961121 AU 1996-63474 19960506
PRAI US 1995-434834 19950504
WO 1996-US6121 19960506
AB Transgenic non-human multicellular organisms contg. expression cassettes

for enzyme involved in post-translational modification of proteins are

are described for use in the manuf. of proteins. The transgenic

organism most often carries genes for enzymes of post-translational modification

and the

gene for a protein of interest that is a substrate for the modification enzyme. Preferably, the genes are regulated, e.g. by development, tissue-type, or by a chem. inducer and the modified protein is secreted

into a bodily fluid. An example provides transgenic mice that

synthesize

human protein C and the processing protease PACE/furin in mammary glands

and secrete both proteins into milk. The genes are placed under control

of the mammary gland-specific promoter of the ***whey***

acidic ***protein*** gene.

L17 ANSWER 5 OF 10 CAPLUS COPYRIGHT 1999 ACS

AN 1996:661119 CAPLUS

DN 125:294771

TI Viral and plasmid vectors encoding mouse mammary tumor virus Naf repressor

or Sag antigen for control of viral infections or lymphocyte gene therapy

IN Guenzburg, Walter H.; Salmons, Brian

PA Bavarian Nordic Research Institute A/s, Den.;

GSF-Forschungszentrum fuer

Umwelt und Gesundheit GmbH

SO PCT Int. Appl. 44 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN CNT 1

PATENT NO. KIND DATE APPLICATION NO.
DATE

PI WO 9628564 AI 19960919 WO 1996-EP1002

19960308

W: AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS,

JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LV, MD, MG,

MK, MN, MW,

MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TJ, TM, TR, TT,

UA, UG,

US, UZ

RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI,

FR, GB, GR,

IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML,

MR, NE, SN, TD, TG

AU 9651040 AI 19961002 AU 1996-51040 19960308

EP 817859 AI 19980114 EP 1996-907399 19960308

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,

IE, SI, LT, LV, FI

JP 11508441 T2 19990727 JP 1996-527260 19960308

PRAI DK 1995-243 19950309

WO 1996-EP1002 19960308

AB The invention refers to novel recombinant vectors useful for gene therapy

of viral infections and of diseases assocd. with B and T cells. The present invention relates, furthermore, to novel usages of the two products of the open reading frame of mouse mammary tumor virus, i.e. the

transcription repressor Naf and superantigen Sag. Procon (promoter conversion) viral vectors may be used to deliver the Naf gene to target

cells and thereby repress expression from heterologous viral

promoters.

Alternatively, the vector may deliver the Sag gene, and, optionally,

a B-

or T-cell-specific therapeutic gene. This will stimulate expansion

of B

and T cells expressing the therapeutic gene.

L17 ANSWER 6 OF 10 CAPLUS COPYRIGHT 1999 ACS

AN 1996:661120 CAPLUS

DN 125:294754

TI Vectors carrying therapeutic genes encoding antimicrobial peptides for

gene therapy

IN Guenzburg, Walter H.; Winder, David; Saller, Robert Michael

PA Bavarian Nordic, Den.; GSF-Forschungszentrum fuer Umwelt

und Gesundheit

GmbH

SO PCT Int. Appl. 54 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN CNT 1

PATENT NO. KIND DATE APPLICATION NO.
DATE

PI WO 9628563 AI 19960919 WO 1996-EP1001

19960308

W: AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS,

JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LV, MD, MG,

MK, MN, MW,

MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TJ, TM, TR, TT,

UA, UG,

US, UZ

RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI,

FR, GB, GR,

IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML,

MR, NE, SN, TD, TG

AU 9651039 AI 19961002 AU 1996-51039 19960308

EP 817858 AI 19980114 EP 1996-907398 19960308

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,

IE, SI, LT, LV, FI

JP 11503305 T2 19990326 JP 1996-527259 19960308

PRAI DK 1995-243 19950309

WO 1996-EP1001 19960308

AB The present invention relates to retroviral vectors carrying sequences

encoding naturally occurring, antimicrobial peptides or derivs. thereof

for the treatment of mammalian tumors, viral infections such as

HIV

infection and bacterial and fungal infections. In particular the

present

invention relates to retroviral vectors which undergo promoter

conversion

(Procon vectors) carrying such sequences. Since these vectors also

carry

tumor or virus specific regulatory elements, the therapeutic

antimicrobial

peptide will be delivered and expressed only in relevant, affected

cells

and not in innocent bystander cells. The U3 region of murine

leukemia

virus-derived vector BAG was replaced with a mouse mammary

tumor virus U3

region without the inverted repeats but contg. the promoter, a

region

conferring responsiveness to glucocorticoid hormones, and a region

contg.

an element directing expression to the mammary gland. A

proprocropin A

gene was inserted next to the promoter to produce vector

p125 CercA. EJ

cells expressing the luciferase gene fused to the HIV LTR and the

Tat gene

displayed luciferase expression. When these recombinant cells

were

infected with p125 CercA there was little luciferase expression.

L17 ANSWER 7 OF 10 CAPLUS COPYRIGHT 1999 ACS

AN 1996:346074 CAPLUS

DN 125:2982

TI Safe, non-self-inactivating retroviral expression vectors using non-LTR

promoters for gene therapy

IN Guenzburg, Walter Henry; Saller, Robert Michael

PA GSF-Forschungszentrum fuer Umwelt und Gesundheit GmbH,

Germany

SO PCT Int. Appl., 40 pp.

CODEN: PIXXD2

DT Patent
LA English
FAN/CNT 1

PATENT NO. KIND DATE APPLICATION NO.
DATE

PI WO 9607748 A1 19960314 WO 1995-EP3445
19950901

W: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU,
IS, JP, KE,
KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW,
MX, NO, NZ, PL,
RO, RU, SD, SG, SI, SK, TJ, TT, UA, UG, US, UZ, VN
RW: KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR,
GB, GR, IE, IT,
LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN,
ML, MR, NE,
SN, TD, TG

CA 2198210 AA 19960314 CA 1995-2198210 19950901
AU 9535201 A1 19960327 AU 1995-35201 19950901
AU 688590 B2 19980312
EP 779929 A1 19970625 EP 1995-931969 19950901

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU,
MC, NL, PT, SE

CN 1159210 A 19970910 CN 1995-194903 19950901
BR 9508664 A 19980106 BR 1995-8664 19950901
HU 76974 A2 19980128 HU 1997-1764 19950901
JP 10507628 T2 19980728 JP 1995-509186 19950901
NO 9700902 A 19970424 NO 1997-902 19970227
FI 9700892 A 19970228 FI 1997-892 19970228
PRAI DK 1994-1017 19940902
WO 1995-EP3445 19950901

AB Retroviral expression vectors for gene therapy with a reduced risk of recombination with helper virus genomes and that use non-retroviral promoters in place of the LTRs are described. These vectors are constructed with non-retroviral regulatory elements in place of the 3'-LTR. After infection, the 3'-LTR region is duplicated and transposed to the 5'-LTR leading to elimination of the viral LTR and expression of the gene from the 5'-LTR. The construct replaces the U3 region of the 3'-LTR with the foreign promoter. This vector will not self-inactivate over time. Vectors using the promoter of the ***whey*** gene or of the mouse mammary tumor virus beta-galactosidase gene are demonstrated.

L17 ANSWER 8 OF 10 CAPLUS COPYRIGHT 1999 ACS
AN 1995:994888 CAPLUS
DN 124:47632
TI Manufacture and secretion into milk of oligosaccharides and

glycoconjugates typical of human milk by mammary gland-specific expression of the human genes for oligosaccharide biosynthetic enzymes

IN Prieto, Pedro Antonio, Smith, David Fletcher, Cummings, Richard Dale;
Kopchik, John Joseph; Mukerji, Pradip; Moremen, Kelley Wilson; Pierce,
James Michael
PA Abbott Laboratories, USA
SO PCT Int. Appl., 51 pp.
CODEN: PIXXD2

DT Patent
LA English
FAN/CNT 1

PATENT NO. KIND DATE APPLICATION NO.
DATE

PI WO 9524495 A1 19950914 WO 1995-US967
19950124

W: AU, CA, FI, JP, MX, NL, NZ
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC,
NL, PT, SE

US 5750176 A 19980512 US 1994-208889 19940309
CA 2184686 AA 19950914 CA 1995-2184686 19950124
AU 9516901 A1 19950925 AU 1995-16901 19950124
AU 697523 B2 19981008
EP 750673 A1 19970102 EP 1995-908663 19950124

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL,
PT, SE

JP 09510094 T2 19971014 JP 1995-523443 19950124
PRAI US 1994-208889 19940309
WO 1995-US967 19950124

AB Methods for genetic engineering of the milk of a non-human mammal is characterized so that it contains heterologous components produced as the secondary gene products of a heterologous gene integrated into the genome of the transgenic non-human mammal are described. The heterologous gene encodes an enzyme such as a human enzyme selected from the group consisting of glycosyltransferases, phosphorylases, hydroxylases, peptidases and sulfotransferases. Esp. useful in the practice of the invention are human glycosyltransferases. The desired heterologous components include oligosaccharides, glycoconjugates. The oligosaccharides and glycoconjugates may be isolated from the milk of the transgenic mammals and used in the prepn. of pharmaceuticals, diagnostic kits, nutritional products and the like. The whole milk may also be used to formulate nutritional products that provide special advantages. The transgenic milk may also be used in the prodn. of specialized enteral

nutritional products. Methods for transforming oocytes and screening preimplantation embryos for the presence of the transforming DNA are described. The cloning and expression of a cDNA for a human fucosyltransferase in transgenic mice using the ***whey*** acid*** protein*** gene promoter to direct mammary gland-specific expression in mice is demonstrated.

L17 ANSWER 9 OF 10 CAPLUS COPYRIGHT 1999 ACS
AN 1995:780439 CAPLUS
DN 123:190527

TI Transgenic production of antibodies in milk and usefulness for diagnostics, therapy, or industry

IN Meade, Harry; DiIulio, Paul; Pollock, Daniel
PA Genzyme Transgenics Corp., USA
SO PCT Int. Appl., 24 pp.
CODEN: PIXXD2

DT Patent
LA English
FAN/CNT 1

PATENT NO. KIND DATE APPLICATION NO.
DATE

PI WO 9517085 A1 19950629 WO 1994-US14795
19941220

W: AU, CA, JP, NZ
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC,
NL, PT, SE

US 5827690 A 19981027 US 1993-170579 19931220
CA 2178941 AA 19950629 CA 1994-2178941 19941220
AU 9515172 A1 19950710 AU 1995-15172 19941220
AU 688845 B2 19980319
EP 741515 A1 19961113 EP 1995-906691 19941220

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC,
NL, PT, SE

JP 09506779 T2 19970708 JP 1994-517602 19941220
US 5849992 A 19981215 US 1995-410887 19950327
AU 9873079 A1 19980820 AU 1998-73079 19980619
PRAI US 1993-170579 19931220
WO 1994-US14795 19941220

AB A method for the prodn. of monoclonal antibodies in mammal's milk, through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.

L17 ANSWER 10 OF 10 CAPLUS COPYRIGHT 1999 ACS
AN 1993:230822 CAPLUS
DN 118:230822

TI Protein composition of rhesus monkey milk: comparison to human milk
AU Kunz, Clemens; Lommerdal, Bo
CS Dep. Nutr., Univ. California, Davis, CA, 95616, USA
SO Comp. Biochem. Physiol., A: Comp. Physiol. (1993), 104A(4), 793-7
CODEN: CBPAB5; ISSN: 0300-9629

DT Journal
LA English
AB Proteins in human milk and Rhesus monkey milk were compared by FPLC gel filtration and anion-exchange chromatog., SDS-PAGE, nitrogen and protein
determ. Mature Rhesus milk is higher in protein concn. (15-20 mg/mL) than human milk but
Rhesus milk but 25-30% in human milk. Secretory IgA, lactoferrin, serum albumin, alpha-lactalbumin and lysozyme are present in Rhesus milk, but at a lower concn. than in human milk. The casein subunit pattern is more complex in Rhesus milk compared to human milk. The ratio of whey proteins to casein is similar in both milks (approx 60/40). A protein with a Mr of 21,600 is a major component in monkey whey but is not found in human milk.

=> s 14

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L18 2 L4

=> dup rem I18

PROCESSING COMPLETED FOR L18
L19 2 DUP REM L18 (0 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 2 ANSWERS -
CONTINUE? Y(N):y

L19 ANSWER 1 OF 2 CAPLUS COPYRIGHT 1999 ACS
AN 1991:625431 CAPLUS
DN 115:225431

TI Production of heterologous polypeptides by recombinant cattle and transgenic methods
IN Heyneker, Herbert L.; Deboer, Herman A.; Strijker, Rein; Plantenburg, Gerard; Lee, Sang He
PA Genpharm International, Inc., USA
SO PCT Int. Appl., 121 pp.
CODEN: PIXXD2

DT Patent
LA English
FAN.CNT 2

PATENT NO. KIND DATE APPLICATION NO.
DATE
PI WO 9108216 A1 19910613 WO 1990-US6874 19901130
W: AU, BR, CA, FI, JP, KR, LK, MC, NO, SU
RW: AT, BE, BF, BJ, CF, CG, CH, CM, DE, DK, ES, FR, GA, GB, GR, IT, LU, ML, MR, NL, SE, SN, TD, TG
CA 2075206 AA 19910602 CA 1990-2075206 19901130
AU 9169608 A1 19910626 AU 1991-69608 19901130
AU 656720 B2 19950216
EP 502976 A1 19920916 EP 1991-901026 19901130
EP 502976 B1 19960703
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE
AT 140027 E 19960715 AT 1991-901026 19901130
EP 737746 A2 19961016 EP 1995-203326 19901130
EP 737746 A3 19961023
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE
ES 2090299 T3 19961016 ES 1991-901026 19901130
RU 2095414 C1 19971110 RU 1990-5052392 19901130
CN 1053446 A 19910731 CN 1990-109733 19901201
NO 9202996 A 19920729 NO 1992-2996 19920729
FI 9203485 A 19920731 FI 1992-3485 19920731
US 5633076 A 19970527 US 1993-154019 19931116
US 5741957 A 19980421 US 1995-461333 19950605
PRAI US 1989-444745 19891201
US 1990-619131 19901127
EP 1991-901026 19901130
WO 1990-US6874 19901130
US 1992-898956 19920615
US 1993-77788 19930615
US 1993-154019 19931116
AB A method for prep. transgenic cows which secrete recombinant proteins into their milk is described. The gene to be expressed in mammary tissue is fused to a mammary tissue-specific promoter, e.g. that of the casein gene, a signal sequence, and a 3' flanking sequence functional in cattle. The chimeric gene is first methylated, e.g. by cloning it in a prokaryotic host. Fertilized oocytes are then transformed with this gene, and the fertilized oocytes are cultured to the preimplantation embryo stage. A cell is removed from the embryo to test for the presence of the desired gene: the chimeric methylated gene is resistant to restriction endonuclease cleavage. The hemiembryo remaining after removing the cell is cloned to prep. multiple embryos which are implanted into a cow to produce transgenic offspring. The milk from the transgenic cows can be used in food formulations, esp. infant formulas. A chimeric gene comprising human lactoferrin cDNA flanked by bovine alpha-SI-***casein*** promoter*** and signal sequence and 3' prep. Transgenic cows secreting lactoferrin into their milk were produced using this gene according to the above procedure.

L19 ANSWER 2 OF 2 CAPLUS COPYRIGHT 1999 ACS
AN 1989:451714 CAPLUS
DN 111:51714

TI Manufacture of recombinant proteins by secretion into milk of transgenic mammals
IN Meade, Harry; Longberg, Nils
PA Biogen N. V., Neth.
SO PCT Int. Appl., 20 pp.
CODEN: PIXXD2

DT Patent
LA English
FAN.CNT 1

PATENT NO. KIND DATE APPLICATION NO.
DATE
PI WO 8810118 A1 19881229 WO 1988-US2134 19880623
W: JP
RW: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE
US 4873316 A 19891010 US 1987-65994 19870623
EP 347431 A1 19891227 EP 1988-906454 19880623
EP 347431 B1 19951004
R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE
JP 02500798 T2 19900322 JP 1988-505800 19880623
AT 128625 E 19951015 AT 1988-906454 19880623
US 5750172 A 19980512 US 1995-460959 19950605
PRAI US 1987-65994 19870623
WO 1988-US2134 19880623
US 1989-332293 19890331
US 1993-109865 19930820
US 1994-322984 19941014
AB A method for producing desired proteins by producing transgenic mammals which secrete the protein into the milk is described. A section of the bovine alpha-S-1 casein gene contg. the promoter and signal sequence was cloned. This DNA sequence was ligated to tissue-type plasminogen activator (tPA) cDNA via DNA contg. RNA processing splice sites (which allow the casein signal sequence RNA to be spliced to the tPA-encoding RNA) to prep. pCASI151. Preimplantation fertilized mice embryos were microinjected with this (linearized) DNA and then implanted in pseudopregnant female mice. Of 262 embryos injected and implanted, 23 live pups were born, 5 of which contained the desired DNA sequences. Male

G0 mice were bred with females. Females of the G1 progeny which contained the IPA sequence produced 0.2-0.5 .mu.g tPA/mL milk.

=> s immunoglobulin# and beta-lactoglobulin/ab,bi

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L20 909 IMMUNOGLOBULIN# AND BETA-LACTOGLOBULIN/AB,BI

=> s 120 and promoter#/ab,bi

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L21 2 L20 AND PROMOTER#/AB,BI

=> dup rem I21

PROCESSING COMPLETED FOR L21
L22 2 DUP REM L21 (0 DUPLICATES REMOVED)

=> d 1 - bib ab

YOU HAVE REQUESTED DATA FROM 2 ANSWERS -
CONTINUE? Y(N)?

L22 ANSWER 1 OF 2 CAPLUS COPYRIGHT 1999 ACS
AN 1999:77669 CAPLUS
DN 130:134970
T1 Heterologous expression of proteins by rescued vector comprising an intron
IN Colman, Alan; Garner, Ian; Dalrymple, Michael Alexander
PA PPL Therapeutics (Scotland) Limited, UK
SO PCT Int. Appl., 58 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN,CNT 1
PATENT NO. KIND DATE APPLICATION NO.
DATE

PI WO 9903981 A1 19990128 WO 1998-CB2130 19980717
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,

UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
AU 9884502 A1 19990210 AU 1998-84502 19980717
PRAJ GB 1997-15064 19970717
WO 1998-CB2130 19980717
AB A nucleic acid expression construct comprising: (a) a ***promoter***;
(b) an intron whose natural position is within the 5'-untranslated region of a gene from which it is derived; (c) a coding sequence; and (d) a 3'-flanking sequence wherein the intron (b) is not derived from the same gene as that from which either the ***promoter*** (a) or the protein-coding sequence (c) is derived and processes, vectors, hosts and uses involving such a construct to obtain inter alia an increase in the level of expression of the coding sequence. To take advantage of "rescue" technol., a pMAD vector was constructed from the ovine .
beta .-
lactoglobulin gene for the cloning of cDNAs. The vector contains the same 5' and 3' flanking sequences preseng in the . ***beta*** .-
lactoglobulin gene which itself always gives rise to high expression in transgenic mice, but lacks all coding sequences and introns of the intact genes. Cloning of cDNAs in the unique EcoRV site between 5' and 3' flanking sequences results in constructs suitable for expression by the "rescue" approach. If instead of the . ***beta*** .- ***lactoglobulin*** first intron, an intron whose natural position is within the 5'-untranslated region of its gene is used (e.g., the bovine or ovine .beta.-casein intron 1 or the cardiac actin intron 1), good expression results. The expression constructs are exemplified for the expression of protein C, antibody fragments, IgG, adhesion mol., and collagen.

L22 ANSWER 2 OF 2 BIOSIS COPYRIGHT 1999 BIOSIS
AN 1999:188239 BIOSIS
DN PREV199900188239
T1 Variable immune response against a developmentally regulated self-antigen.
AU Steinhoff, U. (1); Maloy, K. J.; Burkhardt, C.; Clark, A. J.; Ruelicke, T.; Hengartner, H.; Zinkernagel, R. M.

CS (1) Max-Planck Institute for Infection Biology, Monbijoustr. 2, 10117, Berlin Germany
SO Journal of Autoimmunity, (Feb., 1999) Vol. 12, No. 1, pp. 27-34. ISSN: 0896-8411.
DT Article
LA English
AB We studied the reactivity of T and B cells against a soluble form of the glycoprotein of vesicular stomatitis virus (VSV-G) which was expressed in a transgenic mouse (line 23) under the control of the hormone regulated ***beta*** . ***lactoglobulin*** ***promoter*** . Transgenic mice expressed VSV-G in the thymus, spleen, mammary gland and lung. VSV-G transcripts in the thymus varied with age, i.e., expression was high early in life and decreased with age. VSV-G transgenic mice immunized with recombinant vaccinia virus expressing VSV-G exhibited normal VSV-G-specific IgM-levels, but a 30-fold reduction in IgG response, indicating functional VSV-G-specific B cell activity but impaired T helper cell responses. Interestingly, VSV-G-specific T helper cell activity was reduced only early (4-10 weeks) and late in life (>40 weeks) but was normal in between. Double transgenic mice expressing VSV-G and a VSV-G-specific TCR (line 23 X 7) demonstrated that TCR transgenic CD4+ T cells were partially deleted in earlylife, but then gradually repopulated the periphery and remained constant. These findings suggest that in line 23 two different mechanisms regulated levels of the immune response: clonal reduction/deletion of VSV-G-specific T cells during early life followed by peripheral anergy at a later stage.

=> s I11

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L23 1345 L11

=> s I23 and promoter#/ab,bi

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE
L24 24 L23 AND PROMOTER#/AB,BI
=> dup rem l24

PROCESSING COMPLETED FOR L24
L25 13 DUP REM L24 (11 DUPLICATES REMOVED)
=> d l- bib ab

YOU HAVE REQUESTED DATA FROM 13 ANSWERS -
CONTINUE? Y(N)y

L25 ANSWER 1 OF 13 CAPLUS COPYRIGHT 1999 ACS
AN 1999:77669 CAPLUS
DN 130:134970

TI Heterologous expression of proteins by rescued vector comprising an intron
IN Colman, Alan; Garner, Ian; Dalrymple, Michael Alexander
PA PPL Therapeutics (Scotland) Limited, UK
SO PCT Int. Appl., 58 pp.
CODEN: PXXXD2

DT Patent
LA English
FAN/CNT 1

PATENT NO. KIND DATE APPLICATION NO.
DATE -----

PI WO 9903981 A1 19990128 WO 1998-GB2130
19980717

W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN,
CU, CZ, DE,
DK, EE, ES, FI, GB, GE, GH, GM, GR, HU, ID, IL, IS, JP,
KE, KG,
KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK,
MN, MW, MX,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT,
UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD,
RU, TJ, TM
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH,
CY, DE, DK, ES,
FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF,
CG, CI,
CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
AU 9884502 A1 19990210 AU 1998-84502 19980717
PRAI GB 1997-15064 19970717
WO 1998-GB2130 19980717

AB A nucleic acid expression construct comprising: (a) a
promoter;
(b) an intron whose natural position is within the 5'-untranslated
region
of a gene from which it is derived; (c) a coding sequence; and (d) a
3'-flanking sequence wherein the intron (b) is not derived from the
same
gene as that from which either the ***promoter*** (a) or the

protein-coding sequence (c) is derived and processes, vectors, hosts
and
uses involving such a construct to obtain inter alia an increase in the
level of expression of the coding sequence. To take advantage of
"rescue"
technol., a pMAD vector was constructed from the ovine
beta-lactoglobulin gene for the cloning of cDNAs. The vector
contains
the same 5' and 3' flanking sequences preseng in the
beta-lactoglobulin
gene which itself always gives rise to high level expression in
transgenic
mice, but lacks all coding sequences and introns of the intact genes.
Cloning of cDNAs in the unique EcoRV site between 5' and 3'
flanking
sequences results in constructs suitable for expression by the
"rescue"
approach. If instead of the beta-lactoglobulin first intron, an intron
whose natural position is within the 5'-untranslated region of its
gene is
used (e.g. the bovine or ovine beta- ***casein*** intron 1 or
the
cardiac actin intron 1), good expression results. The expression
constructs are exemplified for the expression of protein C, antibody
fragments, IgG, adhesion mol., and collagen.

L25 ANSWER 2 OF 13 EMBASE COPYRIGHT 1999 ELSEVIER
SC1 B.V.
AN 97257933 EMBASE
DN 1997257933

TI Distinct functional properties of I kappa B alpha. and
I kappa B beta..
AU Tran K.; Merika M.; Thanos D.
CS D. Thanos, DBMB, Columbia University, 630 West 168th St,
New York, NY
10032, United States
SO Molecular and Cellular Biology, (1997) 17/9 (5386-5399).
Refs: 65
ISSN: 0270-7306 CODEN: MCEBD4

CY United States
DT Journal: Article
FS 004 Microbiology
LA English
SL English
AB The biological activity of the transcription factor NF-kappa B is
controlled mainly by the I kappa B alpha. and I kappa B beta.
proteins,
which restrict NF-kappa B to the cytoplasm and inhibit its DNA
binding
activity. Here, we carried out e experiments to determine and
compare the
mechanisms by which I kappa B alpha. and I kappa B beta. inhibit
NF-kappa B-dependent transcriptional activation. First, we found
that in
vivo I kappa B alpha. is a stronger inhibitor of NF-kappa B than is
I kappa B beta. This difference is directly correlated with their
abilities to inhibit NF-kappa B binding to DNA in vitro and in

vivo.
Moreover, I kappa B.alpha., but not I kappa B.beta., can remove
NF-kappa.B from functional preinitiation complexes in vitro
transcription
experiments. Second, we showed that both I kappa.Bs function in
vivo not
only in the cytoplasm but also in the nucleus, where they inhibit
NF-kappa.B binding to DNA. Third, the inhibitory activity of
I.kappa.B.beta., but not that of I.kappa.B.alpha., is facilitated by
phosphorylation of the C-terminal PEST sequence by
casein kinase
II and/or by the interaction of NF-.kappa.B with high-mobility
group
protein I (HMG I) on selected ***promoters***. The
unphosphorylated
form of I.kappa.B.beta. forms stable ternary complexes with
NF-.kappa.B on
the DNA either in vitro or in vivo. These experiments suggest that
I.kappa.B.alpha. works as a postinduction repressor of NF-.kappa.B
independently of HMG I, whereas I.kappa.B.beta. functions
preferentially
in ***promoters*** regulated by the NF-.kappa.B/HMG I
complexes.

L25 ANSWER 3 OF 13 MEDLINE
I AN 96355620 MEDLINE
DN 96355620

TI Stat6 and Jak1 are common elements in platelet-derived growth
factor and
interleukin-4 signal transduction pathways in NIH 3T3 fibroblasts.
AU Patel B K; Wang L M; Lee C C; Taylor W G; Pierce J H;
LaRoche W J
CS Laboratory of Cellular and Molecular Biology, National Cancer
Institute,
Bethesda, Maryland 20892, USA.
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1996 Sep 6) 271
(36) 22175-82.
Journal code: HIV. ISSN: 0021-9258.

CY United States
DT Journal: Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; Cancer Journals
EM 199612

AB Both platelet-derived growth factor (PDGF) and interleukin-4
(IL-4) play
major roles in cell proliferation, differentiation, chemotaxis, and
other
functional responses. Here, we demonstrate that Stat6, previously
shown to
be activated by only IL-4 and IL-3, becomes activated after PDGF
stimulation of NIH 3T3 fibroblasts. PDGF BB, and to a lesser
extent PDGF
AA, rapidly induced DNA binding activity from NIH 3T3 cell
lysates
utilizing the ***immunoglobulin*** heavy chain germ line
epsilon

promoter (lepsilon) that specifically binds to Stat6 in an electrophoretic mobility shift assay. DNA binding activity could be detected within 5 min and reached maximum levels at approximately 20 min in parental NIH 3T3 cells. An identical mobility shift and time course of PDGF-mediated lepsilon binding activity was more pronounced in lysates of NIH 3T3 transfectants overexpressing human Stat6 (NIH 3T3-Stat6). The observed radiolabeled lepsilon mobility shift was competed by lepsilon as well as by the beta- ***casein*** gene ***promoter*** but not by the interferon-alpha-stimulated response element or the interferon-gamma response region of the guanylate-binding protein gene. A Stat6-specific polyclonal antiserum also supershifted the PDGF-induced lepsilon mobility shift. After PDGF BB treatment, a 100-kDa tyrosine phosphorylated species was detected in anti-Stat6 immunoprecipitates. Cycloheximide had little effect on Stat6 tyrosine phosphorylation. In addition to Stat6, Stat5a, and Stat5b, PDGF BB also induced Jak1 tyrosine phosphorylation suggesting a potential pathway for Stat activation. Strikingly, the concurrent addition of IL-4 enhanced PDGF BB-induced lepsilon binding activity. Jak1 tyrosine phosphorylation, and [3H]thymidine incorporation. These results provide evidence that Stat6 and Jak1 are common elements in PDGF and IL-4 signaling pathways and suggest that IL-4 could play a role in potentiating certain known biological responses.

L25 ANSWER 4 OF 13 EMBASE COPYRIGHT 1999 ELSEVIER SCI. B.V.
AN 96096378 EMBASE
DN 1996096378
TI Phosphorylation of I.kappa.B.alpha. in the C-terminal PEST domain by ***casein*** kinase II affects intrinsic protein stability.
AU Lin R.; Beauparlant P.; Makris C.; Meloche S.; Hiscott J.
CS Lady Davis Medical Research Inst., 3755 Cote Ste. Catherine, Montreal, Que.
H3T 1E2, Canada
SO Molecular and Cellular Biology, (1996) 16/4 (1401-1409).
ISSN: 0270-7306 CODEN: MCEBD4
CY United States
DT Journal; Article
FS 029 Clinical Biochemistry
LA English
SL English

AB The NF-kappa.B/Rel transcription factors participate in the activation of immune system regulatory genes and viral early genes including the human immunodeficiency virus type 1 long terminal repeat. NF-kappa.B/Rel proteins are coupled to inhibitory molecules, collectively termed I.kappa.B, which are responsible for cytoplasmic retention of NF-kappa.B. Cell activation leads to the phosphorylation and degradation of I.kappa.B.alpha., permitting NF-kappa.B/Rel translocation to the nucleus and target gene activation. To further characterize the signaling events that contribute to I.kappa.B.alpha. phosphorylation, a kinase activity was isolated from Jurkat T cells that specifically interacted with I.kappa.B.alpha. in an affinity chromatography step and phosphorylated I.kappa.B.alpha. with high specificity in vitro. By using an in-gel kinase assay with recombinant I.kappa.B.alpha. as substrate, two forms of the kinase (43 and 38 kDa) were identified. Biochemical criteria and immunological cross-reactivity identified the kinase activity as the alpha catalytic subunit of ***casein*** kinase II (CKII). Deletion mutants of I.kappa.B.alpha. (DELTA.1 to DELTA.4) localized phosphorylation to the C-terminal PEST domain of I.kappa.B.alpha.. Point mutation of residues T-291, S-283, and T-299 dramatically reduced phosphorylation of I.kappa.B.alpha. by the kinase in vitro. NIH-3T3 cells that stably expressed wild-type I.kappa.B.alpha. (wtl.kappa.B), double-point-mutated I.kappa.B.alpha. (T291A, S283A), or triple-point-mutated I.kappa.B.alpha. (T291A, S283A, T299A) under the control of the tetracycline-responsive ***promoter*** were generated. Constitutive phosphorylation of the triple point mutant was eliminated in vivo, although tumor necrosis factor-inducible I.kappa.B.alpha. degradation was unaffected. In cell lines and in transiently transfected cells, mutation of the CKII sites in I.kappa.B.alpha. resulted in a protein with increased intrinsic stability. Together with results demonstrating a role for N-terminal sites in inducer-mediated phosphorylation and degradation of I.kappa.B.alpha., these studies indicate that CKII sites in the C-terminal PEST domain are important for constitutive phosphorylation and intrinsic stability of I.kappa.B.alpha..

L25 ANSWER 5 OF 13 CAPLUS COPYRIGHT 1999 ACS

AN 1996:609130 CAPLUS
DN 125:273244
TI IL-10 induces DNA binding activity of three STAT proteins (Stat1, Stat3, and Stat5) and their distinct combinatorial assembly in the ***promoters*** of selected genes
AU Wehinger, Jens; Gouilleux, Fabrice; Groner, Bernd; Finke, Juergen; Mertelsmann, Roland; Weber-Nordt, Renate Maria
CS University of Freiburg Medical Center, Department of Hematology and Oncology, Hugstetter Str. 55, Freiburg, 79106, Germany
SO FEBS Lett (1996), 394(3), 365-370
CODEN: FEBLAL; ISSN: 0014-5793
DT Journal
LA English
AB Interaction of IL-10 with its receptor leads to the activation of STAT transcription factors. Herein the authors report the IL-10 dependent simultaneous activation of 3 STAT transcription factors: Stat1, Stat3, and Stat5. Upon IL-10 treatment multiple Stat proteins become simultaneously activated, and bind to different ***promoters*** with equal kinetics but form distinct homo- and heterodimeric transcriptionally active complexes depending on the STAT-consensus elements of a selected gene ***promoter***. Upon IL-10 treatment Stat1, 3, and 5 bind to the GRR of the Fe-gamma.RI gene, activated Stat1 and 3 bind to the SIE sequence of the c-fos ***promoter*** and transcriptionally active Stat5 assembles at the PRL-STAT consensus sequence of the .beta.- ***casein*** gene. Thus, functionally relevant STAT dimerization is influenced by the activated cytokine receptor as well as the specific STAT consensus sequence present in a specific gene ***promoter***.

L25 ANSWER 6 OF 13 MEDLINE
2
AN 96358438 MEDLINE
DN 96358438
TI Regulation of somatostatin gene transcription by cyclic adenosine monophosphate.
AU Montminy M; Brindle P; Arias J; Ferri K; Armstrong R
CS Clayton Foundation Laboratories for Peptide Biology, Salk Institute, La Jolla, CA 92037, USA.
NC GM37828 (NIGMS)
SO METABOLISM: CLINICAL AND EXPERIMENTAL, (1996 Aug) 45 (8 Suppl 1) 4-7.
Ref: 12
Journal code: MUM. ISSN: 0026-0495.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)

General Review; (REVIEW)
(REVIEW, TUTORIAL)

LA English
FS Priority Journals

EW 199702
EW 19970204

AB Cyclic adenosine monophosphate (cAMP) stimulates transcription of

somatostatin and other target genes with burst-attenuation kinetics.

The

kinetics of protein kinase (PK-A)-dependent cAMP response

element binding

protein (CREB) phosphorylation closely parallel the changes in transcription of cAMP-responsive genes by run-on assay. Nuclear translocation of PK-A, visualized by microinjection of fluorescently labeled PK-A holoenzyme, appears to represent the rate-limiting step in

CREB phosphorylation and transcriptional activation. We and others have

recently characterized a CREB-binding protein (CBP), which specifically recognizes sequences within the Ser133 phosphorylated form of

CREB. CBP

does not regulate the DNA binding, dimerization, or nuclear targeting

properties of CREB, but binds selectively to the kinase-inducible

60 amino

acid trans-activation domain (KID) of CREB, critical for

PK-A-inducible

transcription. We developed an antiserum directed against amino

acid

634-648 within the CREB-binding domain of CBP. We detected a

265-kd

polypeptide by Western blot as predicted from the cDNA, which

coincided

with the predominant phospho-CREB-binding activity in HeLa

nuclear

extracts by "Far Western" blot assay. An identical

phospho-CREB-binding

activity was also found in NIH-3T3 cells. This

phospho-CREB-binding

protein appeared to be specific for Ser133-phosphorylated CREB,

because no

such band was detected with CREB labeled to the same specific

activity at

a nonregulatory phosphoacceptor site (Ser156) by ***casein***

kinase

effect on this response. We can now begin reconstituting

PK-A-dependent

transcription in vitro, using well-characterized proteins such as

CREB,

TAF 110, and CBP. The assembly of such factors on

cAMP-regulated

promoters like somatostatin may enable responsiveness to

a variety

of hormonal stimuli that employ cAMP as their second messenger.

L25 ANSWER 7 OF 13 EMBASE COPYRIGHT 1999 ELSEVIER

SC1 B.V DUPLICATE 3

AN 93305660 EMBASE

DN 1993305660

T1 Interleukin 4-inducible phosphorylation of HMG-I(Y) is inhibited

by

rapamycin.

AU Wang D.-Z.; Ray P.; Boothby M.

CS Dept. of Microbiology/Immunology, Vanderbilt Univ. School of

Medicine, Nashville, TN 37232-2363, United States

SO Journal of Biological Chemistry, (1995) 270/39 (22924-22932).

ISSN: 0021-9258 CODEN: JBCHA3

CY United States

DT Journal; Article

FS 026 Immunology, Serology and Transplantation

029 Clinical Biochemistry

LA English

SL English

AB The non-histone chromosomal protein HMG-I(Y) participates in

repression of

transcription directed by a ***promoter*** which confers

interleukin 4

(IL-4)-inducible activation in transfected B cell lines. Metabolic

labeling, phosphoamino acid analyses, and in vitro phosphorylation

studies

DNA. These data further suggest a novel mechanism in which

phosphorylation

triggered by IL-4 or other cytokines could regulate the effects of

HMG-I(Y) on gene transcription

L25 ANSWER 8 OF 13 CAPLUS COPYRIGHT 1999 ACS

AN 1994:210040 CAPLUS

DN 120:210040

T1 Manufacture of foreign proteins in cattle and their accumulation in

milk

IN Deboer, Herman A.; Strijker, Rein; Heyneker, Herbert L.;

Platenburg.

Gerard; Lee, Sang He; Pieper, Frank

PA Genpharm International, Inc., USA

SO PCT Int. Appl., 178 pp.

CODEN: PIXXDZ

DT Patent

LA English

FAN CNT 2

PATENT NO. KIND DATE APPLICATION NO.

DATE

PI WO 9325567 AI 19931223 WO 1993-US5724

19930615

W: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB,

HU, JP, KP,

KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO,

RU, SD, SE,

SK, UA, US, VN

RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC,

NL, PT, SE,

BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD,

TG

AU 9345373 AI 19940104 AU 1993-45373 19930615

EP 652889 AI 19950517 EP 1993-915365 19930615

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LJ, LU, MC,

NL, PT, SE

JP 08504562 T2 19960521 JP 1993-501794 19930615

US 5633076 A 19970527 US 1993-154019 19931116

US 5741957 A 19980421 US 1995-461333 19950605

PRAI US 1992-898956 19920615

US 1989-444745 19891201

US 1990-619131 19901127

US 1993-777788 19930615

WO 1993-US5724 19930615

US 1993-154019 19931116

AB Expression cassettes using regulatory regions functional in

mammary

secretory cells are used to direct the synthesis of proteins in cattle

mammary glands with subsequent accumulation of the protein in

milk

Methods for prep. transgenic cattle by transformation and

implantation of

embryos are described. The transgene is methylated and introduced

into

fertilized oocytes and the oocytes cultured to the pre-implantation

embryo

stage. Cells are then removed from the pre-implantation embryos and the DNA digested with a restriction endonuclease capable of cleaving the methylated transgene but not the unmethylated form. The development of expression vectors using the 5'- and 3'-flanking sequences of the cattle .alpha.S1 ***casein*** gene was demonstrated and an expression cassette for a human lactoferrin cDNA constructed. Transgenic calves carrying the expression cassette were obtained and one showing normal sperm and a lack of mosaicism was obtained.

L25 ANSWER 9 OF 13 MEDLINE DUPLICATE
4

AN 92260636 MEDLINE
DN 92260636
TI Receptor properties of two varicella-zoster virus glycoproteins, gpIV and gpI

AU Litwin V; Jackson W; Grose C
CS Department of Microbiology, University of Iowa College of Medicine, Iowa City 52242..

NC A122795 (NIAID)
SO JOURNAL OF VIROLOGY, (1992 Jun) 66 (6) 3643-51.
Journal code: KCV. ISSN: 0022-538X.

CY United States
DT Journal; Article; (JOURNAL ARTICLE)

FS Priority Journals; Cancer Journals
EM 199208

AB The varicella-zoster virus (VZV) genome contains 70 reading frames (ORF), 5 of which encode the glycoproteins gpI, gpII, gpIV, and gpV. ORF 67 and 68 lie adjacent to each other in the unique short region of the VZV

genome and code for gpIV and gpI, respectively. These two genes, which are contained within the HindIII C fragment of the VZV genome, were subcloned in the correct orientation downstream from the ***promoter*** of the eukaryotic expression vectors pCMV5 and pBI. After transfection, 5 to 20% of the Cos cells bound antibody specific for the given glycoprotein. In this study, it was shown that only the cells transfected with the gpI construct bound to the Fc fragment of human ***immunoglobulin*** G. Neither the transfected gpIV gene product nor the vector only bound to the Fc fragment. Thus, VZV gpI is confirmed to be

the VZV-encoded Fc-binding glycoprotein. Like the wild-type form of gpI expressed in VZV-infected cells, gpI precipitated from transfected cells contained both N-linked and O-linked glycans and was heavily sialated. In addition, the transfected gpI gene product was phosphorylated both in cell culture and in protein kinase assays by mammalian ***casein*** kinases

I and II. Extensive computer-assisted analyses of the VZV gpI sequence, as well as those of alphaherpesviral homolog glycoproteins, disclosed properties similar to those of other cell surface receptors; these included (i) exocytosmic regions rich in cysteine residues, (ii) membrane-proximal regions with potential O-linked glycosylation sites, and (iii) cytoplasmic domains with consensus phosphorylation sites.

L25 ANSWER 10 OF 13 BIOSIS COPYRIGHT 1999 BIOSIS
AN 1992:468244 BIOSIS

DN BR43:89594

TI THE TRANSCRIPTION FACTOR CF1 REGULATES THE C-MYC THE IGH AND THE BETA

CASEIN ***PROMOTERS***
AU MEIER V; SCHMITT-NEY M; GRONER B

CS FRIEDRICH MIESCHER INST., CH-4002, BASEL.
SO 24TH ANNUAL MEETING OF THE SWISS SOCIETIES FOR EXPERIMENTAL BIOLOGY (USGE/USSEB), BASEL, SWITZERLAND, MARCH 19-20, 1992. EXPERIENTIA (BASEL).

(1992) 48 (ABSTR.), A51.

CODEN: EXPEAM. ISSN: 0014-4754.

DT Conference

FS BR; OLD

LA English

L25 ANSWER 11 OF 13 CAPLUS COPYRIGHT 1999 ACS
AN 1991:625431 CAPLUS

DN 115:225431

TI Production of heterologous polypeptides by recombinant cattle and transgenic methods

IN Heyneker, Herbert L.; Deboer, Herman A.; Strijker, Rein; Plantenburg,

Gerard; Lee, Sang He
PA Genpharm International, Inc., USA

SO PCT Int. Appl., 121 pp.
CODEN: PIXXD2

DT Patent

LA English

FAN CNT 2

PATENT NO. KIND DATE APPLICATION NO.

DATE

PI WO 9108216 AI 19910613 WO 1990-US6874

19901130

W: AU, BR, CA, FI, JP, KR, LK, MC, NO, SU
RW: AT, BE, BF, BJ, CF, CG, CH, CM, DE, DK, ES, FR, GA, GB, GR, IT,

LU, ML, MR, NL, SE, SN, TD, TG

CA 2073206 AA 19910602 CA 1990-2075206 19901130
AU 9169608 AI 19910626 AU 1991-69608 19901130

AU 656720 B2 19950216

EP 502976 A1 19920916 EP 1991-901026 19901130

EP 502976 B1 19960703

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE
AT 140027 E 19960715 AT 1991-901026 19901130

EP 737746 A2 19961016 EP 1995-203326 19901130

EP 737746 A3 19961023

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE
ES 2090299 T3 19961016 ES 1991-901026 19901130

RU 2095414 C1 19971110 RU 1990-5052392 19901130

CN 1053446 A 19910731 CN 1990-109733 19901201

NO 9202996 A 19920729 NO 1992-2996 19920729

FI 9203485 A 19920731 FI 1992-3485 19920731

US 5633076 A 19970527 US 1993-154019 19931116

US 5741957 A 19980421 US 1995-461333 19950605

PRAI US 1989-444745 19891201

US 1990-619131 19901127

EP 1991-901026 19901130

WO 1990-US6874 19901130

US 1992-898956 19920615

US 1993-77788 19930615

US 1993-154019 19931116

AB A method for prep. transgenic cows which secrete recombinant proteins

into their milk is described. The gene to be expressed in mammary tissue

is fused to a mammary tissue-specific ***promoter***, e.g. that of the

casein gene, a signal sequence, and a 3' flanking sequence

functional in cattle. The chimeric gene is first methylated, e.g. by cloning it in a prokaryotic host. Fertilized oocytes are then transformed

with this gene, and the fertilized oocytes are cultured to the preimplantation embryo stage. A cell is removed from the embryo to test

for the presence of the desired gene: the chimeric methylated gene is resistant to restriction endonuclease cleavage. The hemembryo remaining

after removing the cell is cloned to prep. multiple embryos which are implanted into a cow to produce transgenic offspring. The milk from the

transgenic cows can be used in food formulations, esp. infant formulas. A

chimeric gene comprising human lactoferrin cDNA flanked by bovine

.alpha.S1- ***casein*** ***promoter*** and signal sequence and 3'

regions was prepd. Transgenic cows secreting lactoferrin into their

milk were produced using this gene according to the above procedure.

L25 ANSWER 12 OF 13 MEDLINE DUPLICATE
5
AN 92107650 MEDLINE
DN 92107650
TI Upstream box/TATA box order is the major determinant of the direction of transcription.
AU Xu L C; Thali M; Schaffner W
CS Institut für Molekularbiologie II, Universität Zürich, Switzerland.
SO NUCLEIC ACIDS RESEARCH, (1991 Dec 25) 19 (24) 6699-704.
Journal code: O8L. ISSN: 0305-1048.
CY ENGLAND: United Kingdom
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; Cancer Journals
EM 199204
AB Mammalian gene ***promoters*** for transcription by RNA polymerase II are typically organized in the following order: upstream sequence motif(s)/TATA box/initiation site. Here we report studies in which the order, orientation and DNA sequences of these three elements are varied to determine how these affect polarity of transcription. We have constructed ***promoters*** with an 'octamer' upstream sequence ATTTCAT (or its complement ATGCAAAAT) in combination with several different TATA boxes and initiation (cap) sites, and tested these ***promoters*** in transfection experiments with cultured cells. TATA boxes derived from the adenovirus major late ***promoter*** (TATAAAA), ***immunoglobulin*** kappa light chain (TTATATA) and heavy chain (TAAATATA) ***promoter*** functioned equally well or even better when inverted. Only the TATA box (CATAAAA) was poorly active when inverted. In addition, a symmetrical TATA box (TATATATA) derived from a ***casein*** gene was very active. Our results suggest that the asymmetry of most TATA boxes (consensus TATAAAA) is not a primary determinant of the polarity of transcription. We also found that the initiation (cap) site, which usually consists of an adenine embedded in a pyrimidine-rich region (PyPyCAPyPyPyPy), was permissive towards sequence alterations; even a randomly composed sequence worked well. However, an inverted, hence

purine-rich, cap site reduced transcript levels to 1/7th, as did an oligo G sequence. Irrespective of the presence of a cap site, the configuration: 'TATA box/octamer' yielded a strong leftward, rather than rightward transcription. From this, we conclude that the polarity of transcription is primarily determined by the linear order of an upstream sequence relative to a TATA box, rather than by the individual orientations of either of these two elements.

L25 ANSWER 13 OF 13 CAPLUS COPYRIGHT 1999 ACS
AN 1989-451714 CAPLUS
DN 111-51714
TI Manufacture of recombinant proteins by secretion into milk of transgenic mammals
IN Meade, Harry; Longberg, Nils
PA Biogen N. V., Neth.
SO PCT Int. Appl., 20 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN CNT 1
PATENT NO. KIND DATE APPLICATION NO.
DATE
PI WO 8810118 AI 19881229 WO 1988-US2134
19880623
W: JP
RW: AT, BE, CH, DE, FR, GB, IT, LU, NL, SE
US 4873316 A 19891010 US 1987-65994 19870623
EP 347431 A1 19891227 EP 1988-906454 19880623
EP 347431 B1 19951004
R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE
JP 02500798 T2 19900322 JP 1988-505800 19880623
AT 128625 E 19951015 AT 1988-906454 19880623
US 5750172 A 19980512 US 1995-460959 19950605
PRAI US 1987-65994 19870623
WO 1988-US2134 19880623
US 1989-332293 19890331
US 1993-109865 19930820
US 1994-322984 19941014
AB A method for producing desired proteins by producing transgenic mammals which secrete the protein into the milk is described. A section of the bovine alpha. S-I ***casein*** gene contg. the ***promoter*** and signal sequence was cloned. This DNA sequence was ligated to tissue-type plasminogen activator (tPA) cDNA via DNA contg. RNA processing splice sites (which allow the ***casein*** signal sequence RNA to be spliced to the tPA-encoding RNA) to prep. pCASI151. Preimplantation

fertilized mice embryos were microinjected with this (linearized) DNA and then implanted in pseudopregnant female mice. Of 262 embryos injected and implanted, 23 live pups were born, 5 of which contained the desired DNA sequences. Male G0 mice were bred with females. Females of the G1 progeny which contained the tPA sequence produced 0.2-0.5 .mu.g tPA/mL milk.
=> s 18
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L26 652 L8
=> s 126 and promoter/ab,bi
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L27 0 L26 AND PROMOTER/AB,BI
=> s 126 and (construct# or vector# or transgen?)/ab,bi
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L28 0 L26 AND (CONSTRUCT# OR VECTOR# OR TRANSGEN?)/AB,BI
=> d his
(FILE 'HOME' ENTERED AT 16:55:03 ON 04 AUG 1999)
FILE 'MEDLINE' ENTERED AT 16:55:09 ON 04 AUG 1999
L1 4 S IMMUNOGLOBULIN AND WHEY ACIDIC PROTEIN/AB,BI
L2 4 S IMMUNOGLOBULIN# AND WHEY ACIDIC PROTEIN/AB,BI
L3 0 S IMMUNOGLOBULIN# AND BETA-LACTOGLOBULIN PROMOTER/AB,BI
L4 0 S IMMUNOGLOBULIN# AND CASEIN PROMOTER/AB,BI
L5 0 S IMMUNOGLOBULIN# AND BETA-CASEIN PROMOTER/AB,BI
L6 0 S IMMUNOGLOBULIN# AND KAPPA-CASEIN PROMOTER/AB,BI
L7 0 S IMMUNOGLOBULIN# AND LACTALBUMIN

PROMOTER/AB,BI
L8 96 S IMMUNOGLOBULIN# AND
LACTALBUMIN/AB,BI
L9 0 S L8 AND (CONSTRUCT# OR VECTOR#)/AB,BI
L10 0 S L8 AND TRANSGEN#/AB,BI
L11 190 S IMMUNOGLOBULIN# AND CASEIN/AB,BI
L12 8 S L11 AND (CONSTRUCT# OR VECTOR#)/AB,BI

FILE MEDLINE, EMBASE, BIOSIS, INPADOC, CAPLUS'
ENTERED AT 17:05:15 ON 04
AUG 1999

L13 13 S L1
L14 6 DUP REM L13 (7 DUPLICATES REMOVED)
L15 23 S L2
L16 10 S L2 NOT L1
L17 10 DUP REM L16 (0 DUPLICATES REMOVED)
L18 2 S L4
L19 2 DUP REM L18 (0 DUPLICATES REMOVED)
L20 909 S IMMUNOGLOBULIN# AND
BETA-LACTOGLOBULIN/AB,BI
L21 2 S L20 AND PROMOTER#/AB,BI
L22 2 DUP REM L21 (0 DUPLICATES REMOVED)
L23 1345 S L11
L24 24 S L23 AND PROMOTER#/AB,BI
L25 13 DUP REM L24 (11 DUPLICATES REMOVED)
L26 652 S L8
L27 0 S L26 AND PROMOTER/AB,BI
L28 0 S L26 AND (CONSTRUCT# OR VECTOR# OR
TRANSGEN#)/AB,BI

=> e meade harry/au

E1 2 MEADE H W/AU
E2 1 MEADE HARLAN DONNELLY/AU
E3 42 --> MEADE HARRY/AU
E4 28 MEADE HARRY M/AU
E5 2 MEADE HASH T J/AU
E6 2 MEADE HASH TARA J/AU
E7 2 MEADE HAZEL/AU
E8 5 MEADE HAZEL W/AU
E9 1 MEADE HAZEL WHITTIG/AU
E10 2 MEADE HOWARD M/AU
E11 3 MEADE HUERTA P/AU
E12 2 MEADE HUERTA PATRICIA/AU

=> s e3-e4

L29 70 ("MEADE HARRY"/AU OR "MEADE HARRY
M"/AU)

=> s i29 and immunoglobulin# and promoter#/ab,bi

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L30 2 L29 AND IMMUNOGLOBULIN# AND

PROMOTER#/AB,BI

=> dup rem i30

PROCESSING COMPLETED FOR L30

L31 2 DUP REM L30 (0 DUPLICATES REMOVED)

=> d i- bib ab

YOU HAVE REQUESTED DATA FROM 2 ANSWERS -
CONTINUE? Y/(N)?

L31 ANSWER 1 OF 2 CAPLUS COPYRIGHT 1999 ACS

AN 1995:780439 CAPLUS

DN 123:190527

TI Transgenic production of antibodies in milk and usefulness for
diagnostics, therapy, or industry

IN ***Meade, Harry*** ; Dittullo, Paul; Pollock, Daniel

PA Genzyme Transgenics Corp., USA

SO PCT Int. Appl., 24 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN CNT 1

PATENT NO. KIND DATE APPLICATION NO.

DATE

PI WO 9517085 AI 19950629 WO 1994-US14795
19941220

W: AU, CA, JP, NZ

RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE

US 5827690 A 19981027 US 1993-170579 19931220

CA 2178941 AA 19950629 CA 1994-2178941 19941220

AU 9515172 AI 19950710 AU 1995-15172 19941220

AU 688845 B2 19980319

EP 741515 AI 19961113 EP 1995-906691 19941220

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC,
NL, PT, SE

JP 09506779 T2 19970708 JP 1994-517602 19941220

US 5849992 A 19981215 US 1995-410887 19950327

AU 9873079 AI 19980820 AU 1998-73079 19980619

PRAI US 1993-170579 19931220

WO 1994-US14795 19941220

AB A method for the prodn. of monoclonal antibodies in mammal's
milk, through
the creation of transgenic animals that selectively express foreign
antibody genes in mammary epithelial cells.

L31 ANSWER 2 OF 2 CAPLUS COPYRIGHT 1999 ACS

AN 1989:451714 CAPLUS

DN 111:51714

TI Manufacture of recombinant proteins by secretion into milk of
transgenic
mammals

IN ***Meade, Harry*** ; Longberg, Nils

PA Biogen N. V., Neth.
SO PCT Int. Appl., 20 pp.
CODEN: PIXXD2

DT Patent

LA English

FAN CNT 1

PATENT NO. KIND DATE APPLICATION NO.
DATE

PI WO 8810118 AI 19881229 WO 1988-US2134
19880623

W: JP

RW: AT, BE, CH, DE, FR, GB, IT, LU, NL, SE

US 4873316 A 19891010 US 1987-65994 19870623

EP 347431 AI 19891227 EP 1988-906454 19880623

EP 347431 BI 19951004

R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

JP 02500798 T2 19900322 JP 1988-505800 19880623

AT 128625 E 19951015 AT 1988-906454 19880623

US 5750172 A 19980512 US 1995-460959 19950605

PRAI US 1987-65994 19870623

WO 1988-US2134 19880623

US 1989-332293 19890331

US 1993-109865 19930820

US 1994-322984 19941014

AB A method for producing desired proteins by producing transgenic
mammals

the which secrete the protein into the milk is described. A section of
the bovine .alpha. S-1 casein gene contg. the ***promoter*** and
signal

sequence was cloned. This DNA sequence was ligated to
tissue-type

plasminogen activator (tPA) cDNA via DNA contg. RNA

processing splice

sites (which allow the casein signal sequence RNA to be spliced to
the

tPA-encoding RNA) to prep. pCASI151. Preimplantation
fertilized mice

embryos were microinjected with this (linearized) DNA and then
implanted

in pseudopregnant female mice. Of 262 embryos injected and
implanted, 23

live pups were born, 5 of which contained the desired DNA
sequences. Male

G0 mice were bred with females. Females of the G1 progeny
which contained

the tPA sequence produced 0.2-0.5 .mu.g tPA/mL milk.

=> d 2 kwic

L31 ANSWER 2 OF 2 CAPLUS COPYRIGHT 1999 ACS

IN ***Meade, Harry*** ; Longberg, Nils

AB . . . which secrete the protein into the milk is described. A
section

of the bovine .alpha. S-1 casein gene contg. the ***promoter***

and
 signal sequence was cloned. This DNA sequence was ligated to
 tissue-type
 plasmidogen activator (IPA) cDNA via DNA contg. RNA. . .
 IT Casens, biological studies
 RL: BIOL (Biological study)
 (gene for, ***promoter*** and signal sequence of, secretion
 of recombinant protein into milk of transgenic mammals in relation
 to)
 IT ***Immunoglobulins***
 RL: PROC (Process)
 (manuf. of, with transgenic mammals, by secretion into milk)
 IT Molecular cloning
 (of chimeric casein ***promoter*** and signal
 sequence-plasminogen
 activator gene, for expression in transgenic mice and sheep)
 IT Gene and Genetic element
 (chimeric, ***promoter*** and signal sequence of
 milk-specific
 protein-contg., in secretion of recombinant protein into milk of
 transgenic mammals)
 IT Plasmid and Episome
 (pCAS1151, chimeric casein ***promoter*** and signal
 sequence-plasminogen activator gene on, expression in transgenic
 mice
 and sheep of)
 => e ditullio paul/au
 E1 1 DITULLIO NW/AU
 E2 19 DITULLIO P/AU
 E3 32 --> DITULLIO PAUL/AU
 E4 1 DITULLIO R/AU
 E5 1 DITULLIO R M/AU
 E6 4 DITULLIO ROBERT J/AU
 E7 1 DITULLIO S/AU
 E8 3 DITULLIO S A/AU
 E9 9 DITULLIO V/AU
 E10 9 DITULLIO VENANZIO/AU
 E11 1 DITULLIO VENENZIO/AU
 E12 1 DITULLIO NICHOLAS W/AU
 => s e2-e3
 L32 51 ("DITULLIO P"/AU OR "DITULLIO PAUL"/AU)
 => s l32 and immunoglob? and promoter#/ab,bi
 'AB' IS NOT A VALID FIELD CODE
 'AB' IS NOT A VALID FIELD CODE
 'AB' IS NOT A VALID FIELD CODE
 'AB' IS NOT A VALID FIELD CODE
 L33 1 L32 AND IMMUNOGLOB? AND PROMOTER#/AB,BI
 => d bib ab

L33 ANSWER I OF I CAPLUS COPYRIGHT 1999 ACS
 AN 1995:780439 CAPLUS
 DN 123:190527
 TI Transgenic production of antibodies in milk and usefulness for
 diagnostics, therapy, or industry
 IN Meade, Harry; ***Ditullio, Paul*** ; Pollock, Daniel
 PA Genzyme Transgenics Corp., USA
 SO PCT Int. Appl., 24 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN,CNT 1
 PATENT NO. KIND DATE APPLICATION NO.
 DATE

 PI WO 9517085 A1 19950629 WO 1994-US14795
 19941220
 W: AU, CA, JP, NZ
 RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC,
 NL, PT, SE
 US 5827690 A 19981027 US 1993-170579 19931220
 CA 2178941 AA 19950629 CA 1994-2178941 19941220
 AU 9515172 A1 19950710 AU 1995-15172 19941220
 AU 688845 B2 19980319
 EP 741515 A1 19961113 EP 1995-906691 19941220
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC,
 NL, PT, SE
 JP 09506779 T2 19970708 JP 1994-517602 19941220
 US 5849992 A 19981215 US 1995-410887 19950327
 AU 9873079 A1 19980820 AU 1998-73079 19980619
 PRAI US 1993-170579 19931220
 WO 1994-US14795 19941220
 AB A method for the prodn. of monoclonal antibodies in mammal's
 milk, through
 the creation of transgenic animals that selectively express foreign
 antibody genes in mammary epithelial cells.
 => e pollock daniel/au
 E1 4 POLLOCK D S/AU
 E2 2 POLLOCK D W/AU
 E3 16 --> POLLOCK DANIEL/AU
 E4 11 POLLOCK DANIEL A/AU
 E5 7 POLLOCK DANIEL D/AU
 E6 12 POLLOCK DARREN A/AU
 E7 1 POLLOCK DARREN R/AU
 E8 1 POLLOCK DAVE/AU
 E9 19 POLLOCK DAVID/AU
 E10 3 POLLOCK DAVID A/AU
 E11 1 POLLOCK DAVID ALEXANDER/AU
 E12 5 POLLOCK DAVID B/AU
 => s e3-e5
 L34 34 ("POLLOCK DANIEL"/AU OR "POLLOCK DANIEL
 A"/AU OR "POLLOCK DANIEL

D"/AU)
 => s l34 and immunoglob? and promoter#/ab,bi
 'AB' IS NOT A VALID FIELD CODE
 'AB' IS NOT A VALID FIELD CODE
 'AB' IS NOT A VALID FIELD CODE
 'AB' IS NOT A VALID FIELD CODE
 L35 1 L34 AND IMMUNOGLOB? AND PROMOTER#/AB,BI
 => d
 L35 ANSWER I OF I CAPLUS COPYRIGHT 1999 ACS
 AN 1995:780439 CAPLUS
 DN 123:190527
 TI Transgenic production of antibodies in milk and usefulness for
 diagnostics, therapy, or industry
 IN Meade, Harry; Ditullio, Paul; ***Pollock, Daniel***
 PA Genzyme Transgenics Corp., USA
 SO PCT Int. Appl., 24 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN,CNT 1
 PATENT NO. KIND DATE APPLICATION NO.
 DATE

 PI WO 9517085 A1 19950629 WO 1994-US14795
 19941220
 W: AU, CA, JP, NZ
 RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC,
 NL, PT, SE
 US 5827690 A 19981027 US 1993-170579 19931220
 CA 2178941 AA 19950629 CA 1994-2178941 19941220
 AU 9515172 A1 19950710 AU 1995-15172 19941220
 AU 688845 B2 19980319
 EP 741515 A1 19961113 EP 1995-906691 19941220
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC,
 NL, PT, SE
 JP 09506779 T2 19970708 JP 1994-517602 19941220
 US 5849992 A 19981215 US 1995-410887 19950327
 AU 9873079 A1 19980820 AU 1998-73079 19980619
 PRAI US 1993-170579 19931220
 WO 1994-US14795 19941220
 (FILE 'HOME' ENTERED AT 16:55:03 ON 04 AUG 1999)
 FILE 'MEDLINE' ENTERED AT 16:55:09 ON 04 AUG 1999
 L1 4 S IMMUNOGLOBULIN AND WHEY ACIDIC
 PROTEIN/AB,BI
 L2 4 S IMMUNOGLOBULIN# AND WHEY ACIDIC
 PROTEIN/AB,BI
 L3 0 S IMMUNOGLOBULIN# AND

THIS PAGE BLANK (USPTO)

BETA-LACTOGLOBULIN PROMOTER/AB,BI
L4 0 S IMMUNOGLOBULIN# AND CASEIN
PROMOTER/AB,BI
L5 0 S IMMUNOGLOBULIN# AND BETA-CASEIN
PROMOTER/AB,BI
L6 0 S IMMUNOGLOBULIN# AND KAPPA-CASEIN
PROMOTER/AB,BI
L7 0 S IMMUNOGLOBULIN# AND LACTALBUMIN
PROMOTER/AB,BI
L8 96 S IMMUNOGLOBULIN# AND
LACTALBUMIN/AB,BI
L9 0 S L8 AND (CONSTRUCT# OR VECTOR#)/AB,BI
L10 0 S L8 AND TRANSGEN#/AB,BI
L11 190 S IMMUNOGLOBULIN# AND CASEIN/AB,BI
L12 8 S L11 AND (CONSTRUCT# OR VECTOR#)/AB,BI

FILE 'MEDLINE, EMBASE, BIOSIS, INPADOC, CAPLUS'
ENTERED AT 17:05:15 ON 04
AUG 1999

L13 13 S L1
L14 6 DUP REM L13 (7 DUPLICATES REMOVED)
L15 23 S L2
L16 10 S L2 NOT L1
L17 10 DUP REM L16 (0 DUPLICATES REMOVED)
L18 2 S L4
L19 2 DUP REM L18 (0 DUPLICATES REMOVED)
L20 909 S IMMUNOGLOBULIN# AND
BETA-LACTOGLOBULIN/AB,BI
L21 2 S L20 AND PROMOTER#/AB,BI
L22 2 DUP REM L21 (0 DUPLICATES REMOVED)
L23 1345 S L11
L24 24 S L23 AND PROMOTER#/AB,BI
L25 13 DUP REM L24 (11 DUPLICATES REMOVED)
L26 652 S L8
L27 0 S L26 AND PROMOTER/AB,BI
L28 0 S L26 AND (CONSTRUCT# OR VECTOR# OR
TRANSGEN#)/AB,BI
E MEADE HARRY/AU
L29 70 S E3-E4
L30 2 S L29 AND IMMUNOGLOBULIN# AND
PROMOTER#/AB,BI
L31 2 DUP REM L30 (0 DUPLICATES REMOVED)
E DITULLIO PAUL/AU
L32 51 S E2-E3
L33 1 S L32 AND IMMUNOGLOB? AND
PROMOTER#/AB,BI
E POLLOCK DANIEL/AU
L34 34 S E3-E5
L35 1 S L34 AND IMMUNOGLOB? AND
PROMOTER#/AB,BI

=>

--Logging off of STN---

=>
Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY SESSION
FULL ESTIMATED COST 164.98 172.27

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)
SINCE FILE TOTAL
ENTRY SESSION
CA SUBSCRIBER PRICE -11.78 -11.78

STN INTERNATIONAL LOGOFF AT 17:15:07 ON 04 AUG 1999

3. 5,827,690, Oct. 27, 1998, Transgenic production of antibodies in milk: Harry Meade, et al., 800/7: 530/867 [IMAGE AVAILABLE]

US PAT NO: 5 852 224 [IMAGE AVAILABLE] LI: 1 of 5

said method comprising producing, by the method of claim 1, non-human animal milk containing at least 2 mg/ml of said α -lactalbumin and extracting said α -lactalbumin from said milk.

CLMS(3)

3. A transgenic non-human mammal whose somatic and germ cells contain a transgene construct, said transgene construct comprising, in the 5' to 3' direction and operatively linked:

(a) at least about 1.8 kb of 5'-flanking sequence from the human α - α -lactalbumin**promoter**;

(b) a DNA sequence encoding

(i) a signal sequence, and

(2) a α -lactalbumin selected from the group consisting of:

(i) human α -lactalbumin; and

(ii) a modified human α -lactalbumin having from one to four of the natural phenylalanine residues substituted by other amino acid residues;

(c) at least about 3 kb of 3'-flanking sequence from the human α -lactalbumin gene;

wherein said transgene construct is integrated into the genome of said mammal in such a way that said DNA sequence is expressed in the mammary gland of said mammal to produce α -lactalbumin in the milk of said mammal at a level of at least 2 mg/ml.

CLMS(4)

4. The transgenic non-human mammal of claim 3 wherein said mammal is a mouse.

CLMS(5)

5. The transgenic non-human mammal of claim 3 wherein said mammal is a cow.

US PAT NO: 5,589,604 [IMAGE AVAILABLE] L1: 4 of 5

CLAIMS:

CLMS(1)

What we claim is:

1. A transgenic non-human mammal that contains and expresses a human protein C DNA construct in the cells of its mammary gland, wherein the DNA construct consists of:

(a) a mammary gland promoter,

(b) a nucleotide sequence that encodes a signal peptide, wherein said signal peptide is effective in directing the secretion of an associated polypeptide into the milk of said transgenic non-human mammal, and wherein said signal peptide-encoding nucleotide sequence is operatively associated with said mammary gland promoter, and

(c) a nucleotide sequence encoding human protein C that is operatively

associated with said signal peptide-encoding nucleotide sequence, wherein human protein C is secreted into the milk of said transgenic non-human mammal, and when purified, said protein C has a specific activity more than about 80% of the specific activity of human protein C isolated from human plasma, as determined by an assay of protein serine protease activity or anticoagulant activity, and wherein said non-human mammal is selected from the group consisting of mouse, pig, sheep, goat and cattle.

CLMS(2)

2. The transgenic non-human mammal of claim 1, wherein said promoter is selected from the group consisting of a whey acidic protein promoter, a casein promoter, a α -lactalbumin**promoter** and a β -lactoglobulin promoter.

CLMS(3)

3. The transgenic non-human mammal of claim 2, wherein promoter is a whey acidic protein promoter or a β -lactoglobulin promoter.

CLMS(4)

4. The transgenic non-human mammal of claim 3, wherein said promoter is a whey acidic protein promoter.

CLMS(5)

5. The transgenic non-human mammal of claim 1, wherein said human protein C isolated from said transgenic non-human mammal has a specific activity that is about 80% to about 100% of the specific activity of human protein C isolated from human plasma.

CLMS(6)

6. The transgenic non-human mammal of claim 5, wherein said specific activity is determined by an activated partial thromboplastin clotting time assay.

CLMS(7)

7. The transgenic non-human mammal of claim 5, wherein said promoter is a whey acidic protein promoter or a β -lactoglobulin promoter.

CLMS(8)

8. A process for the production of protein C, comprising the steps of:

(a) providing a transgenic non-human mammal that contains and

expresses a human protein C DNA construct in the cells of its mammary gland, wherein the DNA construct consists of:

(i) a mammary gland promoter,

(ii) a nucleotide sequence that encodes a signal peptide, wherein said signal peptide is effective in directing the secretion of an associated polypeptide into the milk of said transgenic non-human mammal, and wherein said signal peptide-encoding nucleotide sequence

is operatively associated with said mammary gland promoter, and

(iii) a nucleotide sequence encoding human protein C that is operatively associated with said signal peptide-encoding nucleotide sequence,

wherein human protein C is secreted into the milk of said transgenic non-human mammal, and when purified, said protein C has a specific activity more than about 80% of the specific activity of human protein C isolated from human plasma, as determined by an assay of protein

C serine protease activity or anticoagulant activity, and wherein said non-human mammal is selected from the group consisting of mouse, pig, sheep, goat and cattle,

(b) producing milk from said transgenic non-human mammal,

(c) collecting said milk, and

(d) isolating said protein C from said milk.

CLMS(9)

9. The process of claim 8, wherein said promoter is selected from the group consisting of a whey acidic protein promoter, a casein promoter, a α -lactalbumin**promoter** and a β -lactoglobulin promoter.

CLMS(10)

10. The process of claim 9, wherein promoter is a whey acidic protein promoter or a β -lactoglobulin promoter.

CLMS(11)

11. The process of claim 10, wherein said promoter is a whey acidic protein promoter.

CLMS(12)

12. The transgenic non-human mammal of claim 8, wherein said human protein C isolated from said transgenic non-human mammal has a specific activity that is about 80% to about 100% of the specific activity of human protein C isolated from human plasma.

CLMS(13)

13. The transgenic non-human mammal of claim 12, wherein said specific

activity is determined by an activated partial thromboplastin clotting time assay.

CLMS(14)

14. The transgenic non-human mammal of claim 13, wherein said promoter is a whey acidic protein promoter or a β -lactoglobulin promoter.

CLMS(15)

15. A transgenic non-human mammal that contains and expresses a human protein C DNA construct in the cells of its mammary gland, wherein the DNA construct consists of:

- (a) a mammary gland promoter selected from the group consisting of a whey acidic protein promoter, a casein promoter, a β -lactalbumin** promoter, and a β -lactoglobulin promoter,
- (b) a nucleotide sequence that encodes a signal peptide, wherein said signal peptide is effective in directing the secretion of an associated polypeptide into the milk of said transgenic non-human mammal, and wherein said signal peptide-encoding nucleotide sequence is

operatively

associated with said mammary gland promoter; and

- (c) a nucleotide sequence encoding human protein C that is

operatively

associated with said signal peptide-encoding nucleotide sequence, wherein human protein C is secreted into the milk of said transgenic non-human mammal, and when purified, said protein C has a specific activity more than about 80% of the specific activity of human protein

C

isolated from human plasma, as determined by an assay of protein

serine protease activity or anticoagulant activity, and

wherein said non-human mammal is selected from the group

consisting of

mouse, pig, sheep, goat and cattle.

CLMS(16)

16. The transgenic non-human mammal of claim 15, wherein said human protein C isolated from said transgenic non-human mammal has a specific activity that is about 80% to about 100% of the specific activity of human protein C isolated from human plasma.

CLMS(17)

17. The transgenic non-human mammal of claim 16, wherein said specific activity is determined by an activated partial thromboplastin clotting time assay.

CLMS(18)

18. A process for the production of protein C, comprising the steps of: (a) providing a transgenic non-human mammal that contains and expresses

a human protein C DNA construct in the cells of its mammary gland, wherein the DNA construct consists of:

- (i) a mammary gland promoter selected from the group consisting of a whey acidic protein promoter, a casein promoter, a β -lactalbumin** promoter, and a β -lactoglobulin promoter,
- (ii) a nucleotide sequence that encodes a signal peptide, wherein said

signal peptide is effective in directing the secretion of an associated polypeptide into the milk of said transgenic non-human mammal, and wherein said signal peptide-encoding nucleotide sequence

is operatively associated with said mammary gland promoter, and

- (iii) a nucleotide sequence encoding human protein C that is operatively associated with said signal peptide-encoding nucleotide sequence,

wherein human protein C is secreted into the milk of said transgenic non-human mammal, and when purified, said protein C has a specific activity more than about 80% of the specific activity of human protein

C

isolated from human plasma, as determined by an assay of protein

serine protease activity or anticoagulant activity, and

wherein said non-human mammal is selected from the group

consisting of

mouse, pig, sheep, goat and cattle,

- (b) producing milk from said transgenic non-human mammal,
- (c) collecting said milk, and
- (d) isolating said protein C from said milk.

CLMS(19)

19. The transgenic non-human mammal of claim 18, wherein said

human protein C isolated from said transgenic non-human mammal has a

specific activity that is about 80% to about 100% of the specific activity of

human protein C isolated from human plasma.

CLMS(20)

20. The transgenic non-human mammal of claim 19, wherein said specific activity is determined by an activated partial thromboplastin clotting time assay.

US PAT NO: 5,523,226 [IMAGE AVAILABLE] LI: 5 of 5

CLAIMS:

CLMS(1)

What is claimed is:

- 1. A method of obtaining an embryonic stem cell for incorporation into a

swine embryo to form a chimeric swine, said method comprising: (a) introducing a cell from a culture made by:

- (i) culturing dissociated cells from a swine embryo in conditioned stem

cell medium in the presence or absence of a feeder layer, and (ii) subculturing the culture until a stable culture with morphological features and growth parameters characteristic of an embryonic stem cell culture is established, into a SCID mouse;

- (b) allowing a tumor to form in the mouse from the cell; and
- (c) obtaining an embryonic stem cell from a culture that is shown to be capable of producing a tumor in step b.

CLMS(2)

- 2. The method of claim 1, wherein the embryonic stem cell is characterized by an undifferentiated morphology indistinguishable from the morphology of a cell from the culture of step a of claim 1 from which a cell formed a tumor in step b of claim 1.

CLMS(3)

- 3. A method for determining the cell types in which a genetic complement

is expressed, said method comprising:

- (a) introducing a swine embryonic stem cell Which comprises the genetic complement into an immunocompromised mouse to produce a tumor;
- (b) placing the tumor in suitable conditions to allow the tumor to differentiate into a plurality of recognizable cell types and to express the genetic complement;
- (c) excising the tumor; and
- (d) analyzing the differentiated cell types to determine in which cell types the genetic complement is expressed.

CLMS(4)

- 4. An embryonic stem cell obtained from a culture that is capable of forming a tumor in a SCIDS mouse in accordance with the method of claim 1.

CLMS(5)

- 5. A culture initiated from an embryonic stem cell of claim 4.

=> d 4 kwic

US PAT NO: 5,589,604 [IMAGE AVAILABLE] LI: 4 of 5

CLAIMS:

CLMS(2)

2. . . . 1, wherein said promoter is selected from the group consisting of a whey acidic protein promoter, a casein promoter, a **lactalbumin** promoter** and a .beta.-lactoglobulin promoter.

CLAIMS:

CLMS(9)

9. . . . 8, wherein said promoter is selected from the group consisting of a whey acidic protein promoter, a casein promoter, a **lactalbumin** promoter** and a .beta.-lactoglobulin promoter.

CLAIMS:

CLMS(15)

15. . . . (a) a mammary gland promoter selected from the group consisting of a whey acidic protein promoter, a casein promoter, a **lactalbumin** promoter** and a .beta.-lactoglobulin promoter, (b) a nucleotide sequence that encodes a signal peptide, wherein said signal peptide is effective in directing. . . .

CLAIMS:

CLMS(18)

18. . . . (i) a mammary gland promoter selected from the group consisting of a whey acidic protein promoter, a casein promoter, a **lactalbumin** promoter** and a .beta.-lactoglobulin promoter, (ii) a nucleotide sequence that encodes a signal peptide, wherein said signal peptide is effective in directing. . . .

=> d 4 fro

US PAT NO: 5,589,604 [IMAGE AVAILABLE] L1: 4 of 5

DATE ISSUED: Dec. 31, 1996

TITLE: Expression of human protein C in mammary tissue of transgenic mammals

INVENTOR: William N. Drohan, Springfield, VA

Tracy D. Wilkins, Blacksburg, VA

William H. Velandier, Blacksburg, VA

John L. Johnson, Blacksburg, VA

ASSIGNEE: American Red Cross, Washington, DC (U.S. corp.)
Virginia Intellectual Property Division, Blacksburg, VA (U.S. corp.)

APPL-NO: 08/247,484

DATE FILED: May 23, 1994

REL-US-DATA: Continuation of Ser. No. 638,995, Jan. 11, 1991, abandoned.

INT-CL: [6] C12N 5/00; C12N 15/00; C12N 9/48; C12P 21/04

US-CL-ISSUED: 800/2: 435/69.6, 212

US-CL-CURRENT: 800/7; 435/69.6, 212; 800/14, 15, 16, 17, 18

SEARCH-FLD: 800/2; 435/172.3

REF-CITED:

U.S. PATENT DOCUMENTS

4,775,624 10/1988 Bang et al.

4,873,316 10/1989 Meade et al.

4,992,373 2/1991 Bang et al.

FOREIGN PATENT DOCUMENTS

0264166 4/1988 European Patent Office

0279582 8/1988 European Patent Office

WO88/00239 1/1988 World Intellectual Property Organization

WO88/01648 3/1988 World Intellectual Property Organization

WO90/05188 5/1990 World Intellectual Property Organization

OTHER PUBLICATIONS

Colpan et al (1984) "High Performance Liquid Chromatography of High-Molecular Weight Nucleic Acids. . ." 296, 339-353.

Clark et al (1987) Thbrech 5, 20-24.

Velandier et al (1992) Procd. Natl. Acad. Sci. 89, 12003-12007.

Hemninghausen (1990) Protein Exp. and Purif. 1, 3-8.

Grimnell et al (1990) Regulation and Prod. of Anticoag., 29-63.

Yan et al (1989) TIBS 14, 264-268.

Grimnell et al (1987) Bio/Technology 5, 1189-1192.

Pitrus et al (1988) Procd. Natl. Acad. Sci. 85, 5874-5878.

Hogan et al. (1986) Manipulating the Mouse Embryo, Cold Spring Harbor

Laboratory, Cold Spring Harbor, NY 153-203.

Denman et al. (Sep. 1991) Bio/Technology 9: 839-843. "Transgenic Expression of a Variant of Human tPA in Goat Milk: Purification and Characterization of the Recombinant Enzyme".

Ebert et al. (Sep. 1991) Bio/Technology 9: 835-838. "Transgenic Production of a Variant of Human tPA in Goat Milk: Generation of Transgenic Goats and Analysis of Expression".

Gordon et al. (Nov. 1987) Bio/Technology 5: 1183-1187, "Production of Human Tissue Plasminogen Activator in Transgenic Mouse Milk".

Krimpenfort et al. (Sep. 1991) Bio/Technology 9: 844-847, "Generation of Transgenic Dairy Cattle Using 'In Vitro' Embryo Production".

Walls et al. (1989) Gene 81: 139-149, "Amplification of Multicistronic Plasmids in the Human 293 Cell Line and Secretion of Correctly Processed Recombinant Human Protein C".

Wright et al. (Sep. 1991) Bio/Technology 9: 830, "High Level Expression of Active Human Alpha-1-Antitrypsin in the Milk of Transgenic Sheep".

Yan et al. (Jul. 1990) Bio/Technology: 8 655-661, "Characterization and Novel Purification of Recombinant Human Protein C from Three Mammalian Cell Lines".

ART-UNIT: 184

PRIM-EXMR: Deborah Crouch

LEGAL-REP: Foley & Lardner

ABSTRACT:

Recombinant protein C characterized by a high percentage of active protein can be obtained in the milk of transgenic mammals that incorporate DNAs according to the present invention. Transgenic mammals

of the present invention are produced by introducing into developing embryos DNA that encodes protein C, such that the DNA is stably incorporated in the DNA of germ line cells of the mature mammals and inherited in normal, mendelian fashion.

20 Claims, 5 Drawing Figures

=> d 3 fro

US PAT NO: 5,827,690 [IMAGE AVAILABLE] L1: 3 of 5

DATE ISSUED: Oct. 27, 1998

TITLE: Transgenic production of antibodies in milk

INVENTOR: Harry Meade, Newton, MA

Paul DiTullio, Framingham, MA

Daniel Pollock, Medway, MA

ASSIGNEE: Genzyme Transgenics Corporation, Framingham, MA (U.S. corp.)

APPL-NO: 08/170,579

DATE FILED: Dec. 20, 1993

INT-CL: [6] C12P 21/04; C12N 15/00

US-CL-ISSUED: 435/69.6, 172.3; 530/867; 800/2, DIG.1; 935/60

US-CL-CURRENT: 800/7; 530/867

SEARCH-FLD: 435/172.3, 69.1, 69.6; 530/867; 536/24.1; 800/2, DIG.1; 935/60

REF-CITED:

U.S. PATENT DOCUMENTS

4,816,397 3/1989 Boss et al. 435/68

4,816,567 3/1989 Cabilly et al. 530/387

4,873,316 10/1989 Meade et al. 530/412

5,322,775 6/1994 Clark et al. 435/69.1

FOREIGN PATENT DOCUMENTS

WO 90/04036 10/1989 World Intellectual Property Organization/21/8

WO 92/03918 8/1991 World Intellectual Property Organization

WO 93/12227 12/1992 World Intellectual Property Organization

OTHER PUBLICATIONS

Hemninghausen, L., Sippel, A. (1982) European Journal of Biochemistry 25:131-141.

Gordon, K., Lee, E., Vitale, J., Smith, A., Westphal, H., Hemninghausen, L. (1987) Bio/Technology 5:1183-1187.

Clark, A., Simons, P. Wilmut, I., Lathe, R. (1987) Bio/Technology

- 5:20-24.
- Brunt, J. (1988) *Bio/Technology* 6:1149-1154.
- Clark, A., Bestos, H., Bishop, J., Brown, P., Harris, S., Lathe, R., McClenaghan, M., Prowse, C., Simons, J., Whitelaw, C., Wilmut, L., Meade, H., Gates, L., Lacy, E., Lonberg, N., (1990) *Bio/Technology* 8:443-446.
- Buhler, Th. A., Bruyere, Th., Went, D., Stranzinger, G., Buiki, K., (1990) *Bio/Technology* 8:140-143.
- Weidte, U. H., Lenz, H., Brem, G. (1991) *Gene* 98 (2):185-91.
- DiTullio, P., Cheng, S., Marshall, J., Gregory, R., Ebert, K., Meade, H., (1992) *Bio/Technology* 10:74-77.
- Smith, A., (1992) *Bio/Technology* 10:74-77.
- Roberts, B., DiTullio, P., Vitale, J., Hehir, K., Gordon, K., (1992) *Gene* 121:255-262.
- Soulier et al., (1992) *FEBS Letters*, 297:13.
- Fiat, A. and Jolles, P., "Caseins of Various Origins and Biologically Active Casein Peptides and Oligosaccharides: Structural and Physiological Aspects," *Molecular and Cellular Biochemistry* vol. 87, 5-30 (1989).
- Larson, B., "Biosynthesis and Secretion of Milk Proteins: A Review," *Journal of Dairy Research*, vol. 46, 161-174 (1979).
- McKenzie, H. (editor), *Milk Proteins Chemistry and Molecular Biology*, Academic Press, vol. 1, pp. 3-15, and 26-29 (1970).
- McKenzie, H. (editor), *Milk Proteins Chemistry and Molecular Biology*, Academic Press, vol. 2, pp. 338-339 (1971).
- AH Horwitz et al (1988) *Proc Natl Acad Sci USA* 85: 8678-8682.
- M Sieger et al (1991) *Plant Science* 73: 181-190.
- A Hiatt et al (1989) *Nature* 342: 76-78.
- JP Bradshaw et al (1955) *Biochimica et Biophysica Acta* 847: 344-351.
- KM Ebert et al (1991) *Bio/Technology* 9: 835-838.
- R Bischoff et al (1992) *FEBS Letters* 305: 265-268.
- H Meade et al (1990) *Bio/Technology* 8: 443-446.
- M-A Persuy et al (1992) *European J Biochemistry* 205: 887-893.
- B Roberts et al (1992) *Gene* 121: 255-262.
- NM Greenberg et al (1991) *Proc Natl Acad Sci USA* 88: 8327-8331.
- JS Logan (1993) *Current Opinion in Biotechnology* 4: 591-595.
- JD Watson et al (1987) *Molecular Biology of the Gene* p. 313.
- VG Pursel et al (1993) *J Animal Sci* 71 (Supp 3):10-19.
- RJ Wall (1996) *Theriogenology* 45:57-68.
- LM Houdebine (1994) *J Biotechnol* 34:269-287.
- ART-UNIT: 184
- PRIM-EXMR: Bruce R. Campell
- LEGAL-REP: Lahive & Cockfield, LLP
- ABSTRACT:
A method for the production of monoclonal antibodies in mammal's milk, through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.
- 13 Claims, 4 Drawing Figures
- => d l fro
- US PAT NO: 5,852,224 [IMAGE AVAILABLE] L1: 1 of 5
- DATE ISSUED: Dec. 22, 1998
- TITLE: alpha-lactalbumin gene constructs
- INVENTOR: Julian David Cooper, Blacksburg, VA
- ASSIGNEE: Angelika Elisabeth Schnieke, Edinburgh, United Kingdom PPL Therapeutics (Scotland) Limited, Edinburgh, United Kingdom (foreign corp.)
- APPL-NO: 08/381,691
- DATE FILED: Jan. 31, 1995
- FRN-PRIOR: United Kingdom 9425326 Dec. 15, 1994
- INT-CL: [6] C12N 15/09; C12N 15/11; C12N 15/12; C12P 21/00
- US-CL-ISSUED: 800/2: 435/69.1, 71.1, 172.3; 935/34, 52, 70
- US-CL-CURRENT: 800/7: 435/69.1, 71.1; 800/13, 15, 18
- SEARCH-FLD: 800/2: 536/23.1, 24.1; 435/320.1, 240.2, 177.3, 69.1, 71.1; 350/365; 935/34, 52, 70
- REF-CITED:
- U.S. PATENT DOCUMENTS
- 4,293,583 10/1981 Farr et al. 426/657
- 5,530,177 6/1996 Bleck et al. 800/2
- FOREIGN PATENT DOCUMENTS
- 0 014 3621 8/1980 European Patent Office
- WO 88/01648 3/1988 World Intellectual Property Organization
- WO 93/25567 12/1993 World Intellectual Property Organization
- WO 95/02692 1/1995 World Intellectual Property Organization
- WO 95/18224 7/1995 World Intellectual Property Organization
- OTHER PUBLICATIONS
- Stacey et al., "Use of Double-Replacement Gene Targeting To Replace the Murine alpha-Lactalbumin Gene with Its Human counterpart in Embryonic Stem Cells and Mice", *Molecular and Cellular Biology*, 14(2):1009-1016 (Feb. 1994).
- Colman, A. (1996). *American Journal of Clinical Nutrition* 63, 6395-455.
- Burdon, T. et al (1991). *Mechanism of Development* 36, 57-74.
- Bleck, G. et al (1995). *International Dairy Journal* 5, 619-6320.
- Kappel, C. et al (1992). *Current Opinion: Biotechnology* 3, 548-553.
- Strojek, R. (1988). *Generic Engineering: Principles and Methods* v.10, pp. 221-246. Plenum Press.
- Krimpenfort, P. et al (1991). *Biotechnology* 9, 844-847.
- Hall, L. et al (1987) *Biochem J* 242, 735-742.
- Bleck, G.T. et al (1991) Symposium on Transgenes Development and Disease, Keystone Meeting, Tamaron, Colorado, J. Cell Biochem. Suppl. 0
- (15 Part A), 198.
- Adams, M et al (1993) *Nature Genetics* 4, 256-267.
- Maschio, et al., "Transgenic Mice Carrying The Guinea-Pig alpha-Lactalbumin Gene Transcribe Milk Protein Genes In Their Sebaceous Glands During Lactation", *Biochem. J.*, (1991) 275, 459-467.
- Soulier, et al., "Expression Analysis Of Ruminant alpha-Lactalbumin IN Transgenic Mice: Developmental Regulation And General Location Of Important cis-Regulatory Elements", *FEBS Letters*, vol. 297, No. 1,2, (Feb. 1991) pp. 13-18.
- Hochi, et al., "Secretion Of Bovine alpha-Lactalbumin Into The Milk Of Transgenic Rats", *Molecular Reproduction And Development* 33: 160-164 (1992).
- Vilotte, et al., "Efficient Tissue-Specific Expression Of Bovine alpha-Lactalbumin In Transgenic Mice", *Eur. J. Biochem.* 186, 43-48 (1989).
- Ninomiya, et al., "Functions Of Milk Protein Gene 5' Flanking Regions On Human Growth Hormone Gene", *Molecular Reproduction And Development* 3&2:276-283 (1994).
- F. Maynard, *Journal of Dairy Research*, vol. 59, No. 3, "Identification of a new molecular form of human alpha-lactalbumin" pp. 425-429 (1992).
- Hochi Shin-ichi et al., *Molecular Reproduction and Development*, vol. 33, "Secretion of bovine alpha-lactalbumin into the milk of transgenic rats", pp. 160-163, (1992).
- Jean-Luc Vilotte et al., *Biochimie*, vol. 69, No. 6/7, "Complete nucleotide sequence of bovine alpha-lactalbumin gene: comparison with its rat counterpart", pp. 609-620 (1987).
- ART-UNIT: 162
- PRIM-EXMR: Brian R. Stanton
- LEGAL-REP: Seidel, Gonda, Lavorgna & Monaco, PC
- ABSTRACT:
The present invention utilizes genetic engineering techniques to prepare non-human transgenic mammals that express human alpha-lactalbumin in their milk at a concentration of 2 mg/ml or greater. The invention also includes methods of preparing human alpha-lactalbumin in, for example, mice and cows. Also taught are methods for preparing human alpha-lactalbumin in which from one to four of its natural phenylalanine residues have been substituted by another amino acid.
- 5 Claims, 24 Drawing Figures
- => d his

(FILE 'USPAT' ENTERED AT 14:09:49 ON 05 AUG 1999)
L1 5 S LACTALBUMIN PROMOTER

=> log y

U.S. Patent & Trademark Office LOGOFF AT 14:15:04 ON 05 AUG
1999

SINCE FILE	TOTAL	ENTRY	SESSION
CA SUBSCRIBER PRICE		-2.14	-2.14

STN INTERNATIONAL LOGOFF AT 14:15:17 ON 05 AUG 1999

Connection closed by remote host

requirements are also taken into account, then
Tyr3/9-Tyr31-Tyr80 is preferred. Two DNA constructs, pHA-1 (contg. the human
.alpha.-lactalbumin gene and flanking regions) and pOBHA (contg.
the human
.alpha.-lactalbumin gene driven by the ovine .beta.-lactoglobulin
promoter), were injected into mouse embryos to give rise to
transgenic animals, expressing up to approx. 3 mg/mL
milk. The
bovine .alpha.-lactalbumin gene was also cloned and expressed in
mice.
PCR primer oligonucleotides were designed for site-specific
mutagenesis of
specific phenylalanine codons based on protein modeling,
nutritional
aspects, and on amino acid variants present in either native
.alpha.-lactalbumin or lysozyme genes from different species. The
addn.
of extra poly(Arg) or Arg/Lys residues at the C-terminus of the
.alpha.-lactalbumin assisted purifn. of endogenous protein.
Improved
expression of mutagenized bovine .alpha.-lactalbumin was
achieved by
control with the human .alpha.- ***lactalbumin***
promoter
L3 ANSWER 7 OF 10 CAPLUS COPYRIGHT 1999 ACS
AN 1993:227545 CAPLUS
DN 118:227545
TI Bovine .alpha.-lactalbumin gene promoter and its use in protein
manufacture
with ***transgenic*** female mammals
IN Bleck Gregory T.; Bremel, Robert D.
PA Wisconsin Milk Marketing Board, USA
SO PCT Int. Appl., 58 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN CNT 1
PATENT NO. KIND DATE APPLICATION NO.
DATE
PI WO 9304165 A1 19930304 WO 1992-US6549
19920806
W: AT, AU, BB, BG, BR, CA, CH, CS, DE, DK, ES, FI, GB,
HU, JP, KP,
KR, LK, LU, MG, MN, MW, NL, NO, PL, RO, RU, SD, SE
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC,
NL, SE, BF,
BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG
CA 2093659 AA 19930214 CA 1992-2093659 19920806
AU 9224119 A1 19930316 AU 1992-24119 19920806
AU 663101 B2 19950928
EP 555435 A1 19930818 EP 1992-916978 19920806
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL,
SE
JP 06502550 T2 19940324 JP 1993-504341 19920806

US 5530177 A 19960625 US 1993-71601 19930604
US 5850000 A 19981215 US 1996-621100 19960322
PRAI US 1991-744765 19910813
WO 1992-US6549 19920806
US 593-71601 19930604
AB A variant of the bovine .alpha.- ***lactalbumin***
promoter
which correlates with good milk prodn. is claimed.
Transgenic
female mice contg. the bovine .alpha.-lactalbumin gene contg. this
variation produced high levels of .alpha.-lactalbumin (>1 mg/mL)
in their
milk. Three other potentially significant variations in the steroid
response element and RNA polymerase binding region were noted
L3 ANSWER 8 OF 10 EMBASE COPYRIGHT 1999 ELSEVIER
SCI B V
AN 91246508 EMBASE
DN 1991246508
TI Erratum: The bovine .alpha.- ***lactalbumin***
promoter
directs expression of ovine trophoblast interferon in the mammary
gland of
transgenic mice (FEBS Letters (1991) Vol. 284 (19-22)).
AU Sinnakre M.G.; Volotte J.L.; Soulier S.; L'Haridon R.; Charlier
M.; Gaye
P.; Mercier J.C.
CS Laboratoire de Genetique Biochimique, I.N.R.A., 78350
Jouy-en-Josas,
France
SO FEBS Letters, (1991) 288(1-2) (247).
ISSN: 0014-5793 CODEN: FEBLAL
CY Netherlands
DT Journal; Errata
FS 029 Clinical Biochemistry
LA English
L3 ANSWER 9 OF 10 CAPLUS COPYRIGHT 1999 ACS
AN 1991:672494 CAPLUS
DN 115:272494
TI The bovine .alpha.- ***lactalbumin*** ***promoter***
directs
expression of ovine trophoblast interferon in the mammary gland of
transgenic mice [Erratum to document cited in
CA115(7):66096k]
AU Sinnakre, M. G.; Volotte, J. L.; Soulier, S.; L'Haridon, R.;
Charlier,
M.; Gaye, P.; Mercier, J. C.
CS Lab. Physiol. Comp., Univ. Paris VI, Paris, 75005, Fr.
SO FEBS Lett. (1991), 288(1-2), 247
CODEN: FEBLAL; ISSN: 0014-5793
DT Journal
LA English
AB Errors in the author addresses have been cor. The errors were not
reflected in the abstr. or the index entries.
L3 ANSWER 10 OF 10 MEDLINE
DUPLICATE

3
AN 91285097 MEDLINE
DN 91285097
TI The bovine .alpha.- ***lactalbumin*** ***promoter***
directs
expression of ovine trophoblast interferon in the mammary gland of
transgenic mice [published erratum appears in FEBS Lett
1991 Aug
19:288(1-2):247].
AU Sinnakre M.G.; Volotte J.L.; Soulier S.; L'Haridon R.; Charlier M;
Gaye P;
Mercier J C
CS Laboratoire de Physiologie Comparee, Universite Paris VI,
France..
SO FEBS LETTERS, (1991 Jun 17) 284 (1) 19-22.
Journal code: EUH ISSN: 0014-5793.
CY Netherlands
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals; Cancer Journals
EM 199110
AB A hybrid construct derived from ovine trophoblastin cDNA and
bovine
alpha-lactalbumin-encoding gene, was injected into the pronuclei of
mouse
eggs. In one of the resulting ***transgenic*** mouse lines,
expression
of the hybrid construct was detected and found to be limited to the
mammary gland of lactating females which secreted active ovine
trophoblastin. This strongly suggests that important cis-acting DNA
sequences involved in tissue-specific expression of the bovine gene
are
located within the second half of the 3' untranslated region, or/and
the
proximal 5' and 3' regions flanking the transcriptional unit.
=>
---Logging off of STN---
END
u
Unable to generate the STN prompt.
Exiting the script..
=>
=>
ALL L# QUERIES AND ANSWER SETS ARE DELETED AT
LOGOFF
LOGOFF? (Y)/N/HOLD Y
COST IN U.S. DOLLARS ENTRY SESSION TOTAL
FULL ESTIMATED COST 25.37 25.52
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

substrate for endogenous amidating activity in the mammary gland. Full characterization of the released sCT demonstrated it to be equivalent to synthetic standard in terms of structure, purity, and potency.

L3 ANSWER 4 OF 10 BIOSIS COPYRIGHT 1999 BIOSIS
 AN 1998:532816 BIOSIS
 DN PREV199800532816
 TI Production of ***transgenic*** pigs and mice containing the gene encoding human insulin-like growth factor I (IGF-I) under control of the bovine alpha-***lactalbumin*** **promoter*** and regulatory regions.
 AU Bleck, G. T. (1); Monaco, M. H.; Donovan, S. M.; Wheeler, M. B. (1)
 CS (1) Dep. Animal Sci., Univ. Ill., Urbana, IL USA
 SO Journal of Dairy Science, (1998) Vol. 81, No. SUPPL. 1, pp. 213.
 Meeting Info.: Joint Meeting of the American Dairy Science Association and the American Society of Animal Science Denver, Colorado, USA July 28-31, 1998
 1998 American Society of Animal Science
 ISSN: 0022-0302.
 DT Conference
 LA English

L3 ANSWER 5 OF 10 BIOSIS COPYRIGHT 1999 BIOSIS
 DUPLICATE 2
 AN 1996:186466 BIOSIS
 DN PREV199608742595
 TI Genetic modification of bovine beta-casein and its expression in the milk of ***transgenic*** mice.
 AU Choi, Byung-Kwon; Bleck, Gregory T.; Wheeler, Matthew B.; Jimenez-Flores, Rafael (1)
 CS (1) Dep. Dairy Sci., California Polytechnic State Univ., San Luis Obispo, CA 93407 USA
 SO Journal of Agricultural and Food Chemistry, (1996) Vol. 44, No. 3, pp. 953-960.
 ISSN: 0021-8561.
 DT Article
 LA English
 AB Genomic vectors containing mutant bovine beta-casein with putative glycosylation sites were constructed to study the functional properties of glycosylated beta-casein and its possible effects in milk. The mutation was performed by PCR-based site-directed mutagenesis. The tripeptide Asn-X-Ser, was generated between Asn-68 and Asn-73

in mature beta-casein. The resulting beta-casein mutants were designated pCJB68 and pCJB6873. pCJB68 carries a substitution of Ser-70 for Leu-70 (Asn-68-Ser-69-Ser-70-Pro-71), and pCJB6873 carries a substitution of Ser-70-Ser-71 for Leu-70-Pro-71 (Asn-68-Ser-69-Ser-70-Ser-71). The two mutated genomic constructs were placed under control of the bovine alpha-***lactalbumin*** **promoter***, and lines of mice expressing the pCJB68 and pCJB6873 have been established. The milk from ***transgenic*** mice contained bovine beta-casein at levels up to 2-3 mg/mL. N-Linked glycosylation of bovine beta-casein in the pCJB6873 line was confirmed by peptide-N-glycosidase F treatment, but glycosylation of bovine beta-casein did not occur in pCJB68 mice. In addition, mouse casein micelles containing glycosylated bovine beta-casein showed the largest median diameter and rough outer surface, compared to normal mouse casein micelles and micelles from ***transgenic*** milk containing bovine beta-casein.

L3 ANSWER 6 OF 10 CAPLUS COPYRIGHT 1999 ACS
 AN 1995:511593 CAPLUS
 DN 122:257982
 TI Modified alpha-lactalbumins containing few or no phenylalanines for dietary supplementation in hyperphenylalaninemia
 IN Colman, Alan; Wright, Gordon; Sawyer, Lindsay; Rigden, Daniel John
 PA Pharmaceutical Proteins Ltd., UK
 SO PCT Int. Appl., 71 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN CNT 2
 PATENT NO. KIND DATE APPLICATION NO.
 DATE

PI WO 9502692 A1 19950126 WO 1994-GB1514
 19940713
 W: AM, AU, BB, BG, BR, BY, CA, CN, CZ, FI, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LV, MD, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, US, UZ, VN
 RW: KE, MW, SD, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG

CA 2167155 AA 19950126 CA 1994-2167155 19940713
 AU 9471306 A1 19950213 AU 1994-71306 19940713
 AU 698597 B2 19981105
 EP 711344 A1 19960515 EP 1994-970557 19940713
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE
 CN 1127528 A 19960724 CN 1994-192790 19940713
 JP 09500273 T2 19970114 JP 1994-504407 19940713
 ZA 9405217 A 19960115 ZA 1994-5217 19940715
 WO 9602640 A1 19960201 WO 1995-GB1651 19950712
 W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT
 RW: KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG
 CA 2193513 AA 19960201 CA 1995-2193513 19950712
 AU 9528962 A1 19960216 AU 1995-28962 19950712
 AU 700224 B2 19981224
 EP 765390 A1 19970402 EP 1995-924467 19950712
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE
 CN 1157635 A 19970820 CN 1995-194129 19950712
 JP 10502816 T2 19980317 JP 1995-504802 19950712
 ZA 9505850 A 19960219 ZA 1995-5850 19950713
 PRAJ GB 1993-14802 19930716
 WO 1994-GB1514 19940713
 GB 1994-25326 19941215
 US 1995-381691 19950131
 GB 1995-3822 19950225
 WO 1995-GB1651 19950712
 AB Modified alpha-lactalbumin, e.g. of bovine or human origin, comprising fewer phenylalanine residues than wild-type alpha-lactalbumin may be used as a dietary component for sufferers of hyperphenylalaninemia. Preferably, all of the phenylalanine residues are replaced. The modified alpha-lactalbumin may be expressed in the mammary gland of non-human host animals so as to accumulate in, and if desired be sepd. from, their milk. Thus, residues Phe-9 (or Phe-3 in human), Phe-31, Phe-53, and Phe-80 in bovine alpha-lactalbumin are substituted with Tyr, Leu, Met, Trp, Ile, Ser, or Arg. If selection of the substituting amino acid is made solely on the basis of energy minimization and structural considerations, then Tyr3/9-Leu31-Tyr53-Tyr80 is preferred, if nutritional

***** STN Columbus *****

FILE 'HOME' ENTERED AT 14:12:22 ON 05 AUG 1999

=> file medline embase biosis inpadoc caplus

COST IN U.S. DOLLARS	ENTRY	SINCE FILE	TOTAL
FULL ESTIMATED COST	SESSION	0.15	0.15

FILE 'MEDLINE' ENTERED AT 14:12:34 ON 05 AUG 1999

FILE 'EMBASE' ENTERED AT 14:12:34 ON 05 AUG 1999

COPYRIGHT (C) 1999 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 14:12:34 ON 05 AUG 1999

COPYRIGHT (C) 1999 BIOSIS(R)

FILE 'INPADOC' ENTERED AT 14:12:34 ON 05 AUG 1999

COPYRIGHT (C) 1999 European Patent Office, Vienna (EPO)

FILE 'CAPLUS' ENTERED AT 14:12:34 ON 05 AUG 1999

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 1999 AMERICAN CHEMICAL SOCIETY (ACS)

=> s lactalbumin promoter/ab,bi

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

L1 23 LACTALBUMIN PROMOTER/AB,BI

=> s ll and transgen?/ab,bi

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

L2 17 L1 AND TRANSGEN?/AB,BI

=> dup rem l2

PROCESSING COMPLETED FOR L2

L3 10 DUP REM L2 (7 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 10 ANSWERS - CONTINUE? Y(N)y

L3 ANSWER 1 OF 10 MEDLINE DUPLICATE 1

AN 1999159717 MEDLINE

DN 99159717

TI Production of low-lactose milk by ectopic expression of intestinal lactase

in the mouse mammary gland [see comments].

CM Comment in: Nat Biotechnol 1999 Feb;17(2):135-6

AU Jost B; Vilotte J L; Duluc I; Rodeau J L; Freund J N

CS Institut National de la Sante et de la Recherche Medicale, Unite 381, Strasbourg, France.

SO NATURE BIOTECHNOLOGY, (1999 Feb) 17 (2) 160-4.

Journal code: CQ3. ISSN: 1087-0156.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199906

EW 19990603

AB We have investigated, in mice, an in vivo method for producing low-lactose milk, based on the creation of ***transgenic*** animals carrying a hybrid gene in which the intestinal lactase-philorizin hydrolase cDNA was placed under the control of the mammary-specific alpha-***lactalbumin***

promoter ***Transgenic*** females expressed lactase protein and activity during lactation at the apical side of mammary alveolar cells. Active lactase was also secreted into milk, anchored in the outer membrane of fat globules. Lactase synthesis in the mammary gland caused a significant decrease in milk lactose (50-85%) without obvious changes in fat and protein concentrations. Sucklings nourished with low-lactose milk developed normally. Hence, these data validate the use of ***transgenic*** animals expressing lactase in the mammary gland to produce low-lactose milk in vivo, and they demonstrate that the secretion of an intestinal digestive enzyme into milk can selectively modify its composition.

L3 ANSWER 2 OF 10 CAPLUS COPYRIGHT 1999 ACS

AN 1999482480 CAPLUS

TI Introduction of a proximal stat5 site in the murine alpha-***lactalbumin*** ***promoter*** induces prolactin dependency in vitro and improves expression frequency in vivo

AU Soulier, Solange, Lepourry, Laurence; Stinnakre, Marie-Georges; Langley, Brett; L'Huillier, Phil J.; Paly, Jacqueline; Djiane, Jean; Mercier, Jean-Claude; Vilotte, Jean-Luc

CS Laboratoire de GCntrique Biochimique et de Cytogknétique,

INRA, Jouv-en-Josas, 78352, Fr.

SO Transgenic Res (1999), 8(1), 23-31

CODEN: TRSEES; ISSN: 0962-8819

DT Journal

LA English

AB In order to establish a possible correlation between in vitro prolactin induction and the transcriptional activity of mammary gene promoters in ***transgenic*** mice, a functional Stat5-binding site was created by means of site-directed mutagenesis at position -70 on a 560 bp murine alpha-lactalbumin promoter linked to a CAT reporter gene. Surprisingly, the wild-type promoter was constitutively active in vitro and could not be induced by prolactin. Introducing the proximal Stat5 site abolished this constitutive activity and resulted in prolactin dependence in both CHO-K1- and HCl I-transfected cells. In ***transgenic*** mice, both the frequency of lines expressing the ***transgenic*** and the prevalence of mid to late pregnancy expression were increased.

L3 ANSWER 3 OF 10 BIOSIS COPYRIGHT 1999 BIOSIS

AN 1998479280 BIOSIS

DN PREV199800479280

TI Production of biologically active salmon calcitonin in the milk of ***transgenic*** rabbits.

AU McKee, Colin (1); Gibson, Allan; Dalrymple, Mike; Emslie, Liz; Garner, Ian; Cottingham, Ian

CS (1) PPL Therapeutics Ltd, Roslin, Edinburgh EH25 9PP UK

SO Nature Biotechnology, (July, 1998) Vol. 16, No. 7, pp. 647-651. ISSN: 1087-0156.

DT Article

LA English

AB Salmon calcitonin (sCT) is an example of one of the many bioactive peptides that require amidation of the carboxy terminus for full potency. We describe a method for the production of amidated sCT in the mammary gland of ***transgenic*** rabbits. Expression of a fusion protein comprising human alpha lactalbumin joined by an enterokinase cleavable linker to sCT was directed to the mammary gland under the control of the ovine beta lactoglobulin promoter. C-terminal amidation in vivo was achieved by extending the sCT by a single glycine residue that provides a

production of detectable levels of 2'-fucosyl-lactose in the milk of said mammal.

CLMS(2)

2. The mammal according to claim 1, wherein said mammal is selected from the group consisting of a mouse, a rabbit, a pig, a goat, a sheep and a cow.

=> e meade, harry

E#	FILE	FREQUENCY	TERM
E1	USPAT	1	MEADD/BI
E2	USPAT	306	MEADE/BI
E3	USPAT	0	=> MEADE, HARRY/BI
E4	USPAT	1	MEADELSON/BI
E5	USPAT	6	MEADEN/BI
E6	USPAT	20	MEADER/BI
E7	USPAT	3	MEADERING/BI
E8	USPAT	3	MEADERS/BI
E9	USPAT	1	MEADEVILLE/BI
E10	USPAT	1	MEADHATCHER/BI
E11	USPAT	2	MEADI/BI
E12	USPAT	4	MEADIA/BI

=> e meade, harry/in

E#	FILE	FREQUENCY	TERM
E1	USPAT	2	MEADE, EDWIN M/IN
E2	USPAT	1	MEADE, GEORGE E/IN
E3	USPAT	5	=> MEADE, HARRY/IN
E4	USPAT	3	MEADE, HARRY M/IN
E5	USPAT	1	MEADE, HAZEL/IN
E6	USPAT	3	MEADE, HAZEL W/IN
E7	USPAT	3	MEADE, JAMES H/IN
E8	USPAT	1	MEADE, JAMES M/IN
E9	USPAT	1	MEADE, JAMES P/IN
E10	USPAT	3	MEADE, JAMES R/IN
E11	USPAT	1	MEADE, JEFFREY/IN
E12	USPAT	5	MEADE, JOHN/IN

=> s e3-e4

5 "MEADE, HARRY"/IN
3 "MEADE, HARRY M"/IN

L15 8 ("MEADE, HARRY"/IN OR "MEADE, HARRY M"/IN)

=> d l- fro

US PAT NO: 5,849,992 [IMAGE AVAILABLE] L15: 1 of 8

DATE ISSUED: Dec. 15, 1998

TITLE: Transgenic production of antibodies in milk
INVENTOR: **Harry Meade**, Newton, MA

Paul DiTullio, Framingham, MA
Daniel Pollock, Medway, MA

ASSIGNEE: Genzyme Transgenics Corporation, Framingham, MA
(U.S. corp.)

APPL-NO: 08/410,887

DATE FILED: Mar. 27, 1995

REL-US-DATA: Division of Ser. No. 170,579, Dec. 20, 1993.

INT-CL: [6] C12N 5/00; C12N 15/00

US-CL-ISSUED: 800/2, DIG.1

US-CL-CURRENT: 800/14, 7, 15, 16, 17, 18

SEARCH-FLD: 800/2, DIG.1; 435/172.3

REF-CITED: U.S. PATENT DOCUMENTS

4,816,397	3/1989	Boss et al.	425/68
4,816,567	3/1989	Cabily et al.	530/387
4,873,316	10/1989	Meade et al.	530/412
5,372,775	6/1994	Clark et al.	435/69.1

FOREIGN PATENT DOCUMENTS

WO 90/04036 10/1989 World Intellectual Property Organization/21/8

WO 90/04036 4/1990 World Intellectual Property Organization

WO 92/03918 8/1991 World Intellectual Property Organization

WO 93/12227 12/1992 World Intellectual Property Organization

OTHER PUBLICATIONS

NM Greenberg et al. (1991) Proc Natl Acad Sci USA 88: 8327-8331.

MA Persuy et al. (1992) European J Biochem 205:887-893.

Hennighausen, L., Sippel, A. (1982) European Journal of Biochemistry

25:131-141.

Gordon, K., Lee, E., Vitale, J., Smith, A., Westphal, H., Hennighausen,

L. (1987) Bio/Technology 5:1183-1187.

Clark, A., Simons, P., Wilmut, I., Lathe, R. (1987) Bio/Technology 5:20/24.

Brunt, J. (1988) Bio/Technology 6:1149-1154.

Clark, A., Bessos, H., Bishop, J., Brown, P., Harris, S., Lathe, R., McClenaghan, M., Prowse, C., Simons, J., Whitelaw, C., Wilmut, I., (1989) Bio/Technology 7:487-492.

Meade, H., Gates, L., Lacy, E., Lonberg, N., (1990) Bio/Technology 8:443-446.

Buhler, Th. A., Bruyere, Th., Went, D., Stranzinger, G., Buiki, K., (1990) Bio/Technology 8:140-143.

Weidle, Y. H., Lenz, H., Brem, G. (1991) Gene 98 (2):185-91.

DiTullio, P., Cheng, S., Marshall, J., Grgory, R., Ebert, K., Meade, H., Smith, a., (1992) Bio/Technology 10:74-77.

Roberts, B., DiTullio, P., Vitale, J., Hehir, K., Gordon, K., (1992) Gene. 121:255-262.

Soulter et al., (1992) FEB 5 Letters, 297:13.

Bradshaw, J.P. et al., "The hormonal control of protein N-glycosylation

in the developing rabbit mammary gland and its effect upon transferrin synthesis and secretion" *Biochimica et Biophysica Acta*, 847: 344-351 (1985).

Buler, Th.A. et al., "Rabbit beta.-Casein Promoter Directs Secretion of Human Interleukin-2 Into the Milk of Transgenic Rabbits" *Bio/Technology*, 8: 140-143 (1990).

Clark, A.J. et al., "Expression of Human Anti-Hemophilic Factor IX in the Milk of Transgenic Sheep" *Bio/Technology*, 7: 487-492 (1989).

Clark, A.J. et al., "Pharmaceuticals From Transgenic Livestock" *Trends in Biotechnology*, 5(1): 20-24 (1987).

DiTullio, P. et al., "Production of Cystic Fibrosis Transmembrane Conductance Regulator in the Milk of Transgenic Mice" *Bio/Technology*, 10: 74-77 (1992).

Ebert, K.M. et al., "Transgenic Production of a Variant of Human Tissue-Type Plasminogen Activator in Goat Milk: Generation of Transgenic Goats and Analysis of Expression" *Bio/Technology*, 9: 835-838 (1991).

Fiat, A.-M. and P. Jolles "Caseins of Various Origins and Biologically Active Casein Peptides and Oligosaccharides: Structural and Physiological Aspects" *Molecular and Cellular Biochemistry*, 87: 5-30 (1989).

Gordon, K. et al., "Production of Human Tissue Plasminogen Activator in Transgenic Mouse Milk" *Bio/Technology*, 5: 1183-1187 (1987).

Hennighausen, L. and A.E. Sippel, "Characterization and Cloning of mRNAs Specific for the Lactating Mouse Mammary Gland" *Eur. J. Biochim.*, 131-141 (1982).

Hiatt, A. et al., "Production of antibodies in transgenic plants" *Nature*, 342: 76-78 (1989).

Horwitz, A.H., "Secretion of functional antibody and Fab fragment from yeast cells" *Proc. Natl. Acad. Sci. USA*, 85: 8678-8682 (1988).

Houdebine, L.-M., "Production of pharmaceutical proteins from animals" *Journal of Biotechnology*, 34: 269-287 (1994).

Larson, B.L., "Biosynthesis and Secretion of Milk Proteins: A Review" *Journal of Dairy Research*, 46: 161-174 (1979).

McKenzie, H.A. (editor), "Milk Proteins in Retrospect" in *Milk Proteins: Chemistry and Molecular Biology* (NY:Academic Press) 1: 3-15, 26-29 (1970).

McKenzie, H.A. (editor), *Milk Proteins: Chemistry and Molecular Biology* (NY:Academic Press), 2: 338-339 (1971).

Purset, G. and C.E. Rexroad, Jr., "Status of Research with Transgenic

Farm Animals" J. Anim. Sci., 71(Suppl. 3): 10-19 (1993).
 Roberts, B. et al., "Cloning of the goat beta-casein-encoding gene and expression in transgenic mice" Gene, 121: 255-262 (1992).
 Soulier, S. et al., "Expression analysis of ruminant alpha-lactalbumin in transgenic mice: developmental regulation and general location of important cis-regulatory elements" FEBS, 297(1,2): 13-18 (1992).
 Steiger, N. et al., "Self assembly of immunoglobulins in the cytoplasm of the alga Acetabularia mediterranea" Plant Science, 73: 181-190 (1991).
 an Brunt, J., "Molecular Farming: Transgenic Animals As Bioreactors" Bio/Technology, 6(10): 1149, 1151-2, 1154 (1988).
 Wall, R.J., "Transgenic Livestock: Progress and Prospects for the Future" Theriogenology, 45: 57-68 (1996).
 Watson, J.D. et al., Recombination at the Molecular Level in Molecular Biology of the Gene, 4th ed. (Reading, MA: The Benjamin/Cummings Publishing Co., Inc.) 1: 313 (1987).
 Weidle, U.H. et al., "Genes encoding a mouse monoclonal antibody are expressed in transgenic mice, rabbits and pigs" Gene, 98: 185-191 (1991).
 ART-UNIT: 162
 PRIM-EXMR: Bruce R. Campbell
 LEGAL-REP: Lahive & Cockfield, LLP

ABSTRACT:
 A method for the production of monoclonal antibodies in mammal's milk through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.
 10 Claims, 4 Drawing Figures

US PAT NO: 5,843,705 [IMAGE AVAILABLE] L15: 2 of 8
 DATE ISSUED: Dec. 1, 1998
 TITLE: Transgenically produced antithrombin III
 INVENTOR: Paul DiTullio, Framingham, MA
 Harry Meade, Newton, MA
 Edward S. Cole, Mendon, MA
 ASSIGNEE: Genzyme Transgenic Corporation, Framingham, MA (U.S. corp.)
 APPL-NO: 08/391,743
 DATE FILED: Feb. 21, 1995
 INT-CL: [6] C12P 21/06; C12N 9/48
 US-CL-ISSUED: 435/69.1; 530/393, 392, 386, 360, 412, 832; 514/8, 21; 435/320.1, 172.3, 172.1, 325; 800/2; 424/152.1, 535; 930/240
 US-CL-CURRENT: 800/7; 424/157.1, 535; 435/212, 320.1, 325; 514/8, 21; 530/360, 380, 386, 392, 393, 412, 832; 930/240
 SEARCH-FLD: 435/320.1, 172.3, 240.2, 69.1, 325, 212, 172.1;

530/412, 360, 393, 70, 386, 832, 380; 514/8, 21; 800/2; 424/157.1, 535
 REF-CITED: U.S. PATENT DOCUMENTS
 4,517,294 5/1985 Bock et al. 435/69.4
 4,632,981 12/1986 Bock et al. 514/8
 4,873,316 10/1989 Meade et al. 530/412
 5,366,894 11/1994 Clarke et al. 435/320.1

OTHER PUBLICATIONS
 Bock, S. et al., "Cloning and Expression of the cDNA for Human Antithrombin III", Nucleic Acids Research, vol. 10 (24), pp. 8113-8125 (1982).
 Wall, R., "Transgenic Livestock: Progress and Prospects for the Future", Theriogenology, vol. 45, pp. 57-68 (1996).
 Fan et al J. of Biol. Chem. 268(23):17588-96 1993.
 Cole et al. J. of Cellular Biochemistry Supplement O (180), 1994 p. 265.
 Edmundo et al J. of Cellular Biochemistry Supplement O(180) 1994 p. 265.
 Wall Theriogenology 45:57-68 (1996).
 ART-UNIT: 163
 PRIM-EXMR: Christopher S.F. Low
 LEGAL-REP: Louis Myers

ABSTRACT:
 This invention relates to transgenically produced human Antithrombin III (tgATIII). The human ATIII produced by the transgenic process of the present invention has a monosaccharide composition which comprises N-acetylgalactosamine (GalNAc) along with fucose, N-acetylglucosamine, galactose, mannose, and N-acetylneuraminic acid/N-glycolylneuraminic acid.
 The monosaccharide composition differs with that of plasma derived ATIII (phATIII). It has been found that tgATIII has an increased clearance rate when compared to phATIII.
 13 Claims, 11 Drawing Figures

US PAT NO: 5,827,690 [IMAGE AVAILABLE] L15: 3 of 8
 DATE ISSUED: Oct. 27, 1998
 TITLE: Transgenic production of antibodies in milk
 INVENTOR: **Harry Meade**, Newton, MA
 Paul DiTullio, Framingham, MA
 Daniel Pollock, Medway, MA
 ASSIGNEE: Genzyme Transgenics Corporation, Framingham, MA (U.S. corp.)
 APPL-NO: 08/170,579
 DATE FILED: Dec. 20, 1993

INT-CL: [6] C12P 21/04; C12N 15/00
 US-CL-ISSUED: 435/69.6, 172.3; 530/867, 800/2, DIG.1; 935/60
 US-CL-CURRENT: 800/7; 530/867
 SEARCH-FLD: 435/172.3, 69.1, 69.6; 530/867; 536/24.1; 800/2, DIG.1; 935/60
 REF-CITED: U.S. PATENT DOCUMENTS
 4,816,397 3/1989 Boss et al. 435/68
 4,816,567 3/1989 Cabilly et al. 530/387
 4,873,316 10/1989 Meade et al. 530/412
 5,322,775 6/1994 Clark et al. 435/69.1

FOREIGN PATENT DOCUMENTS
 WO 90/04036 10/1989 World Intellectual Property Organization 21/8
 WO 92/03918 8/1991 World Intellectual Property Organization
 WO 93/12227 12/1992 World Intellectual Property Organization

OTHER PUBLICATIONS
 Henninghausen, L., Sippel, A. (1982) European Journal of Biochemistry 25:131-141.
 Gordon, K., Lee, E., Vitale, J., Smith, A., Westphal, H., Henninghausen, L. (1987) Bio/Technology 5:1183-1187.
 Clark, A., Simons, P. Wilmut, I., Lathe, R. (1987) Bio/Technology 5:20-24.
 Brunt, J. (1988) Bio/Technology 6:1149-1154.
 Clark, A., Bessos, H., Bishop, J., Brown, P., Harris, S., Lathe, R., McClenaghan, M., Prowse, C., Simons, J., Whitelaw, C., Wilmut, L., (1989) Bio/Technology 7:487-492.
 Meade, H., Gates, L., Lacy, E., Lonberg, N., (1990) Bio/Technology 8:443-446.
 Buhler, Th. A., Bruyere, Th., Went, D., Stranzinger, G., Bulki, K., (1990) Bio/Technology 8:140-143.
 Weidle, U. H., Lenz, H., Brem, G. (1991) Gene 98 (2):185-91.
 DiTullio, P., Cheng, S., Marshall, J., Gregory, R., Ebert, K., Meade, H., Smith, A., (1992) Bio/Technology 10:74-77.
 Roberts, B., DiTullio, P., Vitale, J., Hehir, K., Gordon, K., (1992) Gene 121:255-262.
 Soulier et al., (1992) FEB 5 Letters, 297:13.
 Fiat, A. and Jolles, P., "Caseins of Various Origins and Biologically Active Casein Peptides and Oligosaccharides: Structural and Physiological Aspects," Molecular and Cellular Biochemistry vol. 87, 5-30 (1989).
 Larson, B., "Biosynthesis and Secretion of Milk Proteins: A Review," Journal of Dairy Research, vol. 46, 161-174 (1979).
 McKenzie, H. (editor), Milk Proteins Chemistry and Molecular Biology.
 Academic Press, vol. 1, pp. 3-15, and 26-29 (1970).
 McKenzie, H. (editor), Milk Proteins Chemistry and Molecular Biology.
 Academic Press, vol. 2, pp. 338-339 (1971).

Dec 20 1997 1997

AH Horwitz et al (1988) Proc Natl Acad Sci USA 85: 8678-8682.
M Steiger et al (1991) Plant Science 73: 181-190.
A Hatt et al (1989) Nature 342: 76-78.
JP Bradshaw et al (1955) Biochimica et Biophysica Acta 847: 344-351.
KM Ebert et al (1991) Bio/Technology 9: 835-838.
R Bischoff et al (1992) FEBS Letters 305: 265-268.
H Meade et al (1990) Bio/Technology 8: 443-446.
M-A Persuy et al (1992) European J Biochemistry 205: 887-893.
B Roberts et al (1992) Gene 121: 255-262.
NM Greenberg et al (1991) Proc Natl Acad Sci USA 88: 8327-8331.
JS Logan (1993) Current Opinion in Biotechnology 4: 591-595.
JD Watson et al (1987) Molecular Biology of the Gene p. 313.
VG Pursel et al (1993) J Animal Sci 71 (Supp 3): 10-19.
RJ Wall (1996) Theriogenology 45: 57-68.
LM Houdebine (1994) J Biotechnol 34: 269-287.
ART-UNIT: 184
PRIM-EXMR: Bruce R. Campell
LEGAL-REP: Lahive & Cockfield, LLP

ABSTRACT:
A method for the production of monoclonal antibodies in mammal's milk, through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.
13 Claims, 4 Drawing Figures

US PAT NO: 5,750,172 [IMAGE AVAILABLE] L15: 4 of 8

DATE ISSUED: May 12, 1998
TITLE: Transgenic non human mammal milk
INVENTOR: **Harry Meade**, Newton, MA
Nils Lönberg, New York, NY
ASSIGNEE: Pharming B.V., Leiden, Netherlands (foreign corp.)
APPL-NO: 08/460,959
DATE FILED: Jun. 5, 1995
REL-US-DATA: Continuation of Ser. No. 322,984, Oct. 14, 1994, which is a continuation of Ser. No. 109,865, Aug. 20, 1993, abandoned, which is a continuation of Ser. No. 332,293, Mar. 31, 1989, abandoned, which is a division of Ser. No. 65,994, Jun. 23, 1987, Pat. No. 4,873,316.
INT-CL: [6] C12P 21/06; C12P 21/02; C12P 21/04
US-CL-ISSUED: 426/580; 435/69.1, 69.4, 69.51, 69.52, 69.6, 183, 215;
800/2, DIG.1
US-CL-CURRENT: 426/580; 435/69.1, 69.4, 69.51, 69.52, 69.6, 183, 215;
800/7
SEARCH-FLD: 800/2, DIG.1, 3, 4; 435/172.1, 69.1, 69.4, 69.51, 69.52, 69.6, 215, 183; 935/63, 9, 11, 13, 14, 53; 530/832, 833; 426/580
REF-CITED: U.S. PATENT DOCUMENTS
5,304,489 4/1994 Rosen 435/320.1

5,322,775 6/1994 Clark et al.
5,565,362 10/1996 Rosen

OTHER PUBLICATIONS
Van Brunt, "Molecular Farming: Transgenic Animals as Bioreactors", Bio/Technology, vol. 6, No. 10, pp. 1149-1154, Oct. 1988.
Andres et al., "The Ha-Ras Oncogene Directed by a Milk Protein Gene Promoter: Expression and Tumor Induction in Transgenic Mice", Journal of Cellular Biochemistry, Supplement 11C, p. 153, Abstract No. L651, 1987.
Clark et al., "Pharmaceuticals from Transgenic Livestock", Trends in Biotechnology, vol. 5, pp. 20-24, Jan. 1987.
Hanahan, "Heritable Formation of Pancreatic beta-Cell Tumours in Transgenic Mice Expressing Recombinant Insulin/Simian Virus 40 Oncogenes", Nature, vol. 315, pp. 115-122, May 9, 1985.
Overbeek et al., "Lens-specific Expression and Developmental Regulation of the Bacterial Chloramphenicol Acetyltransferase Gene Driven by the Murine alphaA-Crystallin Promoter in Transgenic Mice", Proceedings of the National Academy of Sciences USA, Volum. Dec. 1985.
Yu-Lee et al., "Evolution of the Casein Multigene Family: Conserved Sequences in the 5' Flanking and Exon Regions", Nucleic Acids Research, vol. 14, No. 4, pp. 1883-1902, 1986.
Campbell et al., "Comparison of the Whey Acidic Protein Genes of the Rat and Mouse", Nucleic Acids Research, vol. 12, Nov. 22, pp. 8685-8697, 1984.
ART-UNIT: 189
PRIM-EXMR: Deborah Crouch
LEGAL-REP: Townsend & Townsend & Crew LLP

ABSTRACT:
This invention relates to the production of recombinant proteins, such as coagulation factors VIII and IX, tissue plasminogen activator (TPA), urokinase, growth hormone, insulin, interferons, interleukins, peptide hormones and immunoglobulins, in mammals' milk. Particularly, this invention relates to an expression system which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.
5 Claims, No Drawings

US PAT NO: 5,688,677 [IMAGE AVAILABLE] L15: 5 of 8

DATE ISSUED: Nov. 18, 1997
TITLE: Deoxyribonucleic acids containing inactivated hormone

responsive elements
INVENTOR: Karl M. Ebert, Millbury, MA
Paul DiTullio, Framingham, MA
Seng Hing Cheng, Wellesley, MA
Harry M. Meade, Newton, MA
Alan Edward Smith, Dover, MA

ASSIGNEE: Genzyme Corporation, Framingham, MA (U.S. corp.)
APPL-NO: 08/135,809
DATE FILED: Oct. 13, 1993
INT-CL: [6] C12N 15/06; C12N 15/12
US-CL-ISSUED: 435/240.1; 536/24.1, 23.5
US-CL-CURRENT: 536/23.5, 24.1
SEARCH-FLD: 536/23.5, 24.1; 514/44; 435/6, 172.3, 240.1, 69.1; 800/2
REF-CITED: U.S. PATENT DOCUMENTS
5,240,846 8/1993 Collins et al. 435/240.1

FOREIGN PATENT DOCUMENTS
91/02796 3/1991 World Intellectual Property Organization
91/10734 7/1991 World Intellectual Property Organization
93/12240 6/1993 World Intellectual Property Organization
93/12756 8/1993 World Intellectual Property Organization

OTHER PUBLICATIONS
Hyde, S.C. et al. (1993) "Correction of the Ion Transport Defect in Cystic Fibrosis Transgenic Mice by Gene Therapy" Nature 362:250-255.
Clarke, L.L. et al. (1992) "Defective Epithelial Chloride Transport in a Gene-Targeted Mouse Model of Cystic Fibrosis" Science 257:1125-1128.
Higgins, C.F. et al. (1992) "Cystic Fibrosis Mice Have Arrived" Hum. Mol. Gen. 1(7):459-460.
Snouwaert, J.N. et al. (1992) "An Animal Model For Cystic Fibrosis Made By Gene Targeting" Science 257:1083-1088.
Rosenfeld, M.A. et al. (1992) "In Vivo Transfer of the Human Cystic Fibrosis Transmembrane Conductance Regulator Gene to the Airway Epithelium" Cell 68:143-155.
Yoshimura, K. et al. (1992) "Expression of the Human Cystic Fibrosis Transmembrane Conductance Regulator Gene in the Mouse Lung After In Vivo Intratracheal Plasmid-Mediated Gene Transfer" Nucleic Acids Res. 20(12):3233-3240.
Smith, A.E. (1992) "Emerging Therapies for Cystic Fibrosis" Section V-Topics in Biology in Ann. Rep. Med. Chem. 27:235-243.
Anguiano, A. et al. (1992) "Congenital Bilateral Absence of the Vas Deferens: A Primarily Genital Form of Cystic Fibrosis" JAMA 267:1794-1797.
Anderson, M.P. et al. (1992) "Regulation by ATP and ADP of CFTR Channels That Contain Mutant Nucleotide-Binding Domains"

Science
257:1701-1704.

Kartner, N. et al. (1991) "Expression of the Cystic Fibrosis Gene in Non-Epithelial Invertebrate Cells Produces a Regulated Anion Conductance." *Cell* 64:681-691.

Koller, B.H. et al. (1991) "Toward an Animal Model of Cystic Fibrosis: Targeted Interruption of Exon 10 of the Cystic Fibrosis Transmembrane Regulator Gene in Embryonic Stem Cells." *Proc. Natl. Acad. Sci.* 88:10730-10734.

Rich D.P et al. (1991) "Effect of Deleting the R Domain on CFTR Generated Chloride Channels." *Science* 253:205-207.

Rosenfeld, M.A. et al. (1991) "In Vivo Transfer of the Human Cystic Fibrosis Gene to the Respiratory Epithelium." *Clin. Res.* 39(2):311 A.

Tsui, L.-C. et al. (1991) "Biochemical and Molecular Genetics of Cystic Fibrosis." *Advances in Human Genetics* 20:153-266j.

Tabcharani, J.A. et al. (1991) "Phosphorylation-Regulated Cl-Channel in CHO Cells Stably Expressing the Cystic Fibrosis Gene." *Nature* 352:628-631.

Drumm, M.L. et al. (1990) "Correction of the Cystic Fibrosis Defect in *Vitro* By Retrovirus-Mediated Gene Transfer." *Cell* 62:1227-1233j.

Gregory, R.J. et al. (1990) "Expression and Characterization of the Cystic Fibrosis Transmembrane Conductance Regulator." *Nature* 347:382-386j.

Rich, D.P. et al. (1990) "Expression of the Cystic Fibrosis Transmembrane Conductance Regulator Corrects Defective Chloride Channel Regulation in Cystic Fibrosis Airway Epithelial Cells." *Nature* 347:358-363j.

Huang, M.T.F. et al. (1990) "Intervening Sequences Increase Efficiency of RNA 3' Processing and Accumulation of Cytoplasmic RNA." *Nucleic Acids Res.* 18(4):937-947j.

Riordan, J.R. et al. (1989) "Identification of the Cystic Fibrosis Gene: Cloning and Characterization of Complementary DNA." *Science* 245:1066-1073j.

Rommens, J.H. et al. (1989) "Identification of the Cystic Fibrosis Gene: Chromosome Walking and Jumping." *Science* 245: 1059-1065j.

Kerem, B.S. et al. (1989) "Identification of the Cystic Fibrosis Gene: Genetic Analysis." *Science* 245:1073-1080j.

Yoshimura, K. et al. (1993) "Adenovirus-mediated Augmentation of Cell Transfection with Unmodified Plasmid Vectors." *J. Biol. Chem.* 268(4):2300-2303j.

Engelhardt, J.F. et al. (1993) "Direct Gene Transfer of Human CFTR into Human Bronchial Epithelia of Xenografts with E1-deleted Adenovirus"

Nat. Gen. 4:27-34j.

Flotte, T.R. et al. (1993) "Expression of the Cystic Fibrosis Transmembrane Conductance Regulator from a Novel Adeno-associated Virus Promoter." *J. Biol. Chem.* 268(5):3781-3790j.

Johnson, L.G. et al. (1992) "Efficiency of Gene Transfer for Restoration of Normal Airway Epithelial Function in Cystic Fibrosis." *Nat. Gen.* 2:21-25.

Van Brunt, Bio/Technology 6: 1150 (1988).

ART-UNIT: 184

PRIM-EXMR: James Martinell

ABSTRACT:
A DNA comprising at least one inactivated hormone responsive element and a nucleic acid sequence encoding a membrane-associated protein is described. Therapeutic compositions and cells including the DNA are also described. Other aspects of the invention include methods of treating subjects having cystic fibrosis which include administering an effective amount of the DNA to subjects having cystic fibrosis such that functional cystic fibrosis transmembrane conductance regulator is produced by the subject at a level which is not detrimental to the subject. The present invention also pertains to a method of introducing the DNA into a cell such that the membrane-associated protein is produced at a level which is not detrimental to the cell and cells produced by this method. Still other aspects of the invention include a method of assaying DNA for the presence or absence of a hormone responsive element in a species in which the hormone responsive element is functional and a method of selectively breeding female transgenic mammals which produce a protein of interest.

11 Claims, 14 Drawing Figures

US PAT NO: 5,272,254 [IMAGE AVAILABLE] L15: 6 of 8

DATE ISSUED: Dec. 21, 1993

TITLE: Production of streptavidin-like polypeptides

INVENTOR: **Harry M. Meade**, Newton, MA
Jeffrey L. Garwin, Bedford, MA

ASSIGNEE: Biogen Inc., Cambridge, MA (U.S. corp.)

APPL-NO: 07/800,158

DATE FILED: Nov. 27, 1991

REL-US-DATA: Division of Ser. No. 185,329, Apr. 21, 1988, Pat. No.

5,168,049, which is a continuation of Ser. No. 656,873, Oct. 2, 1984, abandoned.

INT-CL: [5] C07K 13/00; C07K 7/00

US-CL-ISSUED: 530/350, 300, 825

US-CL-CURRENT: 530/350, 300, 825

SEARCH-FLD: 530/350, 825, 300

REF-CITED:

U.S. PATENT DOCUMENTS
4,338,397 7/1982 Gilbert et al. 435/69.1
4,411,994 10/1983 Gilbert et al. 435/69.7
4,839,293 6/1989 Cantor et al.

FOREIGN PATENT DOCUMENTS
0068867A2 1/1983 European Patent Office
2045251A 10/1980 United Kingdom

OTHER PUBLICATIONS
Dailey et al. *J. Biol. Chem.* 253:8203-8209.
Berk, A. J. and P. A. Sharp, "Sizing and Mapping of Early Adenovirus mRNAs by Gel Electrophoresis of S1 Endonuclease-Digested Hybrids," *Cell* 12, pp. 721-732 (Nov. 1977).
Bollen, A. et al., "Cloning and Expression in *Escherichia coli* of Full-Length Complementary DNA Coding for Human α -sub.1 -Antitrypsin," *DNA* 2, pp. 255-264 (1983).
Grunstein, M. and D. S. Hogness, "Colony hybridization: A method for the isolation of cloned DNAs that contain a specific gene," *Proc. Natl. Acad. Sci. USA*, 72, pp. 3961-3965 (Oct. 1975).
Heppel, L. A., "Preparation of Cells of *Escherichia coli* with Altered Permeability," in *Methods in Enzymology*, 12B, pp. 841-847 (1968).
Hohn, B. and J. Collins, "A small cosmid for efficient cloning of large DNA fragments," *Gene*, 11, pp. 291-298 (1980).
Maniatis, T. et al., "Construction of Genomic Libraries in Cosmid Vectors," in *Molecular Cloning: A Laboratory Manual*, pp. 295-305 (1982).
Maxam, A. M. and W. Gilbert, "Sequencing End-Labeled DNA with Base-Specific Chemical Cleavage," in *Methods in Enzymology*, 65, pp. 499-560 (1980).
Pennica, D. et al., "Cloning and expression of human tissue-type plasminogen activator cDNA in *E. coli*," *Nature*, 301, pp. 214-221 (Jan. 1983).
Roberts, T. M. and G. D. Lauer, "Maximizing Gene Expression on a Plasmid Using Recombination in *Vitro*," in *Methods in Enzymology*, 68, pp. 473-482 (1979).
Towbin, H. et al., "Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications," *Proc. Natl. Acad. Sci. USA*, 76, pp. 4350-4354 (Sep. 1979).
Vieira, J. and J. Messing, "The pUC plasmids, an M13mp7-derived system for insertion mutagenesis and sequencing with synthetic universal primers," *Gene*, 19, pp. 259-268 (1982).
Villa-Komaroff, L. et al., "A bacterial clone synthesizing proinsulin," *Proc. Natl. Acad. Sci. USA*, 75, pp. 3727-3731 (Aug. 1978).
Chater, K. et al., "Gene Cloning in Streptomyces," in *Current Topics in Microbiol. and Immunol.*, 96, pp. 69-75 (1982).
Katz, E. et al., "Cloning and Expression of the Tyrosinase Gene from

Streptomycetes antibiotics in Streptomycetes lividans," J. General Microbiol., 129, pp. 2703-2714 (1983).

Stapley, E. O. et al., "Antibiotic MSD-235, I. Production by Streptomycetes

avidinii and Streptomycetes lavendulae," Antimicrobial Agents and Chemotherapy, pp. 20-27 (1964).

Bayer, E. A. and M. Wilchek, "The Use of the Avidin-Biotin Complex as a

Tool in Molecular Biology," In Methods of Biochem. Anal., 26, pp. 1-45

(1980).

Chaiet, L. and F. J. Wolf, "The Properties of Streptavidin, a Biotin-Binding Protein Produced by Streptomyces," Arch. Biochem.

Biophys., 106, pp. 1-5 (1964).

Cuatrecasas, P. and M. Wilchek, "Single-Step Purification of Avidin from

Egg White by Affinity Chromatography on Bioctin-Sepharose Columns,"

Biochem. Biophys. Res. Commun., 33, pp. 235-239 (1968).

Finn, F. M. et al., "Hormone-Receptor studies with Avidin and Biotinylinsulin-Avidin Complexes," J. Biol. Chem., 255 pp. 5742-5746

(1980).

Heney, G. and G. A. Orr, "The Purification of Avidin and Its Derivatives

on 2-Iminobiotin-6-aminohexyl-Sepharose 4B," Anal. Biochem., 114, pp.

92-96 (1981).

Hofmann, K. et al., "Iminobiotin affinity columns and their application to retrieval of streptavidin," Proc. Natl. Acad. Sci. USA 77, pp. 4666-4668 (Aug. 1980).

Korenman, S. G. and B. W. O'Malley, "Newer Methods of Avidin Assay," In

Methods in Enzymology, 18A, pp. 427-430 (1970).

Orr, G. A., "The Use of the 2-Iminobiotin-Avidin Interaction for the Selective Retrieval of Labeled Plasma Membrane Components," J. Bio.

Chem., 256, pp. 761-766 (Jan. 1981).

ART-UNIT: 184

PRIM-EXMR: Robert A. Wax

ASST-EXMR: Gabriele E. Bugaisky

LEGAL-REP: James F. Haley, Denise L. Loring

ABSTRACT:

DNA sequences, hybrid DNA sequences, recombinant DNA molecules and processes for producing streptavidin-like polypeptides and for producing fused proteins consisting of a streptavidin-like polypeptide joined end to end with another protein, polypeptide, peptide or amino acid. The DNA sequences, hybrid DNA sequences and recombinant DNA molecules of this invention are characterized in that they include DNA fragments that code for streptavidin-like polypeptides. These DNA sequences, hybrid

DNA sequences and recombinant DNA molecules and the hosts transformed with them may be employed in the processes of this invention to produce streptavidin-like polypeptides and fused proteins.

5 Claims, 7 Drawing Figures

US PAT NO: 5,168,049 [IMAGE AVAILABLE] L15: 7 of 8

DATE ISSUED: Dec. 1, 1992

TITLE: Production of streptavidin-like polypeptides

INVENTOR: **Harry M. Meade**, Newton, MA

Jeffrey L. Garvin, Bedford, MA

ASSIGNEE: Biogen, Inc., Cambridge, MA (U.S. corp.)

APPL-NO: 07/185,329

DATE FILED: Apr. 21, 1988

REL-US-DATA: Continuation of Ser. No. 656,873, Oct. 2, 1984, abandoned.

INT-CL: [5] C12N 15/00; C12N 15/03; C12N 15/04; C12N 15/05; C12N 21/00; C12P 15/06; C12N 15/11; C12N 15/31; C12N 15/70; C12P 21/00; C12P 21/02

US-CL-ISSUED: 435/69.1, 69.7, 69.8, 172.3, 240.1, 240.2, 240.4, 252.3, 252.33, 252.35, 255, 256, 320.1; 536/27; 935/10, 11

US-CL-CURRENT: 435/69.1, 69.7, 69.8, 252.3, 252.33, 252.35, 254.11, 254.2, 320.1, 366

SEARCH-FLD: 435/68, 70, 71, 91, 172.1, 172.3, 252.3, 252.31-252.35, 255, 256, 320, 69.1, 71.2, 320.1; 514/2; 536/27; 935/10, 11, 22, 29, 33, 38, 39, 47, 48, 66-75

REF-CITED: U.S. PATENT DOCUMENTS

4,338,397 7/1982 Gilbert et al. 435/68

4,411,994 10/1983 Gilbert et al. 435/71

4,839,293 6/1989 Cantor et al. 435/320

OTHER PUBLICATIONS

Hamer et al., "Expression of the chromosomal mouse beta major-globin gene cloned in SV40", Nature 281: 35 (1979).

Maniatis et al., "Construction of Genomic Libraries in Cosmid Vectors", Molecular Cloning, A Laboratory Manual, pp. 295-305 (Maniatis, Fritsch and Sambrook, ed., 1982).

Edlund et al., "Isolation of cDNA sequences coding for a part of human tissue plasminogen activator", Proc. Natl. Acad. Sci. USA 80: 349 (1983).

Hoffmann et al., "Iminobiotin affinity columns and their application to retrieval of streptavidin", Proc. Natl. Acad. Sci. USA 77: 4666 (1980).

Roberts et al., "A general method for maximizing the expression of a cloned gene", Proc. Natl. Acad. Sci. USA 76: 760 (1979).

Hitzeman et al., "Expression of a human gene for interferon in yeast",

Nature 293: 717 (1981).

ART-UNIT: 185

PRIM-EXMR: James Martinell

LEGAL-REP: James F. Haley, Jr., Denise L. Loring

ABSTRACT:

DNA sequences, hybrid DNA sequences, recombinant DNA molecules and processes for producing streptavidin-like polypeptides and for producing fused proteins consisting of a streptavidin-like polypeptide joined end to end with another protein, polypeptide, peptide or amino acid. The DNA sequences, hybrid DNA sequences and recombinant DNA molecules of this invention are characterized in that they include DNA fragments that code for streptavidin-like polypeptides. These DNA sequences, hybrid DNA sequences and recombinant DNA molecules and the hosts transformed with them may be employed in the processes of this invention to produce streptavidin-like polypeptides and fused proteins.

35 Claims, 7 Drawing Figures

US PAT NO: 4,873,316 [IMAGE AVAILABLE] L15: 8 of 8

DATE ISSUED: Oct. 10, 1989

TITLE: Isolation of exogenous recombinant proteins from the milk of transgenic mammals

INVENTOR: **Harry Meade**, Newton, MA

Nils Lounberg, New York, NY

ASSIGNEE: Biogen, Inc., Cambridge, MA (U.S. corp.)

APPL-NO: 07/065,994

DATE FILED: Jun. 23, 1987

INT-CL: [4] C07K 3/02; C07K 3/12; C07K 3/18; C12N 15/00

US-CL-ISSUED: 530/412, 360, 361, 833, 832, 416, 417, 418, 435/68, 172.1, 172.3, 240.2; 935/53, 55, 70, 111; 800/1; 536/27, 28, 29

US-CL-CURRENT: 800/7; 435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8; 530/360, 361, 416, 417, 418, 832, 833; 536/23.1, 23.4, 23.5; 800/18

SEARCH-FLD: 435/68, 172.1, 172.3, 226, 240.2; 530/832, 833, 412, 360, 361, 303; 800/1; 935/53, 55, 70; 536/27, 28, 29

REF-CITED: U.S. PATENT DOCUMENTS

4,018,752 4/1977 Buhler et al. 530/382

4,229,342 10/1980 Mirabel 530/382

4,376,072 3/1983 Connolly 530/382

4,396,601 8/1983 Salser et al. 435/172.3

4,462,932 7/1984 Loneragan 530/382

4,644,056 2/1987 Kothe et al. 530/382

4,736,866 4/1988 Leder et al.

FOREIGN PATENT DOCUMENTS
0117059 8/1984 European Patent Office 435/172.3
0247494 12/1987 European Patent Office
0264166 4/1988 European Patent Office
WO88/00239 1/1988 World Intellectual Property Organization
WO88/01648 3/1988 World Intellectual Property Organization

OTHER PUBLICATIONS

Gordon et al., BioTechnology, 5, 1183-7, (Nov. 1987).
Lovell-Badge, Nature, 315, 628-629, (1985).
Hammer et al., Nature, 415, 680-683, (Jun. 20, 1985).
Garcia et al., Mol. Cell. Biol., 6(6), 1974-82, (1986).
Palmiter et al., J. Cell. Biochem., 8B, p. 25, Ab. #0890, (1984).
Fisher et al., J. B. C., 260 (20), 11223-11230, 1985.
Andres et al., Chem. Abs. 106(17):133024q, 1987 (for PNAS USA, 84(5)).
1299-303, May 1987).
Ross et al., P.N.A.S. (USA), 82, 5880-84, 1985.
Brinster et al., Cell, 27, 223-31, Nov. 1981.
ART-UNIT: 186
PRIM-EXMR: Margaret Moskowitz
ASST-EXMR: Jeff P. Kushan
LEGAL-REP: James F. Haley, Jr., Teresa L. Solomon

ABSTRACT:

This invention relates to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's casein promoter which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.

3 Claims, 1 Drawing Figures

=> d 1 - clms

US PAT NO: 5,849,997 [IMAGE AVAILABLE] L15: 1 of 8

CLAIMS:

CLMS(1)

What is claimed is:

1. A transgenic non-human mammal all of whose germ cells and somatic cells contain a heterologous immunoglobulin protein-coding sequence operatively linked to a promoter sequence that directs the preferential expression of said protein-coding sequence in mammary gland

epithelial cells, thereby providing a heterologous and assembled immunoglobulin in

the milk of said mammal wherein said heterologous and assembled immunoglobulin is in a functional configuration and is produced at levels of at least about 1 mg/ml in the milk of said mammal.

CLMS(2)

2. The transgenic mammal of claim 1 wherein said immunoglobulin comprises a tetrameric antibody directed against a pathogen.

CLMS(3)

3. The transgenic mammal of claim 1 wherein said immunoglobulin comprises a tetrameric antibody directed against a biologically active peptide.

CLMS(4)

4. The transgenic mammal of claim 1 wherein said biologically active peptide is selected from the group consisting of erythropoietin, tissue plasminogen activator and gamma interferon.

CLMS(5)

5. The transgenic mammal of claim 1 wherein said immunoglobulin comprises a tetrameric antibody directed against an enzyme.

CLMS(6)

6. The transgenic mammal of claim 1 wherein said mammal is selected from the group consisting of mice, cows, sheep, goats, and pigs.

CLMS(7)

7. The transgenic mammal of claim 1 wherein said promoter is selected from the group consisting of the casein promoter, the beta lactoglobulin promoter, the whey acid protein promoter, and the lactalbumin promoter.

CLMS(8)

8. The transgenic mammal of claim 1 wherein said immunoglobulin comprises heavy and light chains.

CLMS(9)

9. The transgenic mammal of claim 1 wherein said immunoglobulin is of human origin.

CLMS(10)

10. A transgenic non-human goat all of whose germ cells and somatic cells contain a heterologous immunoglobulin protein-coding sequence operatively linked to a promoter sequence that directs the preferential expression of said protein-coding sequence in mammary gland epithelial cells, thereby providing a heterologous and assembled immunoglobulin in the milk of said goat, wherein said heterologous and assembled immunoglobulin is in a functional configuration and is produced at levels of at least about 1 mg/ml in the milk of said goat.

US PAT NO: 5,843,705 [IMAGE AVAILABLE] L15: 2 of 8

CLAIMS:

CLMS(1)

The invention claimed is:

1. A method for producing human antithrombin III in goat milk, comprising:
a. producing a transgenic goat that expresses in mammary tissue a transgene which encodes a human antithrombin III, wherein the human antithrombin III is secreted into the milk produced by the transgenic goat;
b. collecting milk from the transgenic goat which milk contains the human antithrombin III; and
c. isolating the human antithrombin III from the collected milk, wherein the human antithrombin III isolated from milk has faster clearance time and an increased affinity for heparin both compared to human antithrombin III isolated from human plasma.

CLMS(2)

2. A glycosylated human antithrombin III, wherein a glycosylated human antithrombin III is made by the method of isolating the glycosylated human antithrombin III from milk, wherein the milk is collected from a transgenic goat, which goat expresses in mammary tissue a transgene encoding human antithrombin III and wherein the human antithrombin III is secreted into the milk produced by the transgenic goat; and wherein the glycosylated human antithrombin III has the following properties:
a) monosaccharide glycosylation comprising GalNAc;
b) no O-linked glycosylation;
c) faster plasma clearance time and an increased affinity for heparin both compared to human antithrombin III that is isolated from human plasma.

CLMS(3)

3. A glycosylated human antithrombin III, wherein a glycosylated human antithrombin III is made by the method of isolating the glycosylated human antithrombin III from milk, wherein the milk is collected from a transgenic goat, which goat expresses in mammary tissue a transgene encoding human antithrombin III and wherein the human antithrombin III is secreted into the milk produced by the transgenic goat; and wherein the glycosylated human antithrombin III has the following properties:
- a) monosaccharide glycosylation comprising Fuc, GalNAc, GlcNAc, Man, and NANA/NGNA;
 - b) no O-linked glycosylation;
 - c) faster plasma clearance time and an increased affinity for heparin both compared to human antithrombin III that is isolated from human plasma.

CLMS(4)

4. A glycosylated human antithrombin III, wherein a glycosylated human antithrombin III is made by the method of isolating the glycosylated human antithrombin III from milk, wherein the milk is collected from a transgenic goat, which goat expresses in mammary tissue a transgene encoding human antithrombin III and wherein the human antithrombin III is secreted into the milk produced by the transgenic goat; and wherein the glycosylated human antithrombin III has the following properties:
- a) monosaccharide glycosylation comprising oligomannose and/or hybrid oligosaccharide structures;
 - b) no O-linked glycosylation;
 - c) faster plasma clearance time and an increased affinity for heparin both compared to human antithrombin III that is isolated from human plasma.

CLMS(5)

5. A glycosylated human antithrombin III, wherein a glycosylated human antithrombin III is made by the method of isolating the glycosylated human antithrombin III from milk, wherein the milk is collected from a transgenic goat, which goat expresses in mammary tissue a transgene encoding human antithrombin III and wherein the human antithrombin III is secreted into the milk produced by the transgenic goat; and wherein the glycosylated human antithrombin III has the following properties:
- a) monosaccharide glycosylation comprising primarily an oligomannose or

- hybrid type structure on one site and complex oligosaccharide on the remaining 3 sites;
- b) no O-linked glycosylation;
 - c) faster plasma clearance time and an increased affinity for heparin both compared to human antithrombin III that is isolated from human plasma.

CLMS(6)

6. A glycosylated human antithrombin III, wherein a glycosylated human antithrombin III is made by the method of isolating the glycosylated human antithrombin III from milk, wherein the milk is collected from a transgenic goat, which goat expresses in mammary tissue a transgene encoding human antithrombin III and wherein the human antithrombin III is secreted into the milk produced by the transgenic goat; and wherein the glycosylated human antithrombin III has the following properties:
- a) a monosaccharide composition which is partially sialylated;
 - b) no O-linked glycosylation;
 - c) faster plasma clearance time and an increased affinity for heparin both compared to human antithrombin III that is isolated from human plasma.

CLMS(7)

7. A glycosylated human antithrombin III, wherein a glycosylated human antithrombin III is made by the method of isolating the glycosylated human antithrombin III from milk, wherein the milk is collected from a transgenic goat, which goat expresses in mammary tissue a transgene encoding human antithrombin III and wherein the human antithrombin III is secreted into the milk produced by the transgenic goat; and wherein the glycosylated human antithrombin III has the following properties:
- a) a monosaccharide composition comprising sialic acid which includes NGNA;
 - b) no O-linked glycosylation;
 - c) faster plasma clearance time and an increased affinity for heparin both compared to human antithrombin III that is isolated from human plasma.

CLMS(8)

8. A glycosylated human antithrombin III, wherein a glycosylated human antithrombin III is made by the method of isolating the glycosylated human antithrombin III from milk, wherein the milk is collected from a transgenic goat, which goat expresses in mammary tissue a transgene encoding human antithrombin III and wherein the human antithrombin III is

- secreted into the milk produced by the transgenic goat; and wherein the glycosylated human antithrombin III has the following properties:

- a) monosaccharide glycosylation comprising a fucose on its proximal GlcNAc on each of the sites having oligosaccharides;
- b) no O-linked glycosylation;
- c) faster plasma clearance time and an increased affinity for heparin both compared to human antithrombin III that is isolated from human plasma.

CLMS(9)

9. A glycosylated human antithrombin III, wherein a glycosylated human antithrombin III is made by the method of isolating the glycosylated human antithrombin III from milk, wherein the milk is collected from a transgenic goat, which goat expresses in mammary tissue a transgene encoding human antithrombin III and wherein the human antithrombin III is secreted into the milk produced by the transgenic goat; and wherein the glycosylated human antithrombin III has the following properties:
- a) monosaccharide glycosylation comprising N-acetylglucosamine and mannose;
 - b) no O-linked glycosylation;
 - c) faster plasma clearance time and an increased affinity for heparin both compared to human antithrombin III that is isolated from human plasma.

CLMS(10)

10. A glycosylated human antithrombin III, wherein a glycosylated human antithrombin III is made by the method of isolating the glycosylated human antithrombin III from milk, wherein the milk is collected from a transgenic goat, which goat expresses in mammary tissue a transgene encoding human antithrombin III and wherein the human antithrombin III is secreted into the milk produced by the transgenic goat; and wherein the glycosylated human antithrombin III has the following properties:
- a) monosaccharide glycosylation comprising N-acetylglucosamine, galactose and mannose;
 - b) no O-linked glycosylation;
 - c) faster plasma clearance time and an increased affinity for heparin both compared to human antithrombin III that is isolated from human plasma.

CLMS(11)

11. A glycosylated human antithrombin III, wherein a glycosylated human antithrombin III is made by the method of isolating the glycosylated human antithrombin III from milk, wherein the milk is collected from a

transgenic goat, which goat expresses in mammary tissue a transgene encoding human antithrombin III and wherein the human antithrombin III is secreted into the milk produced by the transgenic goat; and wherein the glycosylated human antithrombin III has the following properties:

- monosaccharide glycosylation comprising N-acetylglucosamine, N-acetylgalactosamine and mannose;
- no O-linked glycosylation;
- faster plasma clearance time and an increased affinity for heparin both compared to human antithrombin III that is isolated from human plasma.

CLMS(12)

12. The glycosylated human antithrombin III of any one of claim 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11, wherein the plasma clearance time of the transgenically produced antithrombin III is at least about 10 times faster than the plasma clearance time of the naturally occurring plasma antithrombin III.

CLMS(13)

13. The glycosylated human antithrombin III of any one of claim 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11, wherein the affinity for heparin of the transgenically produced antithrombin III results in at least about 1000 fold enhanced affinity for thrombin as compared to the naturally occurring plasma antithrombin III.

US PAT NO: 5,827,690 [IMAGE AVAILABLE] L15: 3 of 8

CLAIMS:

CLMS(1)

What is claimed is:

1. A high level expression method for providing a heterologous and assembled immunoglobulin, in the milk of a transgenic mammal comprising:

obtaining milk from a transgenic mammal having introduced into its germline a heterologous immunoglobulin protein-coding sequence operatively linked to a promoter sequence that results in the preferential expression of said protein-coding sequence in mammary gland epithelial cells, thereby providing said heterologous and assembled immunoglobulin in the milk of said mammal, wherein said

heterologous and assembled immunoglobulin is a functional configuration and is produced at level of at least about 1 mg/ml in the milk of said mammal.

CLMS(2)

2. The method of claim 1 wherein said mammal is selected from the group consisting of mice, sheep, and pigs.

CLMS(3)

3. The method of claim 1 wherein said promoter is selected from the group consisting of the beta lactoglobulin promoter, whey acid protein promoter, and the lactalbumin promoter.

CLMS(4)

4. The method of claim 1 wherein said immunoglobulin comprises heavy and light chains.

CLMS(5)

5. The method of claim 1 wherein said immunoglobulin is of human origin.

CLMS(6)

6. The method of claim 1 wherein said immunoglobulin is purified from the milk of said mammal.

CLMS(7)

7. The method of claim 1 wherein said promoter is the casein promoter.

CLMS(8)

8. A high level expression method for providing a heterologous and assembled immunoglobulin, in the milk of a transgenic goat comprising:

obtaining milk from a transgenic goat having introduced into its germline a heterologous immunoglobulin protein-coding sequence operatively linked to a promoter sequence that results in the preferential expression of said protein-coding sequence in mammary gland epithelial cells, thereby providing said heterologous and assembled immunoglobulin in the milk of said goat, wherein said heterologous and assembled immunoglobulin is a functional configuration and is produced at levels of at least about 1 mg/ml in the milk of said goat.

CLMS(9)

9. The method of claim 8 wherein said promoter is selected from the group consisting of the beta lactoglobulin promoter, whey acid protein promoter, and the lactalbumin promoter.

CLMS(10)

10. The method of claim 8 wherein said immunoglobulin comprises

heavy and light chains.

CLMS(11)

11. The method of claim 8 wherein said immunoglobulin is of human origin.

CLMS(12)

12. The method of claim 8 wherein said immunoglobulin is purified from the milk of said goat.

CLMS(13)

13. The method of claim 8 wherein said promoter is the casein promoter.

US PAT NO: 5,750,172 [IMAGE AVAILABLE] L15: 4 of 8

CLAIMS:

CLMS(1)

We claim:

1. Nonhuman mammal's milk comprising detectable levels of a recombinant polypeptide chain, wherein the recombinant polypeptide chain is produced by a nonhuman transgenic mammal whose somatic and germ cells contain an expression system comprising a DNA sequence coding for the recombinant polypeptide chain operably linked to a casein promoter and a signal peptide sequence, wherein the recombinant polypeptide chain is selected from the group consisting of coagulation factors VIII and IX, tissue plasminogen activator (TPA), urokinase, growth hormone, insulin, interferons, interleukins, peptide hormones, immunoglobulins and biologically active fragments thereof.

CLMS(2)

2. The milk of claim 1, wherein the non-human mammal is selected from the group consisting of sheep, goats, pigs and mice.

CLMS(3)

3. The milk of claim 1, wherein the expression system further comprises a 3' untranslated region downstream of the DNA sequence coding for the recombinant polypeptide.

CLMS(4)	4. The milk of claim 1, wherein the expression system further comprises a 5' untranslated region between said promoter and the DNA sequence coding for the signal peptide.	6. The DNA of claim 5 wherein the steroid hormone responsive element is a glucocorticoid responsive element.	polypeptide which is able to bind to biotin or biotin derivatives or analogues; and (c) DNA which, within the degeneracy of the genetic code, encodes the same polypeptide as either (a) or (b); and said second DNA coding for another protein, polypeptide, peptide or amino acid.
CLMS(5)	5. The milk of claim 1, wherein the promoter is an alpha.s1 casein promoter.	7. The DNA of claim 5 wherein the steroid hormone responsive element is an androgen responsive element.	CLMS(2)
US PAT NO: 5,688,677 [IMAGE AVAILABLE]	L15: 5 of 8	8. The DNA of claim 4 wherein the consensus sequence is selected from the group of nucleotide sequences consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:7.	2. A fused protein, according to claim 1, wherein the protein, polypeptide, peptide or amino acid encoded by the second DNA sequence is selected from the group consisting of human and animal interferons, human and animal growth hormones, antigens of FMDV, antigens of HBV, human insulin, human blood factors, tissue plasminogen activator and erythropoietin.
CLAIMS:			
CLMS(1)			CLMS(3)
We claim:			3. The fused protein according to claim 1, wherein the hybrid DNA further comprises a sufficient protein of a signal DNA sequence to cause secretion of the fused protein across the cell membrane of the transformed host.
1. Recombinant DNA comprising a nucleic acid sequence, the sequence including: a consensus sequence of at least one hormone responsive element, wherein the consensus sequence is mutated to render said hormone responsive element inactive; and a sequence encoding a membrane-associated protein.		9. The DNA of claim 8 wherein the consensus sequence is an androgen responsive element.	CLMS(4)
CLMS(2)		CLMS(10)	4. The fused protein according to claim 1 or 3, wherein the hybrid DNA further comprises a sufficient portion of a signal DNA sequence to cause maturation of the fused protein upon secretion of the fused protein across the cell membrane of the transformed host.
2. The DNA of claim 1 wherein the consensus sequence is located within the sequence encoding the membrane-associated protein.		CLMS(11)	CLMS(5)
CLMS(3)		11. A cystic fibrosis-affected cell comprising the DNA of claim 1.	5. A streptavidin, or portion thereof, which is able to bind biotin or biotin derivatives or analogues, said streptavidin containing the streptavidin signal sequence or a portion thereof at the amino terminus and being produced by a host transformed with a recombinant DNA molecule comprising a DNA coding for the streptavidin or portion thereof, the DNA selected from the group consisting of: (a) SA304 and SA307; (b) DNA sequences which hybridize to any of the foregoing DNA sequences in 6X and 0.1% SDS at 30 degree. C. overnight and which code for the polypeptide or portion thereof, which is able to bind to biotin or biotin derivatives or analogues; and (c) DNA which, within the degeneracy of the genetic code, encodes the same polypeptide as either (a) or (b).
3. The DNA of claim 1 wherein the membrane-associated protein is cystic fibrosis transmembrane conductance regulator.		US PAT NO: 5,272,254 [IMAGE AVAILABLE]	
CLMS(4)		8	
4. The DNA of claim 2 wherein the membrane-associated protein is cystic fibrosis transmembrane conductance regulator.		CLAIMS:	
CLMS(5)		CLMS(1)	
5. The DNA of claim 1 wherein the hormone responsive element is asteroid hormone responsive element.		We claim:	
CLMS(6)		1. A fused protein which is produced by a host transformed with a recombinant DNA molecule comprising a hybrid DNA, the hybrid DNA coding for the fused protein and comprising at least two DNAs joined end to end and in the same reading frame, the first DNA coding for streptavidin or a portion thereof, the streptavidin or portion thereof being able to bind to biotin or biotin derivatives or analogues and selected from the group consisting of: (a) SA304, SA307, SA324; (b) DNA which hybridizes to any of the foregoing DNA in 6XSSC and 0.1% SDS at 30.degree. C. overnight and which codes on expression for a	

<p>US PAT NO: 5,168,049 [IMAGE AVAILABLE]</p> <p>L15: 7 of 8</p> <p>CLAIMS:</p> <p>CLMS(1)</p> <p>We claim:</p> <p>1. An isolated DNA sequence coding for streptavidin or a portion thereof, said streptavidin or portion thereof being able to bind to biotin or biotin derivatives or analogues; selected from the group consisting of:</p> <p>(a) SA304, SA307, SA324;</p> <p>(b) DNA sequences which hybridize to any of the foregoing DNA sequences and which code on expression for a polypeptide which is able to bind to biotin or biotin derivatives or analogues; and</p> <p>(c) DNA sequences which code on expression for a polypeptide coded for on expression of any of the foregoing DNA sequences.</p> <p>CLMS(2)</p> <p>2. The DNA sequence according to claim 1, wherein said DNA sequence contains a sufficient portion of a signal DNA sequence to cause, upon expression of said DNA sequence, secretion of the polypeptide encoded by said DNA sequence across the cell membrane of a unicellular host transformed with said DNA sequence.</p> <p>CLMS(3)</p> <p>3. The DNA sequence according to claim 2, wherein said DNA sequence contains a sufficient portion of a signal DNA sequence to cause, upon expression of said DNA sequence, maturation of the polypeptide encoded by said DNA sequence upon secretion of said polypeptide across the cell membrane of a unicellular host transformed with said DNA sequence.</p> <p>CLMS(4)</p> <p>4. A recombinant DNA molecule comprising DNA selected from the group consisting of:</p> <p>(a) a DNA sequence coding for streptavidin or a portion thereof, said streptavidin or portion thereof being able to bind biotin or biotin derivatives or analogues; selected from the group consisting of:</p> <p>(1) SA304, SA307, SA324;</p> <p>(2) DNA sequences which hybridize to any of the foregoing DNA sequences and which code on expression for a polypeptide which is able to bind to biotin or biotin derivatives or analogues; and</p>	<p>(3) DNA sequences which code on expression for a polypeptide coded for on expression of any of the foregoing DNA sequences;</p> <p>(b) DNA comprising any of the foregoing DNA sequences and further comprising a sufficient portion of a signal DNA sequence to cause, upon expression of said DNA sequence, secretion of the polypeptide encoded by said DNA sequence across the cell membrane of a unicellular host transformed with said DNA sequence, and</p> <p>(c) DNA comprising any of the foregoing DNA sequences and further comprising a sufficient portion of a signal DNA sequence to cause, upon expression of said DNA sequence, maturation of the polypeptide encoded by said DNA sequence upon secretion of said polypeptide across the cell membrane of a unicellular host transformed with said DNA sequence.</p> <p>CLMS(5)</p> <p>5. The recombinant DNA molecule according to claim 4, wherein said DNA sequence is operatively linked to an expression control sequence in said molecule.</p> <p>CLMS(6)</p> <p>6. The recombinant DNA molecule according to claim 5 wherein the expression control sequence is selected from the group consisting of the E. coli lac system, the E. coli trp system, the E. coli beta-lac system, the TAC system, the TRC system, the major operator and promoter regions of bacteriophage lambda, the operator and promoter regions of filamentous single-stranded DNA phages, expression control sequences from Streptomyces or other gram positive bacteria, and combinations thereof.</p> <p>CLMS(7)</p> <p>7. The recombinant DNA molecule according to claim 6, selected from the group consisting of pSA304, pSA307 and pSA3721.</p> <p>CLMS(8)</p> <p>8. A unicellular host transformed with at least one recombinant DNA molecule according to claim 5, the expression control sequence in said recombinant DNA molecule being operatively linked to a DNA sequence in said host.</p> <p>CLMS(9)</p>	<p>9. The transformed host according to claim 8, selected from the group consisting of <i>S. lividans</i> (pSA3721), <i>E. coli</i> K12 (pSA304) and <i>E. coli</i> K12 (pSA307).</p> <p>CLMS(10)</p> <p>10. The transformed host according to claim 8, wherein the host transformed is selected from the group consisting of:</p> <p>(a) bacteria;</p> <p>(b) fungi;</p> <p>(c) plant hosts; and</p> <p>(d) animal hosts.</p> <p>CLMS(11)</p> <p>11. The transformed host according to claim 10, wherein the bacteria are selected from the group consisting of:</p> <p>(a) <i>Streptomyces</i>;</p> <p>(b) <i>Bacillus</i>; and</p> <p>(c) <i>E. coli</i>.</p> <p>CLMS(12)</p> <p>12. The transformed host according to claim 10, wherein the fungus is yeast.</p> <p>CLMS(13)</p> <p>13. The transformed host according to claim 10, wherein the animal host is human tissue cells.</p> <p>CLMS(14)</p> <p>14. A method for producing streptavidin or a portion thereof, said streptavidin or portion thereof being able to bind to biotin or biotin derivatives or analogues, comprising the step of culturing a host transformed with a recombinant DNA molecule according to claim 4.</p> <p>CLMS(15)</p> <p>15. The method according to claim 14, wherein the host transformed is selected from the group consisting of:</p> <p>(a) bacteria;</p> <p>(b) fungi;</p> <p>(c) plant hosts; and</p> <p>(d) animal hosts.</p> <p>CLMS(16)</p> <p>16. The method according to claim 15, wherein the bacteria are selected from the group consisting of:</p>
--	---	--

- (a) *Streptomyces*,
(b) *Bacillus*; and
(c) *E. coli*.
- CLMS(17)
17. The method according to claim 15, wherein the fungus is yeast.
- CLMS(18)
18. The method according to claim 15, wherein the animal host is human tissue cells.
- CLMS(19)
19. A hybrid DNA sequence coding for a fused protein, comprising at least two DNA sequences joined end to end and in the same reading frame, said first DNA sequence coding for streptavidin or a portion thereof, said streptavidin or portion thereof being able to bind to biotin or biotin derivatives or analogues, and selected from the group consisting of:
(a) SA304, SA307, SA324;
(b) DNA sequences which hybridize to any of the foregoing DNA sequences, and which code on expression for a polypeptide which is able to bind to biotin or biotin derivatives or analogues; and
(c) DNA sequences which code on expression for a polypeptide coded for on expression of any of the foregoing DNA sequences; and said second DNA sequence coding for another protein, polypeptide, peptide or amino acid.
- CLMS(20)
20. The hybrid DNA sequence according to claim 19, further comprising a sufficient portion of a signal DNA sequence to cause, upon expression of said DNA sequence, secretion of the fused protein across the cell membrane of a unicellular host transformed with said DNA sequence.
- CLMS(21)
21. The hybrid DNA sequence according to claim 20, further comprising a sufficient portion of a signal DNA sequence to cause, upon expression of said DNA sequence, maturation of the fused protein upon secretion of said polypeptide across the cell membrane of a unicellular host transformed with said hybrid DNA sequence.
- CLMS(22)
22. A hybrid DNA sequence according to claim 19, 20 or 21, in which said second DNA sequence codes for tissue plasminogen activator, said hybrid DNA sequence being selected from the group consisting of:
(a) SAT9724; and
(b) SAT7021.
- CLMS(23)
23. The hybrid DNA sequence according to claim 19, 20 or 21, wherein the second DNA sequence encodes polypeptides selected from the group consisting of human interferons, human growth hormone, animal growth hormones, antigens of FMDV, human insulin, and tissue plasminogen activator.
- CLMS(24)
24. A recombinant DNA molecule comprising a hybrid DNA sequence according to claim 19, 20 or 21, wherein said hybrid DNA sequence is operatively linked to an expression control sequence in said molecule.
- CLMS(25)
25. The recombinant DNA molecule according to claim 24, wherein said hybrid DNA sequence contains a second DNA sequence encoding polypeptides selected from the group consisting of human interferons, human growth hormone, animal growth hormones, antigens of FMDV, human insulin, and tissue plasminogen activator.
- CLMS(26)
26. The recombinant DNA molecule according to claim 24, wherein the expression control sequence is selected from the group consisting of the *E. coli* lac system, the *E. coli* trp system, the *E. coli* beta-lac system, the TAC system, the TRC system, the major operator and promoter regions of bacteriophage lambda, the operator and promoter regions of filamentous single-stranded DNA phages, expression control sequences from *Streptomyces* or other gram positive bacteria, and combinations thereof.
- CLMS(27)
27. A recombinant DNA molecule according to claim 26, selected from the group consisting of:
(a) *Streptomyces*,
(b) *Bacillus*; and
(c) *E. coli*.
- CLMS(28)
28. A method for producing a fused protein comprising the step of culturing a host transformed with a recombinant DNA molecule of claim 24.
- CLMS(29)
29. The method of claim 28, wherein the hybrid DNA sequence contains a second DNA sequence, said second DNA sequence encoding polypeptides selected from the group consisting of human interferons, human growth hormone, animal growth hormones, antigens of FMDV, human insulin, and tissue plasminogen activator.
- CLMS(30)
30. A unicellular host transformed with at least one recombinant DNA molecule according to claim 24 the expression control sequence in said DNA molecule being operatively linked to a DNA sequence in said host.
- CLMS(31)
31. A transformed host according to claim 30, selected from the group consisting of *E. coli* HB101 (pSAT9724) and *S. lividans* (pSAT7026).
- CLMS(32)
32. The transformed host according to claim 30, wherein the host transformed is selected from the group consisting of:
(a) bacteria;
(b) fungi;
(c) plant hosts; and
(d) animal hosts.
- CLMS(33)
33. The transformed host according to claim 32, wherein the bacteria are selected from the group consisting of:
(a) *Streptomyces*;
(b) *Bacillus*; and
(c) *E. coli*.
- CLMS(34)
34. The transformed host according to claim 32, wherein the fungus is yeast.
- CLMS(35)

35. The transformed host according to claim 32, wherein the animal host is human tissue cells.

US PAT NO: 4,873,316 [IMAGE AVAILABLE]

L15: 8 of 8

CLAIMS:

CLMS(1)

We claim:

1. A process for the production and secretion into mammal's milk of an exogenous recombinant protein comprising the steps of:

a. producing milk in a transgenic mammal characterized by an expression system comprising a casein promoter operatively linked to an exogenous DNA sequence coding for the recombinant protein through a DNA sequence coding for a signal peptide effective in secreting and maturing the recombinant protein in mammary tissue;

b. collecting the milk; and

c. isolating the exogenous recombinant protein from the milk.

CLMS(2)

2. The process according to claim 1, wherein said expression system also includes a 3' untranslated region downstream of the DNA sequence coding for the recombinant protein.

CLMS(3)

3. The process according to claim 1, wherein said expression system also includes a 5' untranslated region between said promoter and the DNA sequence coding for the signal peptide.

=> d his

(FILE 'USPAT' ENTERED AT 16:49:49 ON 05 AUG 1999)

L1 107 S MAMMARY(10A)(IMMUNOGLOB? OR ANTIBOD?)

L2 80 S L1 AND MILK

L3 69 S L2 AND EXPRESS?

L4 42 S L3 AND (VECTOR# OR CONSTRUCT# OR PLASMID#)

L5 296 S MAMMARY(SA)(PROMOTER#)

L6 1 S L5(10A)(ANTIBOD? OR IMMUNOGLOB?)

L7 59 S L5 AND (CASEIN# OR WHEY ACID OR LACTALBUMIN OR LACTOGLOB

UL1

L8

L9

L10

L11

L12

L13

L14

L15

=> log y

U.S. Patent & Trademark Office LOGOFF AT 17:21:38 ON 05 AUG 1999

55 S L7 AND (IMMUNOGLOB? OR ANTIBOD?)

44 S L8 AND MILK

359 S MAMMARY(10A)(PROMOTER#)

40 S L10(10A)(MILK)

0 S L11(10A)(ANTIBOD? OR IMMUNOGLOB?)

37 S L10(P)(ANTIBOD? OR IMMUNOGLOB?)

20 S L13(P)(MILK)

E MEADE, HARRY

E MEADE, HARRY/IN

8 S E3-E4

transgenic non-human mammal. An especially useful catalytic entity is human glycosyltransferases which produce oligosaccharides and glycoconjugates. Specifically exemplified, is the production of 2'-fucosyl-lactose in the milk of transgenic mice which contain and express a transgene encoding .alpha.-1,2-fucosyltransferase operatively linked to a mammary gland specific **promoter**. A method of obtaining humanized milk is disclosed. The method comprises the steps of (a) inserting into the genome of a non-human mammal a heterologous gene encoding the production of a human catalytic entity wherein said catalytic entity produces a secondary gene product in the milk of said non-human mammal; and (b) milking said non-human mammal. The humanized milk may be used in the preparation of an enteral nutritional product useful in the nutritive maintenance of an animal.

5. 5,888,774, Mar. 30, 1999, Recombinant DNA molecules and expression **vectors** for erythropoietin; Genevieve Delcuve, 435/69 6, 320 1, 455, 456 [IMAGE AVAILABLE]

US PAT NO: 5,888,774 [IMAGE AVAILABLE] L4: 5 of 16

ABSTRACT:
A recombinant DNA molecule adapted for transfection of a host cell comprising a nucleic acid molecule encoding mammalian erythropoietin, an expression control sequence operatively linked thereto and at least one SAR element. The invention also relates to expression **vectors** having the recombinant DNA molecule and to mammalian cells transformed with the expression **vector**. The mammalian cells lack multiple copies of an amplified amplification gene and are capable of expressing recombinant EPO in vitro at levels of at least 1,500 u/10 sup.6 cells in 24 hours. The invention further relates to a method of expressing recombinant mammalian erythropoietin using the expression **vectors** and to a transgenic non-human animal or embryo whose germ cells and somatic cells contain a DNA **construct** having the recombinant DNA molecule of the invention.

6. 5,833,982, Nov. 10, 1998, Modified factor VII; Kathleen L. Berkner, et al., 424/94 64; 435/212, 226; 514/12, 530/384 [IMAGE AVAILABLE]

US PAT NO: 5,833,982 [IMAGE AVAILABLE] L4: 6 of 16

ABSTRACT:

cells; Russel E. Kaufman, et al., 435/69 1, 252 3, 320 1, 325; 530/350; 536/23 1, 23 5 [IMAGE AVAILABLE]

US PAT NO: 5,912,142 [IMAGE AVAILABLE] L4: 2 of 16

ABSTRACT:
The present invention relates, in general, to a cancer-related protein and to a nucleic acid sequence encoding same. In particular, the invention relates to a protein over expressed in certain neoplastic cells, including breast and ovarian cancer cells, to its encoding sequence, and to diagnostic and treatment methodologies based on same.

3. 5,892,070, Apr. 6, 1999, Transgenic non-human mammals producing oligosaccharides and glycoconjugates; Pedro Antonio Prieto, et al., 800/14; 435/69 1, 800/15, 16, 17, 18 [IMAGE AVAILABLE]

US PAT NO: 5,892,070 [IMAGE AVAILABLE] L4: 3 of 16

ABSTRACT:
The invention relates to transgenic non-human mammals characterized in that the genome of said mammals contain at least one heterologous gene encoding for the production of heterologous catalytic entity selected from the group consisting of enzymes and antibodies, and wherein said catalytic entity produces a second heterologous product in the milk of said mammal. Especially useful in the practice of the invention are human glycosyltransferases and transgenic sheep, goats and cows. The heterologous product includes oligosaccharides and glycoconjugates. Specifically exemplified, is the production of 2'-fucosyl-lactose in the milk of transgenic mice which contain and express a transgene encoding .alpha.-1,2-fucosyltransferase operatively linked to a mammary gland specific **promoter**.

4. 5,891,698, Apr. 6, 1999, Oligosaccharides and glycoproteins produced in milk of transgenic non-human mammals; Pedro Antonio Prieto, et al., 800/7; 435/100; 800/14, 15, 16, 17, 18, 25 [IMAGE AVAILABLE]

US PAT NO: 5,891,698 [IMAGE AVAILABLE] L4: 4 of 16

ABSTRACT:
The invention relates to humanized milk. The milk is produced by a non-human transgenic mammal wherein the genome of said transgenic non-human mammal contains at least one heterologous gene encoding for a human catalytic entity and wherein the catalytic entity produces oligosaccharides and glycoconjugates that are present in the milk of said

FILE 'USPAT' ENTERED AT 17:22:03 ON 04 AUG 1999

* U. S. P A T E N T T E X T F I L E *
*
* THE WEEKLY PATENT TEXT AND IMAGE DATA IS CURRENT *
* THROUGH AUGUST 3,1999 *
* *

=> s immunoglob?
L1 11569 IMMUNOGLOB?
=> s I1 and whey acidic protein
4328 WHEY
98021 ACIDIC
73025 PROTEIN
45 WHEY ACIDIC PROTEIN
(WHEY(W)ACIDIC(W)PROTEIN)
L2 18 L1 AND WHEY ACIDIC PROTEIN
=> s I2 and promoter#
36024 PROMOTER#
L3 16 L2 AND PROMOTER#
=> s I3 and (construct# or vector# or plasmid#)
96404 CONSTRUCT#
77317 VECTOR#
16546 PLASMID#
L4 16 L3 AND (CONSTRUCT# OR VECTOR# OR PLASMID#)
=> d I - cit ab
1. 5,919,997, Jul. 6, 1999, Transgenic mice having modified cell-cycle regulation; David H. Beach, et al., 800/18; 435/91 2, 320 1, 325, 455, 463, 467; 800/3, 22, 25 [IMAGE AVAILABLE]

US PAT NO: 5,919,997 [IMAGE AVAILABLE] L4: 1 of 16

ABSTRACT:
The present invention relates to transgenic mice in which the biological function of at least one cell cycle regulatory proteins of the INK4 family is altered.

2. 5,912,142, Jun. 15, 1999, Gene product over expressed in cancer

The catalytic active site of Factor VII is modified to produce a compound which effectively interrupts the blood coagulation cascade. The modifications render Factor VIIa substantially unable to activate plasma Factors X or IX. Pharmaceutical compositions of the modified Factor VII are used to treat a variety of coagulation-related disorders, including platelet deposition, vascular thrombosis, ischemic reperfusion, acute closure of a coronary artery, vascular restenosis secondary to balloon angioplasty, endarterectomy, reductive atherectomy, stent placement, laser therapy or rotoablation.

7. 5,817,788, Oct. 6, 1998, Modified factor VII; Kathleen L. Berkner, et al., 536/23 2; 435/212, 226, 325 [IMAGE AVAILABLE] L4: 7 of 16

US PAT NO: 5,817,788 [IMAGE AVAILABLE] L4: 7 of 16

ABSTRACT:
The catalytic active site of Factor VII is modified to produce a compound which effectively interrupts the blood coagulation cascade. The modifications render Factor VIIa substantially unable to activate plasma Factors X or IX. Pharmaceutical compositions of the modified Factor VII are used to treat a variety of coagulation-related disorders.

8. 5,788,965, Aug. 4, 1998, Modified factor VII; Kathleen L. Berkner, et al., 424/94.64; 435/212, 226; 514/12, 822; 530/384 [IMAGE AVAILABLE] L4: 8 of 16

US PAT NO: 5,788,965 [IMAGE AVAILABLE] L4: 8 of 16

ABSTRACT:
The catalytic active site of Factor VII is modified to produce a compound which effectively interrupts the blood coagulation cascade. The modifications render Factor VIIa substantially unable to activate plasma Factors X or IX. Pharmaceutical compositions of the modified Factor VII are used to treat a variety of coagulation-related disorders.

9. 5,776,773, Jul. 7, 1998, Yeast artificial chromosomes and their use in the control of gene expression; Marianne Bruggemann, 435/325, 320.1, 449 [IMAGE AVAILABLE] L4: 9 of 16

US PAT NO: 5,776,773 [IMAGE AVAILABLE] L4: 9 of 16

ABSTRACT:

Embryonic stem cells that are essentially free of yeast DNA are prepared from suitably marked yeast artificial chromosomes and used to transfer DNA segments of considerable size into organisms.

10. 5,750,176, May 12, 1998, Transgenic non-human mammal milk comprising 2'-fucosyl-lactose; Pedro Antonio Prieto, et al., 426/580; 424/530; 426/556, 587, 588; 530/832; 800/7 [IMAGE AVAILABLE] L4: 10 of 16

US PAT NO: 5,750,176 [IMAGE AVAILABLE] L4: 10 of 16

ABSTRACT:
The invention relates to the milk of a transgenic non-human mammal. The milk is characterized in that it contains heterologous components produced as the secondary gene products of a heterologous gene contained in the genome of the transgenic non-human mammal. The heterologous gene encodes a heterologous catalytic entity such as a human enzyme selected from the group consisting of glycosyltransferases, phosphorylases, hydroxylases, peptidases and sulfotransferases. Especially useful in the practice of the invention are human glycosyltransferases. The desired heterologous components include oligosaccharides, glycoconjugates. Specifically exemplified, is the production of 2'-fucosyl-lactose in the milk of transgenic mice which contain and express a transgene encoding alpha.-1,2-fucosyltransferase operatively linked to a mammary gland specific **promoter**. The oligosaccharides and glycoconjugates may be isolated from the milk of the transgenic mammals and used in the preparation of pharmaceuticals, diagnostic kits, nutritional products and the like. The whole transgenic milk may also be used to formulate nutritional products that provide special advantages. The transgenic milk may also be used in the production of specialized enteral nutritional products.

11. 5,700,671, Dec. 23, 1997, Methods of making transgenic animals producing oligosaccharides and glycoproteins; Pedro Antonio Prieto, et al., 800/25; 435/6, 69.1, 193 [IMAGE AVAILABLE] L4: 11 of 16

US PAT NO: 5,700,671 [IMAGE AVAILABLE] L4: 11 of 16

ABSTRACT:
The invention relates to transgenic non-human mammals characterized in that the genome of said mammals contain at least one heterologous gene encoding for the production of heterologous catalytic entity selected from the group consisting of enzymes and antibodies, and wherein said

catalytic entity produces a second heterologous product in the milk of said mammal. Especially useful in the practice of the invention are human glycosyltransferases and transgenic sheep, goats and cows. The heterologous product includes oligosaccharides and glycoconjugates.

12. 5,648,243, Jul. 15, 1997, Human serum albumin expression **construct**; David R. Hurwitz, et al., 435/69.6, 320.1; 536/23.1, 23.5, 24.1, 24.2 [IMAGE AVAILABLE] L4: 12 of 16

US PAT NO: 5,648,243 [IMAGE AVAILABLE] L4: 12 of 16

ABSTRACT:
The present invention provides DNA **constructs** comprising a **promoter** DNA sequence and a DNA sequence coding for human serum albumin. In one embodiment the human serum albumin sequence comprises at least one, but not all, of the introns in the naturally occurring gene encoding for the HSA protein. In another embodiment the DNA **constructs** comprise a 5' regulatory sequence which directs the expression and secretion of HSA protein in the milk of a transgenic animal. Preferably, the **promoter** gene is a milk protein **promoter** sequence such as beta.-lactoglobulin. The present invention also provides transgenic animals which secrete HSA in the milk of lactating females. The present invention also provides **vectors** comprising the **constructs** of the present invention.

13. 5,476,995, Dec. 19, 1995, Peptide production; Anthony J. Clark, et al., 800/16; 435/69.1, 317.1, 320.1 [IMAGE AVAILABLE] L4: 13 of 16

US PAT NO: 5,476,995 [IMAGE AVAILABLE] L4: 13 of 16

ABSTRACT:
A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for polypeptide into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta.-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.

14. 5,366,894, Nov. 22, 1994, Peptide production; Anthony J. Clark, et al., 435/320.1, 69.1, 325 [IMAGE AVAILABLE]	ABSTRACT: A transgenic mouse offspring produced by the mating of a first transgenic mouse carrying a transresponder transgene whose expression is regulated by a viral gene product of HSV-1 and a second transgenic mouse carrying a transactivator transgene. A process for expressing a gene of interest which comprises the mating of a first transgenic mouse carrying a transresponder transgene whose expression is regulated by a viral gene product of HSV-1 and a second transgenic mouse carrying a transactivator transgene.	Chimeric and humanized IL4 MABs derived from high affinity MABs, pharmaceutical compositions containing same, and methods of treatment are provided.
US PAT NO: 5,366,894 [IMAGE AVAILABLE] L4: 14 of 16		2. 5,914,110, Jun. 22, 1999, Recombinant IL4 antibodies useful in treatment of IL4 mediated disorders; Stephen D. Holmes, et al., 424/133.1, 141.1, 152.1; 435/7.1, 70.21, 326, 328, 335; 530/350, 387.3; 388.15, 388.23, 391.1 [IMAGE AVAILABLE]
ABSTRACT: A method of producing a substance comprising a peptide, involves incorporating a DNA sequence coding for the peptide into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The substance may be an (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The substance will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.	=> s immunoglob? and casein 11569 IMMUNOGLOB? 17607 CASEIN L5 908 IMMUNOGLOB? AND CASEIN => s l5 and promoter# 36024 PROMOTER# L6 410 L5 AND PROMOTER# => s l6 and (construct# or vector# or plasmid#) 96404 CONSTRUCT# 77317 VECTOR# 16546 PLASMID# L7 387 L6 AND (CONSTRUCT# OR VECTOR# OR PLASMID#) => s immunoglob? and casein promoter 11569 IMMUNOGLOB? 17607 CASEIN 27862 PROMOTER 34 CASEIN PROMOTER (CASEIN(W)PROMOTER) L8 20 IMMUNOGLOB? AND CASEIN PROMOTER => d l - cit ab	US PAT NO: 5,914,110 [IMAGE AVAILABLE] L8: 2 of 20
15. 5,322,775, Jun. 21, 1994, Peptide production; Anthony J. Clark, et al., 435/69.1, 69.6, 69.7, 317.1, 320.1; 530/412 [IMAGE AVAILABLE]	ABSTRACT: A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for a polypeptide chain of said compound into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.	ABSTRACT: Chimeric and humanized IL4 MABs derived from high affinity MABs, pharmaceutical compositions containing same, and methods of treatment are provided.
US PAT NO: 5,322,775 [IMAGE AVAILABLE] L4: 15 of 16		3. 5,877,010, Mar. 2, 1999, Thymidine kinase mutants; Lawrence A. Loeb, et al., 435/320.1, 243, 325; 536/23.2, 23.5, 23.72, 24.1 [IMAGE AVAILABLE]
ABSTRACT: A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for a polypeptide chain of said compound into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.	The present invention provides isolated nucleic acid molecules encoding a Herpesviridae thymidine kinase enzyme comprising one or more mutations, at least one of the mutations encoding an amino acid substitution upstream from a DRH nucleoside binding site which increases a biological activity of the thymidine kinase, as compared to unmutated thymidine kinase. Within another aspect, one of the mutations is an amino acid substitution within a DRH nucleoside binding site which increases a biological activity of the thymidine kinase, as compared to unmutated thymidine kinase. Also provided are vectors suitable for expressing such DNA molecules, as well as methods for utilizing such vectors.	US PAT NO: 5,877,010 [IMAGE AVAILABLE] L8: 3 of 20
16. 5,221,778, Jun. 22, 1993, Multiplex gene regulation; Guerard W. Byrne, et al., 800/4, 424/231.1, 435/193, 317.1, 948, 800/18, 22 [IMAGE AVAILABLE]	ABSTRACT: A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for a polypeptide chain of said compound into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.	ABSTRACT: Chimeric, humanized and other IL-5 mAbs, derived from high affinity neutralizing mAbs, pharmaceutical compositions containing same, methods of treatment and diagnostics are provided.
US PAT NO: 5,221,778 [IMAGE AVAILABLE] L4: 16 of 16		4. 5,851,525, Dec. 22, 1998, Recombinant IL-5 antagonists useful in treatment of IL-5 mediated disorders; Robert S. Ames, Jr., et al., 424/145.1, 152.1, 158.1, 172.1; 530/387.1, 387.3, 388.23 [IMAGE AVAILABLE]
	1. 5,928,904, Jul. 27, 1999, DNA encoding recombinant IL4 antibodies useful in treatment of IL4 mediated disorders; Stephen D. Holmes, et al., 435/69.6, 70.21, 71.1, 320.1, 326, 328, 335; 530/300, 350, 387.3, 388.23; 536/23.5, 23.53 [IMAGE AVAILABLE]	US PAT NO: 5,851,525 [IMAGE AVAILABLE] L8: 4 of 20
	US PAT NO: 5,928,904 [IMAGE AVAILABLE] L8: 1 of 20	
	ABSTRACT:	

THIS PAGE BLANK (USPTO)

5. 5,849,992, Dec. 15, 1998, Transgenic production of antibodies in milk; Harry Meade, et al., 800/14, 7, 15, 16, 17, 18 [IMAGE AVAILABLE]		hormones and **immunoglobulins**, in mammals' milk. Particularly, this invention relates to an expression system which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.	
US PAT NO: 5,849,992 [IMAGE AVAILABLE]	L8: 5 of 20		
ABSTRACT: A method for the production of monoclonal antibodies in mammal's milk through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.			
6. 5,827,690, Oct. 27, 1998, Transgenic production of antibodies in milk; Harry Meade, et al., 800/7; 530/867 [IMAGE AVAILABLE]			
US PAT NO: 5,827,690 [IMAGE AVAILABLE]	L8: 6 of 20		
ABSTRACT: A method for the production of monoclonal antibodies in mammal's milk, through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.			
7. 5,783,184, Jul. 21, 1998, Method for treatment and diagnosis of IL-5 mediated disorders; Edward Robert Appelbaum, et al., 424/130.1, 133.1, 141.1, 145.1; 435/7.1; 530/388.1, 388.23 [IMAGE AVAILABLE]			
US PAT NO: 5,783,184 [IMAGE AVAILABLE]	L8: 7 of 20		
ABSTRACT: The present invention relates to treatment and diagnosis of conditions mediated by IL-5 and excess eosinophil production, and more specifically to mAbs and other altered antibodies such as Fabs, chimeric, human and humanized antibodies that do not block binding of human IL-5 to the alpha.-chain of the human IL-5 receptor.			
8. 5,750,172, May 12, 1998, Transgenic non human mammal milk; Harry Meade, et al., 426/580; 435/69.1, 69.4, 69.51, 69.52, 69.6, 183, 215; 800/7 [IMAGE AVAILABLE]			
US PAT NO: 5,750,172 [IMAGE AVAILABLE]	L8: 8 of 20		
ABSTRACT: This invention relates to the production of recombinant proteins, such as coagulation factors VIII and IX, tissue plasminogen activator (TPA), urokinase, growth hormone, insulin, interferons, interleukins, peptide			
9. 5,741,957, Apr. 21, 1998, Transgenic bovine; Herman A. Deboer, et al., 800/7; 435/69.1; 800/15, 25 [IMAGE AVAILABLE]			
US PAT NO: 5,741,957 [IMAGE AVAILABLE]	L8: 9 of 20		
ABSTRACT: A transgenic bovine is disclosed whose somatic and germ cells contain a transgene, wherein the transgene comprising a mammary gland specific promoter, a mammary gland specific enhancer, a DNA sequence encoding a signal sequence functional in bovine mammary gland secretory cells and a DNA sequence encoding a heterologous polypeptide of interest wherein the transgenic bovine expresses the transgene such that the polypeptide of interest is detectable in milk produced by the transgenic bovine.			
10. 5,736,388, Apr. 7, 1998, Bacteriophage-mediated gene transfer systems capable of transfecting eukaryotic cells; Sunil Chada, et al., 435/320.1; 424/93.6; 435/235.1; 514/44 [IMAGE AVAILABLE]			
US PAT NO: 5,736,388 [IMAGE AVAILABLE]	L8: 10 of 20		
ABSTRACT: Lamboid bacteriophage capable of specifically interacting with and delivering nucleic acid molecules to eukaryotic cells are disclosed. Such bacteriophage-derived gene transfer systems target one or more specific receptors on eukaryotic cells, for instance by incorporating mutant tail fiber proteins or by incorporating known ligands for specific eukaryotic receptors into lambda phage. Also disclosed are methods for identifying and producing modified bacteriophage tail fiber polypeptides capable of specifically interacting with eukaryotic transmembrane proteins. Methods of treating diseases using such gene transfer systems are also disclosed.			
11. 5,721,367, Feb. 24, 1998, Homologous recombination in mammalian cells; Robert M. Kay, et al., 800/18; 435/69.1, 69.7, 463, 465; 800/15 [IMAGE AVAILABLE]			
hormones and **immunoglobulins**, in mammals' milk. Particularly, this invention relates to an expression system which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.			
US PAT NO: 5,721,367 [IMAGE AVAILABLE]	L8: 11 of 20		
ABSTRACT: The invention relates to methods for intracellularly producing DNA segments by homologous recombination of smaller overlapping DNA fragments and transgenic mammalian cells and transgenic non-human mammals produced by such methods.			
12. 5,693,323, Dec. 2, 1997, Recombinant IL-5 antagonists useful in treatment of IL-5 mediated disorders; Robert S. Anes, Jr., et al., 424/145.1; 435/328, 335; 530/387.3, 388.23 [IMAGE AVAILABLE]			
US PAT NO: 5,693,323 [IMAGE AVAILABLE]	L8: 12 of 20		
ABSTRACT: Chimeric, humanized and other IL-5 mAbs, derived from high affinity neutralizing mAbs, pharmaceutical compositions containing same, methods of treatment and diagnostics are provided.			
13. 5,688,677, Nov. 18, 1997, Deoxyribonucleic acids containing inactivated hormone responsive elements; Karl M. Ebert, et al., 536/23.5, 24.1 [IMAGE AVAILABLE]			
US PAT NO: 5,688,677 [IMAGE AVAILABLE]	L8: 13 of 20		
ABSTRACT: A DNA comprising at least one inactivated hormone responsive element and a nucleic acid sequence encoding a membrane-associated protein is described. Therapeutic compositions and cells including the DNA are also described. Other aspects of the invention include methods of treating subjects having cystic fibrosis which include administering an effective amount of the DNA to subjects having cystic fibrosis such that functional cystic fibrosis transmembrane conductance regulator is produced by the subject at a level which is not detrimental to the subject. The present invention also pertains to a method of introducing the DNA into a cell such that the membrane-associated protein is produced at a level which is not detrimental to the cell and cells produced by this method. Still other aspects of the invention include a method of assaying DNA for the presence or absence of a hormone responsive element in a species in which the hormone responsive element is functional and a method of selectively breeding female transgenic mammals which produce a protein of			

interest.

14. 5,683,892, Nov. 4, 1997, DNA encoding recombinant IL-5 antagonists
useful in treatment of IL-5 mediated disorders; Robert S. Ames, Jr., et
al., 435/69.1, 69.3, 70.21, 252.3, 320.1, 328; 536/23.53 [IMAGE
AVAILABLE]

US PAT NO: 5,683,892 [IMAGE AVAILABLE] L8: 14 of
20

ABSTRACT:
DNA encoding chimeric, humanized and other IL-5 mAbs, derived
from high
affinity neutralizing mAbs, pharmaceutical compositions containing
same,
methods of treatment and diagnostics are provided.

15. 5,681,746, Oct. 28, 1997, Retroviral delivery of full length factor
VIII; Mordchai Bodner, et al., 435/350, 320.1, 366, 371; 536/23.5
[IMAGE
AVAILABLE]

US PAT NO: 5,681,746 [IMAGE AVAILABLE] L8: 15 of
20

ABSTRACT:
Retroviral vectors for directing expression of full length factor VIII in
transduced host cells, plasmids encoding the same, and host cells
transformed, transfected, or transduced therewith are disclosed. Also
disclosed are retroviral particles comprising such retroviral vectors, as
are methods for making such particles in suitable packaging cells.
Retroviral particles so produced may be amphotropic, ecotropic,
polytropic, or xenotropic; alternatively, they may comprise chimeric or
hybrid envelope proteins to alter host range. Also described are
retroviral particles comprising retroviral vectors for directing full
length factor VIII expression which are complement resistant.
Pharmaceutical compositions comprising retroviral particles of the
invention are also disclosed, as are methods of treating mammals,
particularly humans, afflicted with hemophilia.

16. 5,633,076, May 27, 1997, Method of producing a transgenic
bovine or
transgenic bovine embryo; Herman A. DeBoer, et al., 800/25 [IMAGE
AVAILABLE]

US PAT NO: 5,633,076 [IMAGE AVAILABLE] L8: 16 of
20

ABSTRACT:
A method is disclosed for the production of a transgenic bovine or a
transgenic bovine embryo comprising obtaining an ovum from bovine
ovaries, maturing the ovum in vitro, fertilizing the mature ovum or ova
in vitro to form a zygote, introducing a transgene into the zygote in
vitro and maturing the zygote to a preimplantation stage embryo in
vitro.
To produce the transgenic bovine, the embryo is transplanted into a

recipient female bovine, wherein the female bovine gestates the
embryo to
produce a transgenic bovine.

17. 5,612,205, Mar. 18, 1997, Homologous recombination in
mammalian
cells; Robert M. Kay, et al., 435/463, 465 [IMAGE AVAILABLE]

US PAT NO: 5,612,205 [IMAGE AVAILABLE] L8: 17 of
20

ABSTRACT:
The invention relates to methods for intracellularly producing DNA
segments by homologous recombination of smaller overlapping DNA
fragments
and transgenic mammalian cells and transgenic non-human mammals
produced
by such methods.

18. 5,525,708, Jun. 11, 1996, Covalent dimer of kit ligand; Karl H.
Nocka, et al., 530/409, 351, 399, 417 [IMAGE AVAILABLE]

US PAT NO: 5,525,708 [IMAGE AVAILABLE] L8: 18 of
20

ABSTRACT:
A modified form of KL, the ligand for the c-Kit proto-oncogene, has
been
prepared wherein the protein is stabilized by an intermolecular
covalent
linkage. The protein can be prepared by expression of a recombinant
protein which is dissolved in denaturant and refolded under conditions
resulting in a disulfide linked dimer. Examples demonstrate the
purification and characterization of this disulfide-linked cysteine dimer
kit ligand (KL-CD) which contains at least one intermolecular
disulfide
bond and has at least ten-fold greater activity in promoting cell
proliferation than native, non-covalently linked KL, as measured in in
vitro assays.

19. 5,268,275, Dec. 7, 1993, Vitamin K-dependent carboxylase;
Darrel W.
Stafford, et al., 435/69.1, 69.6, 232, 252.3, 320.1, 352, 354, 358, 366;
536/23.2 [IMAGE AVAILABLE]

US PAT NO: 5,268,275 [IMAGE AVAILABLE] L8: 19 of
20

ABSTRACT:
Isolated DNA encoding a vitamin K dependent carboxylase is
disclosed. The
carboxylase is selected from the group consisting of: (a) isolated DNA
which encodes bovine or human vitamin K dependent carboxylase; (b)
isolated DNA which hybridizes to isolated DNA of (a) above and
which
encodes a vitamin K dependent carboxylase; and (c) isolated DNA

differing
from the isolated DNAs of (a) and (b) above in nucleotide sequence
due to
the degeneracy of the genetic code, and which encodes a vitamin K
dependent carboxylase. Also disclosed are vectors and host cells
containing the aforesaid DNA, methods of using the same, and
purified
protein coded for by the aforesaid DNA.

20. 4,873,316, Oct. 10, 1989, Isolation of exogenous recombinant
proteins from the milk of transgenic mammals; Harry Meade, et al.,
800/7;
435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8; 530/360, 361, 416, 417, 418,
832;
833; 536/23.1, 23.4, 23.5; 800/18 [IMAGE AVAILABLE]

US PAT NO: 4,873,316 [IMAGE AVAILABLE] L8: 20 of
20

ABSTRACT:
This invention relates to the production of recombinant proteins in
mammals' milk. Particularly, this invention relates to an expression
system comprising the mammal's **cascin** **promoter** which
when
transgenically incorporated into a mammal permits the female species
of
that mammal to produce the desired recombinant protein in or along
with
its milk. This invention also relates to the transgenic mammal that
produces the desired recombinant product in its milk.

=> d his

(FILE 'USPAT' ENTERED AT 17:22:03 ON 04 AUG 1999)
L1 11569 S IMMUNOGLOB?
L2 18 S L1 AND WHEY ACIDIC PROTEIN
L3 16 S L2 AND PROMOTER#
L4 16 S L3 AND (CONSTRUCT# OR VECTOR# OR
PLASMID#)
L5 908 S IMMUNOGLOB? AND CASEIN
L6 410 S L5 AND PROMOTER#
L7 387 S L6 AND (CONSTRUCT# OR VECTOR# OR
PLASMID#)
L8 20 S IMMUNOGLOB? AND CASEIN PROMOTER

=> s immunoglob? and lactoglobulin

11569 IMMUNOGLOB?
635 LACTOGLOBULIN
L9 208 IMMUNOGLOB? AND LACTOGLOBULIN

=> s immunoglob? and lactoglobulin promoter

11569 IMMUNOGLOB?
635 LACTOGLOBULIN
27862 PROMOTER

<p>15 LACTOGLOBULIN PROMOTER (LACTOGLOBULIN(W)/PROMOTER) L10 8 IMMUNOGLOB? AND LACTOGLOBULIN PROMOTER => d l - cit ab</p>	<p>construct; David R. Hurwitz, et al., 435/69.6, 320.1; 536/23.1, 23.5, 24.1, 24.2 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,648,243 [IMAGE AVAILABLE] L10: 4 of 8</p> <p>ABSTRACT: The present invention provides DNA constructs comprising a promoter DNA sequence and a DNA sequence coding for human serum albumin. In one embodiment the human serum albumin sequence comprises at least one, but not all, of the introns in the naturally occurring gene encoding for the HSA protein. In another embodiment the DNA constructs comprise a 5' regulatory sequence which directs the expression and secretion of HSA protein in the milk of a transgenic animal. Preferably, the promoter gene is a milk protein promoter sequence such as .beta.-lactoglobulin. The present invention also provides transgenic animals which secrete HSA in the milk of lactating females. The present invention also provides vectors comprising the constructs of the present invention.</p> <p>5. 5,476,995, Dec. 19, 1995, Peptide production; Anthony J. Clark, et al., 800/16; 435/69.1, 317.1, 320.1 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,476,995 [IMAGE AVAILABLE] L10: 5 of 8</p> <p>ABSTRACT: A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for polypeptide into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.</p> <p>6. 5,366,894, Nov. 22, 1994, Peptide production; Anthony J. Clark, et al., 435/320.1, 69.1, 325 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,366,894 [IMAGE AVAILABLE] L10: 6 of 8</p> <p>ABSTRACT: This invention relates to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's casein promoter which when incorporated into a mammal permits the female species of that</p>	<p>A method of producing a substance comprising a peptide, involves incorporating a DNA sequence coding for the peptide into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The substance may be an (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The substance will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.</p> <p>7. 5,322,775, Jun. 21, 1994, Peptide production; Anthony J. Clark, et al., 435/69.1, 69.6, 69.7, 317.1, 320.1; 530/412 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,322,775 [IMAGE AVAILABLE] L10: 7 of 8</p> <p>ABSTRACT: A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for a polypeptide chain of said compound into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.</p> <p>8. 4,873,316, Oct. 10, 1989, Isolation of exogenous recombinant proteins from the milk of transgenic mammals; Harry Meade, et al., 800/7; 435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8; 530/360, 361, 416, 417, 418, 832, 833; 536/23.1, 23.4, 23.5; 800/18 [IMAGE AVAILABLE]</p> <p>US PAT NO: 4,873,316 [IMAGE AVAILABLE] L10: 8 of 8</p> <p>ABSTRACT: This invention relates to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's casein promoter which when incorporated into a mammal permits the female species of that</p>
<p>15 LACTOGLOBULIN PROMOTER (LACTOGLOBULIN(W)/PROMOTER) L10 8 IMMUNOGLOB? AND LACTOGLOBULIN PROMOTER => d l - cit ab</p>	<p>1. 5,849,992, Dec. 15, 1998, Transgenic production of antibodies in milk; Harry Meade, et al., 800/14, 7, 15, 16, 17, 18 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,849,992 [IMAGE AVAILABLE] L10: 1 of 8</p> <p>ABSTRACT: A method for the production of monoclonal antibodies in mammal's milk through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.</p> <p>2. 5,827,690, Oct. 27, 1998, Transgenic production of antibodies in milk; Harry Meade, et al., 800/7; 530/867 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,827,690 [IMAGE AVAILABLE] L10: 2 of 8</p> <p>ABSTRACT: A method for the production of monoclonal antibodies in mammal's milk, through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.</p> <p>3. 5,750,172, May 12, 1998, Transgenic non human mammal milk; Harry Meade, et al., 426/580; 435/69.1, 69.4, 69.51, 69.52, 69.6, 183, 215; 800/7 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,750,172 [IMAGE AVAILABLE] L10: 3 of 8</p> <p>ABSTRACT: This invention relates to the production of recombinant proteins, such as coagulation factors VIII and IX, tissue plasminogen activator (TPA), urokinase, growth hormone, insulin, interferons, interleukins, peptide hormones and **immunoglobulins**, in mammals' milk. Particularly, this invention relates to an expression system which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.</p> <p>4. 5,648,243, Jul. 15, 1997, Human serum albumin expression</p>	<p>construct; David R. Hurwitz, et al., 435/69.6, 320.1; 536/23.1, 23.5, 24.1, 24.2 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,648,243 [IMAGE AVAILABLE] L10: 4 of 8</p> <p>ABSTRACT: The present invention provides DNA constructs comprising a promoter DNA sequence and a DNA sequence coding for human serum albumin. In one embodiment the human serum albumin sequence comprises at least one, but not all, of the introns in the naturally occurring gene encoding for the HSA protein. In another embodiment the DNA constructs comprise a 5' regulatory sequence which directs the expression and secretion of HSA protein in the milk of a transgenic animal. Preferably, the promoter gene is a milk protein promoter sequence such as .beta.-lactoglobulin. The present invention also provides transgenic animals which secrete HSA in the milk of lactating females. The present invention also provides vectors comprising the constructs of the present invention.</p> <p>5. 5,476,995, Dec. 19, 1995, Peptide production; Anthony J. Clark, et al., 800/16; 435/69.1, 317.1, 320.1 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,476,995 [IMAGE AVAILABLE] L10: 5 of 8</p> <p>ABSTRACT: A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for polypeptide into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.</p> <p>6. 5,366,894, Nov. 22, 1994, Peptide production; Anthony J. Clark, et al., 435/320.1, 69.1, 325 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,366,894 [IMAGE AVAILABLE] L10: 6 of 8</p> <p>ABSTRACT: This invention relates to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's casein promoter which when incorporated into a mammal permits the female species of that</p>

mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.

=> s 11 and lactalbumin

1127 LACTALBUMIN
L11 249 L1 AND LACTALBUMIN

=> s 11 and lactalbumin promoter#

1127 LACTALBUMIN
36024 PROMOTER#
7 LACTALBUMIN PROMOTER#
(LACTALBUMIN(W)PROMOTER#)
L12 2 L1 AND LACTALBUMIN PROMOTER#

=> d 1- cit ab

1. 5,849,992, Dec. 15, 1998, Transgenic production of antibodies in milk; Harry Meade, et al., 800/14, 7, 15, 16, 17, 18 [IMAGE AVAILABLE]

US PAT NO: 5,849,992 [IMAGE AVAILABLE] L12: 1 of 2

ABSTRACT:

A method for the production of monoclonal antibodies in mammal's milk through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.

2. 5,827,690, Oct. 27, 1998, Transgenic production of antibodies in milk; Harry Meade, et al., 800/7; 530/867 [IMAGE AVAILABLE]

US PAT NO: 5,827,690 [IMAGE AVAILABLE] L12: 2 of 2

ABSTRACT:

A method for the production of monoclonal antibodies in mammal's milk, through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.

=> d 1 2 clms

US PAT NO: 5,849,992 [IMAGE AVAILABLE] L12: 1 of 2

CLAIMS:

CLMS(1)

What is claimed is:

1. A transgenic non-human mammal all of whose germ cells and somatic cells contain a heterologous **immunoglobulin** protein-coding sequence operatively linked to a promoter sequence that directs the preferential expression of said protein-coding sequence in mammary gland epithelial cells, thereby providing a heterologous and assembled **immunoglobulin** in the milk of said mammal wherein said heterologous and assembled **immunoglobulin** is in a functional configuration and is produced at levels of at least about 1 mg/ml in the milk of said mammal.

CLMS(2)

2. The transgenic mammal of claim 1 wherein said **immunoglobulin** comprises a tetrameric antibody directed against a pathogen.

CLMS(3)

3. The transgenic mammal of claim 1 wherein said **immunoglobulin** comprises a tetrameric antibody directed against a biologically active peptide.

CLMS(4)

4. The transgenic mammal of claim 1 wherein said biologically active peptide is selected from the group consisting of erythropoietin, tissue plasminogen activator and gamma interferon.

CLMS(5)

5. The transgenic mammal of claim 1 wherein said **immunoglobulin** comprises a tetrameric antibody directed against an enzyme.

CLMS(6)

6. The transgenic mammal of claim 1 wherein said mammal is selected from the group consisting of mice, cows, sheep, goats, and pigs.

CLMS(7)

7. The transgenic mammal of claim 1 wherein said promoter is selected from the group consisting of the casein promoter, the beta lactoglobulin promoter, the whey acid protein promoter, and the **lactalbumin** promoter**.

CLMS(8)

8. The transgenic mammal of claim 1 wherein said immunoglobulin comprises heavy and light chains.

CLMS(9)

9. The transgenic mammal of claim 1 wherein said **immunoglobulin** is of human origin.

CLMS(10)

10. A transgenic non-human goat all of whose germ cells and somatic cells contain a heterologous **immunoglobulin** protein-coding sequence operatively linked to a promoter sequence that directs the preferential expression of said protein-coding sequence in mammary gland epithelial cells, thereby providing a heterologous and assembled **immunoglobulin** in the milk of said goat, wherein said heterologous and assembled **immunoglobulin** is in a functional configuration and is produced at levels of at least about 1 mg/ml in the milk of said goat.

US PAT NO: 5,827,690 [IMAGE AVAILABLE] L12: 2 of 2

CLAIMS:

CLMS(1)

What is claimed is:

1. A high level expression method for providing a heterologous and assembled **immunoglobulin** in the milk of a transgenic mammal comprising: obtaining milk from a transgenic mammal having introduced into its germline a heterologous **immunoglobulin** protein-coding sequence operatively linked to a promoter sequence that results in the preferential expression of said protein-coding sequence in mammary gland epithelial cells, thereby providing said heterologous and assembled **immunoglobulin** in the milk of said mammal, wherein said heterologous and assembled **immunoglobulin** is a functional configuration and is produced at level of at least about 1 mg/ml in the milk of said mammal.

CLMS(2)

2. The method of claim 1 wherein said mammal is selected from the group consisting of mice, sheep, and pigs.

CLMS(3)

3. The method of claim 1 wherein said promoter is selected from the group consisting of the beta lactoglobulin promoter, whey acid protein promoter, and the **lactalbumin** **promoter**.

CLMS(4)

4. The method of claim 1 wherein said **immunoglobulin** comprises heavy and light chains.

CLMS(5)

5. The method of claim 1 wherein said **immunoglobulin** is of human origin.

CLMS(6)

6. The method of claim 1 wherein said **immunoglobulin** is purified from the milk of said mammal.

CLMS(7)

7. The method of claim 1 wherein said promoter is the casein promoter.

CLMS(8)

8. A high level expression method for providing a heterologous and assembled **immunoglobulin**, in the milk of a transgenic goat comprising:
obtaining milk from a transgenic goat having introduced into its genome a heterologous **immunoglobulin** protein-coding sequence
operatively linked to a promoter sequence that results in the preferential expression of said protein-coding sequence in mammary gland epithelial cells, thereby providing said heterologous and assembled **immunoglobulin** in the milk of said goat, wherein said heterologous and assembled **immunoglobulin** is a functional configuration and is produced at levels of at least about 1 mg/ml in the milk of said goat.

CLMS(9)

9. The method of claim 8 wherein said promoter is selected from the group consisting of the beta lactoglobulin promoter, whey acid protein promoter, and the **lactalbumin** **promoter**.

CLMS(10)

10. The method of claim 8 wherein said **immunoglobulin** comprises heavy and light chains.

CLMS(11)

11. The method of claim 8 wherein said **immunoglobulin** is of human origin.

CLMS(12)

12. The method of claim 8 wherein said **immunoglobulin** is purified from the milk of said goat.

CLMS(13)

13. The method of claim 8 wherein said promoter is the casein promoter.

=> s 536/23.1/ccls

L13 2866 536/23.1/CCLS

=> s 536/23.5/ccls

L14 2126 536/23.5/CCLS

=> s 536/23.53/ccls

L15 192 536/23.53/CCLS

=> s 536/24.1/ccls

L16 1349 536/24.1/CCLS

=> 514/44/ccls

'514' IS NOT A RECOGNIZED COMMAND

=> s 514/44/ccls

L17 733 514/44/CCLS

=> s 435/320.1/ccls

L18 6152 435/320.1/CCLS

=> s 113-118

L19 8837 (L13 OR L14 OR L15 OR L16 OR L17 OR L18)

=> s 119 and 11

L20 2207 L19 AND L1

=> s 120 and (whey acidic protein or casein or lactoglobulin or lactalbumin)

4328 WHEY
98021 ACIDIC
73025 PROTEIN
45 WHEY ACIDIC PROTEIN
(WHEY(W)ACIDIC(W)PROTEIN)
17607 CASEIN
635 LACTOGLOBULIN
1127 LACTALBUMIN
L21 295 L20 AND (WHEY ACIDIC PROTEIN OR CASEIN
OR LACTOGLOBULIN OR
LAC
TALBUMIN)

=> s 120 and (whey acidic protein or casein or lactoglobulin or lactalbumin)(w)(promoter#)

4328 WHEY
98021 ACIDIC
73025 PROTEIN
45 WHEY ACIDIC PROTEIN
(WHEY(W)ACIDIC(W)PROTEIN)
17607 CASEIN
635 LACTOGLOBULIN
1127 LACTALBUMIN
36024 PROMOTER#
47 (WHEY ACIDIC PROTEIN OR CASEIN OR
LACTOGLOBULIN OR LACTALBU
MIN
(W)(PROMOTER#)
L22 12 L20 AND (WHEY ACIDIC PROTEIN OR CASEIN OR
LACTOGLOBULIN OR
LAC
TALBUMIN)(W)(PROMOTER#)

=> d 1- cit ab

1. 5,928,904, Jul. 27, 1999, DNA encoding recombinant IL4 antibodies useful in treatment of IL4 mediated disorders; Stephen D. Holmes, et al., 435/69.6, 70.21, 71.1, **320.1**, 326, 328, 335; 530/300, 350, 387.3, 388.23; **536/23.5**, **23.53** [IMAGE AVAILABLE]

US PAT NO: 5,928,904 [IMAGE AVAILABLE] L22: 1 of 12

ABSTRACT:

Chimeric and humanized IL4 MAbs derived from high affinity MAbs, pharmaceutical compositions containing same, and methods of treatment are provided.

2. 5,877,010, Mar. 2, 1999, Thymidine kinase mutants; Lawrence A. Loeb, et al., **435/320.1** 243, 325; 536/23.2, **23.5**, 23.72, **24.1** [IMAGE AVAILABLE]

- described. Therapeutic compositions and cells including the DNA are also described. Other aspects of the invention include methods of treating subjects having cystic fibrosis which include administering an effective amount of the DNA to subjects having cystic fibrosis such that functional cystic fibrosis transmembrane conductance regulator is produced by the subject at a level which is not detrimental to the subject. The present invention also pertains to a method of introducing the DNA into a cell such that the membrane-associated protein is produced at a level which is not detrimental to the cell and cells produced by this method. Still other aspects of the invention include a method of assaying DNA for the presence or absence of a hormone responsive element in a species in which the hormone responsive element is functional and a method of selectively breeding female transgenic mammals which produce a protein of interest.
5. 5,683,892, Nov. 4, 1997, DNA encoding recombinant IL-5 antagonists useful in treatment of IL-5 mediated disorders; Robert S. Ames, Jr., et al., 435/69.1, 69.3, 70.21, 252.3, **320.1** 328, **536/23.53** [IMAGE AVAILABLE]
- US PAT NO: 5,736,388 [IMAGE AVAILABLE] L22: 3 of 12
- ABSTRACT:
Lamboid bacteriophage capable of specifically interacting with and delivering nucleic acid molecules to eukaryotic cells are disclosed. Such bacteriophage-derived gene transfer systems target one or more specific receptors on eukaryotic cells, for instance by incorporating mutant tail fiber proteins or by incorporating known ligands for specific eukaryotic receptors into lambda phage. Also disclosed are methods for identifying and producing modified bacteriophage tail fiber polypeptides capable of specifically interacting with eukaryotic transmembrane proteins. Methods of treating diseases using such gene transfer systems are also disclosed.
4. 5,688,677, Nov. 18, 1997, Deoxyribonucleic acids containing inactivated hormone responsive elements; Karl M. Ebert, et al., **536/23.5** 24.1** [IMAGE AVAILABLE]
- US PAT NO: 5,688,677 [IMAGE AVAILABLE] L22: 4 of 12
- ABSTRACT:
A DNA comprising at least one inactivated hormone responsive element and a nucleic acid sequence encoding a membrane-associated protein is described. Therapeutic compositions and cells including the DNA are also described. Other aspects of the invention include methods of treating subjects having cystic fibrosis which include administering an effective amount of the DNA to subjects having cystic fibrosis such that functional cystic fibrosis transmembrane conductance regulator is produced by the subject at a level which is not detrimental to the subject. The present invention also pertains to a method of introducing the DNA into a cell such that the membrane-associated protein is produced at a level which is not detrimental to the cell and cells produced by this method. Still other aspects of the invention include a method of assaying DNA for the presence or absence of a hormone responsive element in a species in which the hormone responsive element is functional and a method of selectively breeding female transgenic mammals which produce a protein of interest.
5. 5,683,892, Nov. 4, 1997, DNA encoding recombinant IL-5 antagonists useful in treatment of IL-5 mediated disorders; Robert S. Ames, Jr., et al., 435/69.1, 69.3, 70.21, 252.3, **320.1** 328, **536/23.53** [IMAGE AVAILABLE]
- US PAT NO: 5,683,892 [IMAGE AVAILABLE] L22: 5 of 12
- ABSTRACT:
DNA encoding chimeric, humanized and other IL-5 mAbs, derived from high affinity neutralizing mAbs, pharmaceutical compositions containing same, methods of treatment and diagnostics are provided.
6. 5,681,746, Oct. 28, 1997, Retroviral delivery of full length factor VIII; Mordechai Bodner, et al., 433/350, **320.1** 366, 371; **536/23.5** [IMAGE AVAILABLE]
- US PAT NO: 5,681,746 [IMAGE AVAILABLE] L22: 6 of 12
- ABSTRACT:
Retroviral vectors for directing expression of full length factor VIII in transduced host cells, plasmids encoding the same, and host cells transformed, transfected, or transduced therewith are disclosed. Also disclosed are retroviral particles comprising such retroviral vectors, as are methods for making such particles in suitable packaging cells. Retroviral particles so produced may be amphotropic, ecotropic, polytropic, or xenotropic; alternatively, they may comprise chimeric or hybrid envelope proteins to alter host range. Also described are retroviral particles comprising retroviral vectors for directing full length factor VIII expression which are complement resistant.
7. 5,648,243, Jul. 15, 1997, Human serum albumin expression construct; David R. Hurwitz, et al., 435/69.6, **320.1** 536/23.1** 23.5** 24.1** 24.2 [IMAGE AVAILABLE]
- US PAT NO: 5,648,243 [IMAGE AVAILABLE] L22: 7 of 12
- ABSTRACT:
The present invention provides DNA constructs comprising a promoter DNA sequence and a DNA sequence coding for human serum albumin. In one embodiment the human serum albumin sequence comprises at least one, but not all, of the introns in the naturally occurring gene encoding for the HSA protein. In another embodiment the DNA constructs comprise a regulatory sequence which directs the expression and secretion of HSA protein in the milk of a transgenic animal. Preferably, the promoter gene is a milk protein promoter sequence such as .beta.-lactoglobulin. The present invention also provides transgenic animals which secrete HSA in the milk of lactating females. The present invention also provides vectors comprising the constructs of the present invention.
8. 5,476,995, Dec. 19, 1995, Peptide production; Anthony J. Clark, et al., 800/16, 435/69.1, 317.1, **320.1** [IMAGE AVAILABLE]
- US PAT NO: 5,476,995 [IMAGE AVAILABLE] L22: 8 of 12
- ABSTRACT:
A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for polypeptide into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.
9. 5,366,894, Nov. 22, 1994, Peptide production; Anthony J. Clark, et
- Pharmaceutical compositions comprising retroviral particles of the invention are also disclosed, as are methods of treating mammals, particularly humans, afflicted with hemophilia.
7. 5,648,243, Jul. 15, 1997, Human serum albumin expression construct; David R. Hurwitz, et al., 435/69.6, **320.1** 536/23.1** 23.5** 24.1** 24.2 [IMAGE AVAILABLE]
- US PAT NO: 5,648,243 [IMAGE AVAILABLE] L22: 7 of 12
- ABSTRACT:
The present invention provides DNA constructs comprising a promoter DNA sequence and a DNA sequence coding for human serum albumin. In one embodiment the human serum albumin sequence comprises at least one, but not all, of the introns in the naturally occurring gene encoding for the HSA protein. In another embodiment the DNA constructs comprise a regulatory sequence which directs the expression and secretion of HSA protein in the milk of a transgenic animal. Preferably, the promoter gene is a milk protein promoter sequence such as .beta.-lactoglobulin. The present invention also provides transgenic animals which secrete HSA in the milk of lactating females. The present invention also provides vectors comprising the constructs of the present invention.
8. 5,476,995, Dec. 19, 1995, Peptide production; Anthony J. Clark, et al., 800/16, 435/69.1, 317.1, **320.1** [IMAGE AVAILABLE]
- US PAT NO: 5,476,995 [IMAGE AVAILABLE] L22: 8 of 12
- ABSTRACT:
A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for polypeptide into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.
9. 5,366,894, Nov. 22, 1994, Peptide production; Anthony J. Clark, et

al., **435/320.1**, 69.1, 325 [IMAGE AVAILABLE]
 US PAT NO: 5,366,894 [IMAGE AVAILABLE] L22: 9 of 12

ABSTRACT:
 A method of producing a substance comprising a peptide, involves incorporating a DNA sequence coding for the peptide into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The substance may be an (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The substance will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.

10, 5,322,775, Jun. 21, 1994, Peptide production; Anthony J. Clark, et al., 435/69.1, 69.6, 69.7, 317.1, **320.1**, 530/412 [IMAGE AVAILABLE]
 US PAT NO: 5,322,775 [IMAGE AVAILABLE] L22: 10 of 12

ABSTRACT:
 A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for a polypeptide chain of said compound into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.

11, 5,268,275, Dec. 7, 1993, Vitamin K-dependent carboxylase; Daniel W. Stafford, et al., 435/69.1, 69.6, 232, 252.3, **320.1**, 352, 354, 358, 366, 536/23.2 [IMAGE AVAILABLE]
 US PAT NO: 5,268,275 [IMAGE AVAILABLE] L22: 11 of 12

ABSTRACT:
 Isolated DNA encoding a vitamin K dependent carboxylase is disclosed. The carboxylase is selected from the group consisting of: (a) isolated DNA which encodes bovine or human vitamin K dependent carboxylase; (b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a vitamin K dependent carboxylase; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) above in nucleotide sequence due to the degeneracy of the genetic code, and which encodes a vitamin K dependent carboxylase. Also disclosed are vectors and host cells containing the aforesaid DNA, methods of using the same, and purified protein coded for by the aforesaid DNA.

12, 4,873,316, Oct. 10, 1989, Isolation of exogenous recombinant proteins from the milk of transgenic mammals; Harry Meade, et al., 435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8, 530/360, 361, 416, 417, 418, 832, **536/23.1**, 23.4, **23.5**, 800/18 [IMAGE AVAILABLE]
 US PAT NO: 4,873,316 [IMAGE AVAILABLE] L22: 12 of 12

ABSTRACT:
 This invention relates to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's **casein** **promoter** which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.

=> d 12 kwic

US PAT NO: 4,873,316 [IMAGE AVAILABLE] L22: 12 of 12
 US-CL-CURRENT: 800/7; 435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8; 530/360, 361, 416, 417, 418, 832, 833; **536/23.1**, 23.4, **23.5**, 800/18

ABSTRACT:
 This . . . to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's **casein** **promoter** which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein. . . .

DETDSC:

al., **435/320.1**, 69.1, 325 [IMAGE AVAILABLE]
 US PAT NO: 5,366,894 [IMAGE AVAILABLE] L22: 9 of 12

ABSTRACT:
 A method of producing a substance comprising a peptide, involves incorporating a DNA sequence coding for the peptide into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The substance may be an (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The substance will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.

10, 5,322,775, Jun. 21, 1994, Peptide production; Anthony J. Clark, et al., 435/69.1, 69.6, 69.7, 317.1, **320.1**, 530/412 [IMAGE AVAILABLE]
 US PAT NO: 5,322,775 [IMAGE AVAILABLE] L22: 10 of 12

ABSTRACT:
 A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for a polypeptide chain of said compound into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.

11, 5,268,275, Dec. 7, 1993, Vitamin K-dependent carboxylase; Daniel W. Stafford, et al., 435/69.1, 69.6, 232, 252.3, **320.1**, 352, 354, 358, 366, 536/23.2 [IMAGE AVAILABLE]
 US PAT NO: 5,268,275 [IMAGE AVAILABLE] L22: 11 of 12

ABSTRACT:
 Isolated DNA encoding a vitamin K dependent carboxylase is disclosed. The carboxylase is selected from the group consisting of: (a) isolated DNA which encodes bovine or human vitamin K dependent carboxylase; (b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a vitamin K dependent carboxylase; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) above in nucleotide sequence due to the degeneracy of the genetic code, and which encodes a vitamin K dependent carboxylase. Also disclosed are vectors and host cells containing the aforesaid DNA, methods of using the same, and purified protein coded for by the aforesaid DNA.

12, 4,873,316, Oct. 10, 1989, Isolation of exogenous recombinant proteins from the milk of transgenic mammals; Harry Meade, et al., 435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8, 530/360, 361, 416, 417, 418, 832, **536/23.1**, 23.4, **23.5**, 800/18 [IMAGE AVAILABLE]
 US PAT NO: 4,873,316 [IMAGE AVAILABLE] L22: 12 of 12

ABSTRACT:
 This invention relates to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's **casein** **promoter** which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.

=> d 12 kwic

US PAT NO: 4,873,316 [IMAGE AVAILABLE] L22: 12 of 12
 US-CL-CURRENT: 800/7; 435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8; 530/360, 361, 416, 417, 418, 832, 833; **536/23.1**, 23.4, **23.5**, 800/18

ABSTRACT:
 This . . . to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's **casein** **promoter** which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein. . . .

DETDSC:

al., **435/320.1**, 69.1, 325 [IMAGE AVAILABLE]
 US PAT NO: 5,366,894 [IMAGE AVAILABLE] L22: 9 of 12

ABSTRACT:
 A method of producing a substance comprising a peptide, involves incorporating a DNA sequence coding for the peptide into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The substance may be an (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The substance will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.

10, 5,322,775, Jun. 21, 1994, Peptide production; Anthony J. Clark, et al., 435/69.1, 69.6, 69.7, 317.1, **320.1**, 530/412 [IMAGE AVAILABLE]
 US PAT NO: 5,322,775 [IMAGE AVAILABLE] L22: 10 of 12

ABSTRACT:
 A method of producing a proteinaceous compound, involves incorporating a DNA sequence coding for a polypeptide chain of said compound into a gene of a mammal (such as a sheep) coding for a milk whey protein in such a way that the DNA sequence is expressed in the mammary gland of the adult female mammal. The proteinaceous compound may be a (optionally modified) protein such as a blood coagulation factor. The DNA sequence is preferably inserted into the first exon of a gene coding for a whey protein such as beta-lactoglobulin. The proteinaceous compound will generally be recovered from milk of the female mammal, but may (for example if it is an enzyme) be used in situ.

11, 5,268,275, Dec. 7, 1993, Vitamin K-dependent carboxylase; Daniel W. Stafford, et al., 435/69.1, 69.6, 232, 252.3, **320.1**, 352, 354, 358, 366, 536/23.2 [IMAGE AVAILABLE]
 US PAT NO: 5,268,275 [IMAGE AVAILABLE] L22: 11 of 12

ABSTRACT:
 Isolated DNA encoding a vitamin K dependent carboxylase is disclosed. The carboxylase is selected from the group consisting of: (a) isolated DNA which encodes bovine or human vitamin K dependent carboxylase; (b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a vitamin K dependent carboxylase; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) above in nucleotide sequence due to the degeneracy of the genetic code, and which encodes a vitamin K dependent carboxylase. Also disclosed are vectors and host cells containing the aforesaid DNA, methods of using the same, and purified protein coded for by the aforesaid DNA.

12, 4,873,316, Oct. 10, 1989, Isolation of exogenous recombinant proteins from the milk of transgenic mammals; Harry Meade, et al., 435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8, 530/360, 361, 416, 417, 418, 832, **536/23.1**, 23.4, **23.5**, 800/18 [IMAGE AVAILABLE]
 US PAT NO: 4,873,316 [IMAGE AVAILABLE] L22: 12 of 12

ABSTRACT:
 This invention relates to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's **casein** **promoter** which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.

=> d 12 kwic

US PAT NO: 4,873,316 [IMAGE AVAILABLE] L22: 12 of 12
 US-CL-CURRENT: 800/7; 435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8; 530/360, 361, 416, 417, 418, 832, 833; **536/23.1**, 23.4, **23.5**, 800/18

ABSTRACT:
 This . . . to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's **casein** **promoter** which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein. . . .

DETDSC:

APPL-NO: 07/065,994
DATE FILED: Jun. 23, 1987
INT-TITLE: [4] C07K 3/02; C07K 3/18; C12N 15/00
US-CL-ISSUED: 530/412, 360, 361, 833, 832, 416, 417, 418; 435/68, 172.1,
172.3, 240.2; 935/53, 55, 70, 111; 800/1; 536/27, 28, 29
US-CL-CURRENT: 800/7; 435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8;
530/360,
361, 416, 417, 418, 832, 833; **536/23.1** 23.4,
23.5 800/18
SEARCH-FLD: 435/68, 172.1, 172.3, 226, 240.2; 530/832, 833,
412, 360,
361, 303; 800/1; 935/53, 55, 70; 536/27, 28, 29
REF-CITED:
U.S. PATENT DOCUMENTS
4,018,752 4/1977 Buhler et al. 530/382
4,229,342 10/1980 Mirabel 530/382
4,376,072 3/1983 Connolly 530/382
4,396,601 8/1983 Salser et al. 435/172.3
4,462,932 7/1984 Loneragan 530/382
4,644,056 2/1987 Kothe et al. 530/382
4,736,866 4/1988 Leder et al.
FOREIGN PATENT DOCUMENTS
0117059 8/1984 European Patent Office 435/172.3
0247494 12/1987 European Patent Office
0264166 4/1988 European Patent Office
WO88/00239 1/1988 World Intellectual Property Organization
WO88/01648 3/1988 World Intellectual Property Organization
OTHER PUBLICATIONS
Gordon et al., BioTechnology, 5, 1183-7, (Nov. 1987).
Lovell-Badge, Nature, 315, 628-629, (1985).
Hammer et al., Nature, 415, 680-683, (Jun. 20, 1985).
Garcia et al., Mol. Cell. Biol., 6(6), 1974-82, (1986).
Palmiter et al., J. Cell. Biochem., 8B, p. 25, Ab. #0890, (1984).
Fisher et al., J. B. C., 260 (20), 11223-11230, 1985.
Andres et al., Chem. Abs. 106(17):133024q, 1987 (for PNAS USA, 84(5)).
1299-303, May 1987).
Ross et al., P.N.A.S. (USA), 82, 5880-84, 1985.
Brinster et al., Cell, 27, 223-31, Nov. 1981.
ART-UNIT: 186
PRIM-EXMR: Margaret Moskowitz
ASST-EXMR: Jeff P. Kushan
LEGAL-REP: James F. Haley, Jr., Teresa L. Solomon

ABSTRACT:
This invention relates to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's **cascin** **promoter** which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along

with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.
3 Claims, 1 Drawing Figures
=> file jpoab
FILE 'JPOABS' ENTERED AT 17:42:26 ON 04 AUG 1999

* J A P A N E S E P A T E N T A B S T R A C T S *
*
* DATA IS LOADED THROUGH DECEMBER 24, 1996, FOR THE JAPANESE *
* PATENT OFFICE ABSTRACT (JPOABS) FILE. NEW RECORDS ARE NOT *
* BEING ADDED. PLEASE USE THE GPI-JPO FILE (JPO) WHICH IS *
* CURRENT THROUGH MARCH 31, 1999 (SEE BELOW). *

* GLOBAL PATENT INFORMATION-JAPANESE PATENT OFFICE FILE *
* (GPI-JPO FILE) *
* THE FILE IS CURRENT THROUGH MARCH 31, 1999. *

IRD CNOABS

=> s 11

L23 505 IMMUNOGLOB?

=> s 12

505 IMMUNOGLOB?
580 WHEY
14269 ACIDIC
10862 PROTEIN
0 WHEY ACIDIC PROTEIN
(WHEY(W)ACIDIC(W)PROTEIN)
0 L1 AND WHEY ACIDIC PROTEIN

=> s 13

505 IMMUNOGLOB?
580 WHEY
14269 ACIDIC
10862 PROTEIN
0 WHEY ACIDIC PROTEIN
(WHEY(W)ACIDIC(W)PROTEIN)
2534 PROMOTER#
0 L2 AND PROMOTER#

L25

=> s 14

505 IMMUNOGLOB?
580 WHEY
14269 ACIDIC
10862 PROTEIN
0 WHEY ACIDIC PROTEIN
(WHEY(W)ACIDIC(W)PROTEIN)
2534 PROMOTER#
7370 CONSTRUCT#
15691 VECTOR#
2014 PLASMIID#
L26 0 L3 AND (CONSTRUCT# OR VECTOR# OR PLASMIID#)

=> s 15

505 IMMUNOGLOB?
1448 CASEIN
L27 4 IMMUNOGLOB? AND CASEIN

=> d 1 - cit ab

1. 07-285885, Oct. 31, 1995, PRODUCTION OF
IMMUNOGLOBULIN.; HIFUMI
OISHI, et al., A61K 39/395; B01D 61/14; C07K 16/06
07-285885 L27: 1 of 4

ABSTRACT:

PURPOSE: To provide **immunoglobulin** and a method for producing an
immunoglobulin monomer from the resultant
immunoglobulin
(antigen).

CONSTITUTION: **Casein** is subjected to an isoelectric point (pH4.5 to 4.6) precipitation treatment or treated with an enzyme such as rennin to obtain whey. Ethylenediaminetetraacetic acid or glycine is added to the whey so that the concentration of the respective components may be 0.5 to 10mM or 50 to 500mM. pH of the whey is adjusted to 6.0 to 6.5 by using a sodium citrate solution and a cation-exchange resin is brought into contact therewith. The treated material is further treated by using an ultrafiltration module having 50000 or 100000 dalton separation limitation, thus obtaining the objective **immunoglobulin**.

COPYRIGHT: (C)1995,JPO

2. 63-135336, Jun. 7, 1988, DRUG FOR INTESTINAL DISORDER;
TOSHIRO HORI,

et al., A61K 39/395; A61K 35/20

63-135336 L27: 2 of 4

ABSTRACT:

PURPOSE: To obtain an agent for controlling intestinal disorder, by compounding **immunoglobulin** of cattle whey as an active component.

CONSTITUTION: The objective intestinal disorder controlling agent can be

produced by using **immunoglobulin** of cattle whey (whole **immunoglobulin** existing in cattle whey) in an amount of .gforeq.50%

of the whole solid component and properly mixing the **immunoglobulin**

with a vehicle. It is necessary to administer the **immunoglobulin** of cattle whey at a dose of .gforeq.10mg/kg and the dose is preferably .lforeq.1g for person taking the easiness of administration into consideration. The **immunoglobulin** of cattle whey used in the agent

can be produced by treating normal milk of healthy cow with an enzyme

and/or acid, removing precipitated **casein**, subjecting the obtained whey part to ion-exchange treatment, etc., to remove low-molecular weight

soluble salts, lactose, etc., and separating and concentrating the **immunoglobulin** fraction. The **immunoglobulin** may be used in the

form of powder, liquid, etc., however, use of pulverized globulin is preferable for the convenience of storage and handling.

3. 63-135323, Jun. 7, 1988, COSMETIC; TOSHIRO HORI, et al., A61K 7/40

63-135323 L27: 3 of 4

ABSTRACT:

PURPOSE: To obtain a cosmetic suitably useful for treating pimples, by blending a cosmetic base with **immunoglobulin** of bovine milk serum as

an active ingredient.

CONSTITUTION: Bovine milk serum part obtained by treating ordinary milk

casein form the milk is subjected to ion exchange, gel filtration, affinity chromatography or ultrafiltration with 100,000mol.wt. of elimination limit to remove low-molecular soluble salt, lactose, protein

having tens thousands molecular weight, etc. The prepared

immunoglobulin fraction is separated and concentrated to give **immunoglobulin** of bovine milk serum. A cosmetic is blended with the

immunoglobulin as an active ingredient. The amount of the **immunoglobulin** in the cosmetic is preferably .gforeq.5wt% calculated

as a solid substance.

4. 60-75433, Apr. 27, 1985, METHOD FOR CONCENTRATING **IMMUNOGLOBULIN**

IN COW'S COLOSTRUM; YOSHIYUKI MINAMI, et al., A61K 39/395

60-75433 L27: 4 of 4

ABSTRACT:

PURPOSE: To concentrate efficiently the titled globulin, by defatting cow's colostrum, adding an acid to the defatted cow's colostrum to adjust

the pH to a specific value, adding pepsin or rennin thereto to coagulate and separate **casein**, and ultrafiltering the resultant whey.

CONSTITUTION: A colostrum collected within about one week after the

delivery of a cow is defatted with a cream separator, and an acid, e.g. acetic or propionic acid, is added to the defatted colostrum to adjust the pH to 5.0 approx.6.0. Pepsin or rennin is then added thereto

coagulate **casein**. In the process, the defatted colostrum is preferably heated to 50 approx.55 degrees C. Crud is flocculated by heating, and whey is collected. The recovered whey is then centrifuged at

a low speed, and the remaining fine particles are removed. The whey is then preferably quenched to 10 degrees C. or below. The resultant whey is

concentrated in an ultrafiltration apparatus, e.g. plate form, etc. The resultant whey containing concentrated **immunoglobulin** is freeze-dried

or spray-dried to give a raw material used to prepare a feed for young animals.

=> s 19

505 IMMUNOGLOB? 30 LACTOGLOBULIN

L28 0 IMMUNOGLOB? AND LACTOGLOBULIN

=> s 111

505 IMMUNOGLOB? 39 LACTALBUMIN

L29 0 LI AND LACTALBUMIN

=> file epoab

FILE 'EPOABS' ENTERED AT 17:44:16 ON 04 AUG 1999

* EUROPEAN PATENT ABSTRACTS *

FILE 'EPOABS' ENTERED AT 17:44:16 ON 04 AUG 1999

=> s 123

L30 894 IMMUNOGLOB?

=> s 124

894 IMMUNOGLOB?

852 WHEY
7450 ACIDIC
12508 PROTEIN

3 WHEY ACIDIC PROTEIN
(WHEY(W)ACIDIC(W)PROTEIN)

L31 0 LI AND WHEY ACIDIC PROTEIN

=> s 126

894 IMMUNOGLOB?

852 WHEY
7450 ACIDIC
12508 PROTEIN

3 WHEY ACIDIC PROTEIN
(WHEY(W)ACIDIC(W)PROTEIN)

L32 0 L3 AND (CONSTRUCT# OR VECTOR# OR PLASMID#)

=> s 127

894 IMMUNOGLOB?

510 CASEIN
5 IMMUNOGLOB? AND CASEIN

L33 5 IMMUNOGLOB? AND CASEIN

=> d l - cit ab

894 IMMUNOGLOB?

510 CASEIN
5 IMMUNOGLOB? AND CASEIN

L33 5 IMMUNOGLOB? AND CASEIN

=> d l - cit ab

894 IMMUNOGLOB?

510 CASEIN
5 IMMUNOGLOB? AND CASEIN

L33 5 IMMUNOGLOB? AND CASEIN

=> d l - cit ab

894 IMMUNOGLOB?

510 CASEIN
5 IMMUNOGLOB? AND CASEIN

L33 5 IMMUNOGLOB? AND CASEIN

=> d l - cit ab

894 IMMUNOGLOB?

510 CASEIN
5 IMMUNOGLOB? AND CASEIN

L33 5 IMMUNOGLOB? AND CASEIN

=> d l - cit ab

<CHG DATE=19960327 STATUS=O>The present invention provides an improved biotin-avidin conjugate in which a biotinylated antibody conjugate or an avidin-enzyme conjugate is present in a suitable diluent for immunohistochemical staining. The diluent additionally comprises ****casein**** in an amount sufficient to prevent charge interactions of the conjugate with a tissue section. In a preferred embodiment, the ****immunoglobulin**** is from the same species as the biotinylated antibody conjugate. The formulation effectively reduces overall unwanted binding, irrespective of the source of the binding.

2. WO 09508562A1, Mar. 30, 1995, METHOD OF OBTAINING ****IMMUNOGLOBULINS**** FROM COLOSTRUM AND THEIR USE IN PHARMACEUTICAL COMPOSITION; CONOR JOHN GRAHAM, C07K 1/14; C07K 1/30; C07K 16/04; A61K 9/20; A61K 39/395

WO 09508562A1 L33: 2 of 5

ABSTRACT:

A method of obtaining a high purity ****immunoglobulin**** preparation from an antibody rich colostrum which includes: (i) removing milk fat from the colostrum to obtain a low-fat colostrum; (ii) pasteurising the low-fat colostrum; (iii) coagulating the pasteurised, low-fat colostrum and removing milk curd containing ****casein****; (iv) centrifuging remaining liquid to remove precipitates; (v) removing lactose, minerals and water to obtain an antibody containing fraction; (vi) dissolving the antibody containing fraction in THRESH buffer and idolizing against the same buffer; and (vii) concentrating the antibody containing solution to obtain a 10 % by weight antibody solution. A pharmaceutical composition including a core element which includes an active antibody component derived by the above method, wherein the core element is in the form of a tablet, and wherein the compression forces used to prepare the tablet are such that they do not injure or denature the active antibodies.

3. WO 08910139A1, Nov. 2, 1989, PREPARATION WITH ANTIBODY ACTIVITY AND BROAD SPECTRUM; HERBERT DICHTELMEUELLER, et al., A61K 39/395; /C07K 15/06; C07K 3/02

WO 08910139A1 L33: 3 of 5

ABSTRACT:

<CHG DATE=19940730 STATUS=O>A preparation with antibody activity is prepared from colostrum of non-immunized mammals extracted during the first 30 hours, preferably however during the first 10 hours, following parturition. The colostrum is diluted with water, pasteurized, and after removal of the ****casein**** and fat, concentrated and stabilized. The preparation has a high ****immunoglobulin**** content (>80 %) and low anticomplementary activity. It can be administered orally in humans and intravenously in veterinary medicine. It can be used successfully, alone or in combination with other pharmaceutical substances, to treat bacteria- or toxin-induced diseases, in particular severe diarrhea in AIDS patients and other immunological disorders, travellers' diarrhea and toxin-induced infantile diarrhea, gastric and intestinal ulcers, as well as chronic and acute Yersinia infections, and to combat protozoa.

4. GB 02188526A, Oct. 7, 1987, Whey protein, JOHN BURTON, et al., A23C 9/146; A23J 1/20

GB 02188526A L33: 4 of 5

ABSTRACT:

****casein****, comprising a polypeptide or mixture of polypeptides substantially free of native alpha-, beta- and kappa-****casein****, serum albumin, alpha-lactalbumin, beta-lactoglobulin and the ****immunoglobulins****, and ****immunoglobulin**** remains in solution at pH 4.6 to pH 5.3 at 20 DEG C, and ****immunoglobulin**** is anionic at pH 4.6 to pH 5.3; and ****immunoglobulin**** forms a gel when an aqueous solution containing at least 12% w/v of the proteinaceous material at 20 DEG C and pH 4.5 or below is allowed to stand for 18 to 24 h. ****immunoglobulin**** milk whey at pH 4 to 6 may be contacted with an anion exchange resin, the resin may be eluted with HCl or NaCl and the product may be concentrated by ultrafiltration or thermal evaporation and/or spray dried or freeze-dried.

5. GB 02179947A, Mar. 18, 1987, Process for the extraction of proteins

from milk; PIERRE FREDERIC EMMANUE MONSAN, et al., C07K 3/22; C07K 3/02; C07K 3/28

GB 02179947A L33: 5 of 5

ABSTRACT:

****immunoglobulin****, comprising a polypeptide or mixture of polypeptides substantially free of native alpha-, beta- and kappa-****casein****, serum albumin, alpha-lactalbumin, beta-lactoglobulin and the ****immunoglobulins****, and ****immunoglobulin**** remains in solution at pH 4.6 to pH 5.3 at 20 DEG C, and ****immunoglobulin**** forms a gel when an aqueous solution containing at least 12% w/v of the proteinaceous material at 20 DEG C and pH 4.5 or below is allowed to stand for 18 to 24 h. ****immunoglobulin**** milk whey at pH 4 to 6 may be contacted with an anion exchange resin, the resin may be eluted with HCl or NaCl and the product may be concentrated by ultrafiltration or thermal evaporation and/or spray dried or freeze-dried.

6. US 04849241A, Jul. 18, 1989, Novel process for lowering the concentration of beta-****lactoglobulin**** in cheese whey, SHALAN A AL-MASHIKI, et al., A23C 21/10

US 04849241A L34: 1 of 3

ABSTRACT:

<CHG DATE=19940730 STATUS=O>A process for lowering the concentration of beta-****lactoglobulin**** in cheese whey while retaining the ****immunoglobulins**** in said cheese whey which comprises treating said cheese whey with a polyphosphate, such as sodium hexametaphosphate, within a pH range of from about 3.8 to about 4.7.

7. US 04112123A, Sep. 5, 1978, Nutritionally balanced single food composition and method of production; WILLARD LEWIS ROBERTS, A23C 21/00

US 04112123A L34: 2 of 3

***** STN Columbus *****

FILE 'HOME' ENTERED AT 10:20:31 ON 10 AUG 1999

=> file medline

COST IN U.S. DOLLARS	ENTRY	SINCE FILE	TOTAL
FULL ESTIMATED COST	SESSION	0.15	0.15

FILE 'MEDLINE' ENTERED AT 10:20:36 ON 10 AUG 1999

FILE LAST UPDATED: 6 AUG 1999 (19990806/UP). FILE
COVERS 1960 TO DATE.

MEDLINE has been reloaded to reflect the annual MeSH changes
made by
the National Library of Medicine for 1999. Enter HELP RLOAD for
details.

OLDMEDLINE, data from 1960 through 1965 from the Cumulated
Index
Medicus (CIM), has been added to MEDLINE. See HELP
CONTENT for details.

Left, right, and simultaneous left and right truncation are available in
the
Basic Index. See HELP SFIELDS for details.

THIS FILE CONTAINS CAS REGISTRY NUMBERS FOR EASY
AND ACCURATE
SUBSTANCE IDENTIFICATION.

=> s immunoglob? or antibod?/ab,bi

'AB' IS NOT A VALID FIELD CODE
108967 IMMUNOGLOB?

532337 ANTIBOD?/BI

L1 583923 IMMUNOGLOB? OR ANTIBOD?/AB,BI

=> s l1(5a)(construct# or plasmid# or vector#)/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 CONSTRUCT#/AB

26368 CONSTRUCT#/BI

0 PLASMID#/AB

71642 PLASMID#/BI

0 VECTOR#/AB

59945 VECTOR#/BI

L2 843 L1(5A)(CONSTRUCT# OR PLASMID# OR
VECTOR#/AB,BI

=> s l2(5)(milk)/ab,bi

MISSING OPERATOR 'L2(5'

The search profile that was entered contains terms or
nested terms that are not separated by a logical operator.

=> s l2(5a)(milk)/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 (MILK)/AB

48429 (MILK)/BI

L3 2 L2(5A)(MILK)/AB,BI

=> d l- bib ab

YOU HAVE REQUESTED DATA FROM 2 ANSWERS -
CONTINUE? Y(N);y

L3 ANSWER 1 OF 2 MEDLINE
AN 93078105 MEDLINE

DN 93078105

TI Concentration of ***milk*** secretory

immunoglobulin A

against Shigella virulence ***plasmid*** -associated antigens as
a

predictor of symptom status in Shigella-infected breast-fed infants.
AU Hayati K C; Guerrero M L; Morrow A L; Gomez H F; Winsor D
K; Ruiz-Palacios

CS Department of Pediatrics, University of Texas Medical School,
Houston

77030.

NC 5-PO1-HD-13021 (NICHHD)

SO JOURNAL OF PEDIATRICS, (1992 Dec) 121 (6) 852-6.

Journal code: J.LZ. ISSN: 0022-3476.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Abridged Index Medicus Journals; Priority Journals; Cancer
Journals

EM 199303

AB We conducted a prospective, community-based study of healthy
breast-fed

Mexican infants to determine the protective effects of anti-Shigella
secretory IgA antibodies in milk. Milk samples were collected
monthly, and

stool culture specimens were obtained weekly and at the time of
episodes

of diarrhea. Nineteen breast-fed infants were found to have Shigella

flexneri, Shigella boydii, or Shigella sonnei in stool samples. Ages

of

the 10 infants with symptomatic infection and the nine with

asymptomatic

infection did not differ significantly. Milk samples collected up to

12

weeks before infection were evaluated by enzyme-linked

immunosorbent assay

for secretory IgA antibodies against lipopolysaccharides of S.

flexneri.

S. boydii serotype 2, S. sonnei, and virulence plasmid-associated
antigens. The geometric mean titers of anti-Shigella

antibodies

to virulence ***plasmid*** -associated antigens in ***milk***

received before infection were eightfold higher in infants who

remained

well than in those in whom diarrhea developed. The significance of

milk

secretory IgA directed against lipopolysaccharide was less clear.

We

conclude that human milk protects infants against symptomatic

shigella

infection when it contains high concentrations of secretory IgA

against

virulence plasmid-associated antigens.

L3 ANSWER 2 OF 2 MEDLINE

AN 91093893 MEDLINE

DN 91093893

TI Human ***milk*** secretory ***immunoglobulin*** A to

Shigella

virulence ***plasmid*** -coded antigens.

AU Cleary T G; West M S; Ruiz-Palacios G; Winsor D K; Calva J J;
Guerrero M

L; Van R

CS Department of Pediatrics and Microbiology, University of Texas
Medical

School at Houston 77030.

NC 5-PO1-HD-13021 (NICHHD)

SO JOURNAL OF PEDIATRICS, (1991 Jan) 118 (1) 34-8.

Journal code: J.LZ. ISSN: 0022-3476.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Abridged Index Medicus Journals; Priority Journals; Cancer
Journals

EM 199104

AB Although antibodies to the lipopolysaccharide antigens of

Shigella have

been demonstrated in human milk, such antibodies do not explain

the

putative protective effect of breast-feeding against symptomatic

Shigella

infection. Shigella species do not share related lipopolysaccharides,

but

they do possess closely related virulence plasmids that code for the

proteins essential for cell invasion. We therefore sought to

determine the

frequency, amount, and duration of excretion of human

milk

antibodies to these shared virulence ***plasmid***

-associated

antigens in populations of different rates of Shigella infection

frequency

(Mexico City, high; Houston, low). Such antibodies were present in

the milk of virtually all the Mexican women but also were present in a large proportion of milk samples from the women living in Houston. The amounts of these antibodies were highest in colostrum but after 2 weeks of lactation fell to stable levels. The frequency and persistence of these antibodies in the milk of the women from Houston suggest that the memory and drive for secretion of these antibodies is extremely long lived.

=> s l1(5a)(mammary)

L4 36573 MAMMARY
357 L1(5A)(MAMMARY)

=> s l4(10a)(recombinant)

L5 143021 RECOMBINANT
0 L4(10A)(RECOMBINANT)

=> s l1(5a)(mammary)/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 (MAMMARY)/AB
36573 (MAMMARY)/BI
L6 357 L1(5A)(MAMMARY)/AB,BI

=> s l6(5a)(recombinant)/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 (RECOMBINANT)/AB
143021 (RECOMBINANT)/BI
L7 0 L6(5A)(RECOMBINANT)/AB,BI

=> s l6(5a)(specific)/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 (SPECIFIC)/AB
601385 (SPECIFIC)/BI
L8 33 L6(5A)(SPECIFIC)/AB,BI

=> s l8 and promoter#/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 PROMOTER#/AB
66697 PROMOTER#/BI
L9 0 L8 AND PROMOTER#/AB,BI

=> s l2(3a)(recombinant)/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 (RECOMBINANT)/AB
174143 (RECOMBINANT)/BI

L10 57 L2(5A)(RECOMBINANT)/AB,BI

=> s l10(5a)(milk or mammary)/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 MILK/AB
48429 MILK/BI
0 MAMMARY/AB
36573 MAMMARY/BI
L11 0 L10(5A)(MILK OR MAMMARY)/AB,BI

=> s l10(10a)(function? or assembl?)/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 FUNCTION?/AB
838844 FUNCTION?/BI
0 ASSEMBL?/AB
34637 ASSEMBL?/BI
L12 2 L10(10A)(FUNCTION? OR ASSEMBL?)/AB,BI

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 2 ANSWERS -
CONTINUE? Y(N),y

L12 ANSWER 1 OF 2 MEDLINE
AN 1998455662 MEDLINE
DN 98455662

TI Isolation and recombinant expression of an MHV-JHM
neutralising monoclonal
antibody.

AU Kolb A F, Lechermaier M; Heister A, Toksoy A, Siddell S G
CS Institute of Virology and Immunology, University of Wurzburg,
Germany.
SO ADVANCES IN EXPERIMENTAL MEDICINE AND
BIOLOGY, (1998) 440 657-64.

Journal code: 2LU. ISSN: 0065-2598.

CY United States
DT Journal; Article; (JOURNAL ARTICLE)

LA English
FS Priority Journals

EM 199903
EW 19990303

AB The monoclonal antibody A1 (mab A1) efficiently neutralises the
infection of susceptible cells by the murine hepatitis virus MHV-JHM in
vitro and in vivo (Wege et al., 1984). The variable regions of mab A1 were
amplified from mRNA of the respective hybridoma cell line by RT-PCR and
integrated into different eukaryotic expression vectors. The biological
function of the ***recombinant*** ***antibody***
constructs was verified by virus neutralisation assays.

Whereas a complete recombinant antibody (mab A1 rec.) expressed in

transfected murine myeloma cells inhibited the MHV-JHM infection as well as the parental antibody, a single-chain Fv derived from mab A1 did not show any neutralising activity.

L12 ANSWER 2 OF 2 MEDLINE
AN 97041563 MEDLINE
DN 97041563

TI Lung cancer-reacting human recombinant antibody AE6F4:
potential

usefulness in the sputum cytodiagnosis

AU Shoji M; Kawamoto S; Seki K; Teruya K; Setoguchi Y;
Mochizuki K; Kato M;

Hashizume S; Hanagiri T; Yoshimatsu T; Nakanishi K; Yasumoto
K; Nagashima

A; Nakahashi H; Suzuki T; Imai T; Shirahata S; Nomoto K;
Murakami H

CS Morinaga Institute of Biological Science, Yokohama, Japan.
SO HUMAN ANTIBODIES AND HYBRIDOMAS, (1996) 7 (1)
27-36.

Journal code: A6A. ISSN: 0956-960X.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199706

EW 19970601

AB Human monoclonal antibody (hMAb) AE6F4 has been shown to
be potentially useful for immunocytological detection of lung cancer cells in
sputum. By

recombinant DNA technology, IgM type hMAb AE6F4 was
switched to IgG. The
IgG mimic ***recombinant*** AE6F4 ***antibody***
expression

plasmid was ***assembled*** using the
antibody heavy
chain gene, which ligated the gene encoding VH and CH1(mu)

domains of hMAb
AE6F4 heavy chain to the gene encoding CH2(gamma 1) and
CH3(gamma 1)

domains of human IgG heavy chain, and the antibody light chain
gene of

hMAb AE6F4. The recombinant antibody expressed by baby
hamster kidney

(BHK)-21 cells showed molecular size equivalence to IgG, and
consisted of
human mu-gamma hybrid heavy and kappa light chains. The
immunological

specificity of the recombinant antibody was the same as that of
hMAb AE6F4

by immunoblotting analysis to the 14-3-3 protein, the putative
antigen of

hMAb AE6F4, and by immunohistochemical and
immunocytological analyses

using tissue sections and sputa of lung cancer patients. The

transfected
BHK-21 cells produced the recombinant antibody persistently and
the
productivity was greater than 20 times that by human-human
hybridoma
producing hMAb AE6F4.

=> s l2(5a)(mammary)(w)(tumor or tumour)/ab,bi

'AB' IS NOT A VALID FIELD CODE
36573 MAMMARY
0 TUMOR/AB
374204 TUMOR/BI
0 TUMOUR/AB
64119 TUMOUR/BI
L13 0 L2(5A)(MAMMARY)(W)(TUMOR OR
TUMOUR)/AB,BI

=> s l1(5a)(mammary)(w)(tumor or tumour)/ab,bi

'AB' IS NOT A VALID FIELD CODE
36573 MAMMARY
0 TUMOR/AB
374204 TUMOR/BI
0 TUMOUR/AB
64119 TUMOUR/BI
L14 74 L1(5A)(MAMMARY)(W)(TUMOR OR
TUMOUR)/AB,BI

=> s l14(5a)(promoter#)/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 (PROMOTER#)/AB
66697 (PROMOTER#)/BI
L15 0 L14(5A)(PROMOTER#)/AB,BI

=> s l14(5a)(virus)/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 (VIRUS)/AB
275885 (VIRUS)/BI
L16 42 L14(5A)(VIRUS)/AB,BI

=> s l16 and milk/ab,bi

'AB' IS NOT A VALID FIELD CODE
0 MILK/AB
48429 MILK/BI
L17 2 L16 AND MILK/AB,BI

=> d 1 - bib ab

YOU HAVE REQUESTED DATA FROM 2 ANSWERS -
CONTINUE? Y(N)?y

L17 ANSWER 1 OF 2 MEDLINE
AN 81061682 MEDLINE
DN 81061682
TI Ubiquity of natural ***antibodies*** to the ***mammary***

tumour ***virus*** in mice.
AU Benivelzen P; Brinkhof J
NC NO1 CP43328 (NCI)
SO ARCHIV FUR GESCHWULSTFORSCHUNG, (1980) 50 (3)
193-203.

Journal code: 746. ISSN: 0003-911X.
CY GERMANY, EAST: German Democratic Republic
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 198103
AB Sera of female and male mice from eleven inbred mouse strains
collected at

either 4, 12, 36 or 60 weeks of age were tested for the presence of
natural ***antibodies*** to the murine ***mammary***
tumour ***virus*** by means of the Sepharose bead
immunofluorescence assay. Antibodies to the virus proved to be
ubiquitous,
but pronounced strain differences were found in titer and onset of
antibody production. These differences were related to neither
release of

virus in the ***milk*** nor susceptibility to spontaneous
mammary
tumour development of a given strain. Immunological specificity of

the
observed reactions was concluded from a) the failure to block the
reaction
by absorption with fetal calf serum, mouse ***milk*** or sheep
erythrocytes, while absorption with purified virus abolished the
reactivity; b) the lack of reactivity of rat sera with the mouse
mammary

tumour virus in this system; c) the negative response of mouse sera
with
Sepharose beads coated with ovalbumin; d) the lack of correlation
between

antibody titers to Rauscher murine leukemia virus and mammary
tumour virus
in this system; e) the retaining of activity to highly purified viral
polypeptides; f) blocking of the reaction by preincubation with
rabbit

anti-mouse immunoglobulin serum or Protein A from
Staphylococcus aureus.
Since germfree mice of various strains also have such antibodies, it

is
concluded that the reactions are not due to horizontal transmission
of the

virus. From the lack of correlation between antibody titers and
tumour
incidences, it is concluded that various systems overshadow the
potential
immunosurveillance role of such natural antiviral antibodies.

L17 ANSWER 2 OF 2 MEDLINE
AN 76137969 MEDLINE
DN 76137969

TI Human ***antibodies*** binding to the mouse
mammary
tumour ***virus*** : a nonspecific reaction?
AU Newgard K W; Cardiff R D; Blair P B
SO CANCER RESEARCH, (1976 Feb) 36 (2 pt 2) 765-8.
Journal code: CNF. ISSN: 0008-5472.

CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 197607
AB Specific rabbit antisera and over 100 human sera were found to
precipitate
iodinated mouse mammary tumor virus (MTV). The specificity of
these

reactions was tested in competitive inhibition studies. Three classes
of

reaction could be distinguished. The Class 1 reaction was the most
specific; it could be inhibited only by MTV and was observed
exclusively

with rabbit anti-MTV. The Class 2 reaction was apparently against
mouse

cell determinants; it could be inhibited not only by MTV but also
by mouse

lactating mammary gland and was characteristic of rabbit
anti-mouse

lactating mammary gland. The Class 3 reaction was the least
specific; it

could be inhibited not only by MTV and mouse lactating mammary
gland but
also by dog ***milk***. All of the human sera tested exhibited
Class 3
reactivities.

=> file medline embase biosis inpadoc caplus

COST IN U.S. DOLLARS	ENTRY	SINCE FILE	TOTAL
FULL ESTIMATED COST	SESSION	5.22	5.37

FILE 'MEDLINE' ENTERED AT 10:28:48 ON 10 AUG 1999

FILE 'EMBASE' ENTERED AT 10:28:48 ON 10 AUG 1999
COPYRIGHT (C) 1999 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 10:28:48 ON 10 AUG 1999
COPYRIGHT (C) 1999 BIOSIS(R)

FILE 'INPADOC' ENTERED AT 10:28:48 ON 10 AUG 1999
COPYRIGHT (C) 1999 European Patent Office, Vienna (EPO)

FILE 'CAPLUS' ENTERED AT 10:28:48 ON 10 AUG 1999
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER

AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 1999 AMERICAN CHEMICAL SOCIETY (ACS)

=> s 117

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L18 13 L17

=> dup rem 118

PROCESSING COMPLETED FOR L18
L19 10 DUP REM L18 (3 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 10 ANSWERS -
CONTINUE? Y(N)y

L19 ANSWER 1 OF 10 CAPLUS COPYRIGHT 1999 ACS
AN 1990:589384 CAPLUS
DN 113:189384
TI Mouse mammary tumor virus (MMTV) infection in SWISS and RIII mice.

Correlation between resistance to exogenous infection and anti-MMTV serum

response
AU Hainaut, P.; Vaira, Dolores; Francois, Camille; Calberg-Bacq, Claire

CS Inst. Pathol., Univ. Liege, Liege, B-4000, Belg.
SO Arch. Virol. (1990), 113(1-2), 35-52

CODEN: ARVIDF; ISSN: 0304-8608

DT Journal
LA English

AB Host-virus relationships were examd. in mice from the mouse mammary tumor virus (MMTV)-infected strains SWISS MB+ and RIII, which harbor the same

MMTV variant, and from the derived sublines Swiss MB- and RIIIf, which were freed of ***milk*** -borne MMTV by foster-nursing.

These 2 strains are not phylogenetically related, the SWISS strain bearing the endogenous Mtv-3 locus in its DNA. In RIII and SWISS MB+ mice, the

incidence of early mammary tumors, which was of 96% and 8%, resp., was correlated to the level of MMTV expression in ***milk***. In

the SWISS MB-line, a non-coordinate expression of the provirus assocd. with

the Mtv-3 locus was obsd. in the mammary glands, the salivary glands, and the spleen. This expression was not tumorigenic and was characterized by the presence of the p28 gag antigen and the absence of gp52 env antigen, except, however, in mammary glands of elder mice where traces of gp52 were found. In the mammary glands of SWISS MB+ mice, the expression of the

Mtv-3 locus was masked by large amts. of antigens resulting from exogenous

virus expression. RIIIf mice were MMTV-neg. Viral antigens coexisted

with anti-MMTV antibodies in the serum of infected and tumor-bearing mice, but not in the form of immune complexes. An anti-MMTV serum reactivity

was also detected in SWISS MB- and RIIIf mice. However, the response was higher in the 2 SWISS lines than in the 2 RIII lines. Except in tumor-bearing mice, the anti-MMTV response was not modified by the

presence of exogenous virus and thus resulted essentially from exposure to

endogenous MMTV expression. In expl. infection studies, RIII mice were

more susceptible to MMTV infection than SWISS mice. The correlation

between resistance to MMTV infection and serum response to endogenous MMTV

expression, suggests that the non-tumorigenic expression of an provirus can protect at least partially, against exogenous MMTV infection.

L19 ANSWER 2 OF 10 BIOSIS COPYRIGHT 1999 BIOSIS
AN 1988:505990 BIOSIS
DN BA86:126674

TI THE PRESENCE OF ANTIBODIES SPECIFIC FOR MMTV STRUCTURAL PROTEINS AMONG ANTIBODIES FROM CIRCULATING IMMUNE COMPLEXES OF BREAST CANCER PATIENTS.

AU LITVINOV S V; MALIVANOVA T F; CHUEV YU V; KRYUKOVA I N

CS ALL-UNION ONCOL. SCI. CENT., ACAD. MED. SCI. USSR, MOSCOW, USSR.

SO BYULL EKSP BIOL MED, (1988) 105 (4), 475-477.

CODEN: BERMAE; ISSN: 0365-9615.

FS BA: OLD

LA Russian

AB Circulating immune complexes were precipitated from breast cancer

patients' sera using 2.5% polyethylenglycol. CIC isolated from 70 ml of

sera from 15 patients were dissociated and

immunoglobulin-containing fraction was prepared by chromatography on Sephadex G-200 column. The

fraction contained IgG specific for MuMTV structural proteins, as revealed

by ELISA. CIC preparations from 22 sera of breast cancer patients were

digested with pepsin; Fab' fragment preparations were also analysed by

ELISA, only one of them was MMTV-specific. IgG and Fab' fragments isolated

from CIC reacted specifically with MMTV proteins, the reaction was not

blocked by virus-free murine ***milk*** or other retroviruses (Ra-MuLV

and MPMV).

L19 ANSWER 3 OF 10 BIOSIS COPYRIGHT 1999 BIOSIS

AN 1985:278669 BIOSIS

DN BA79:58665

TI AUTOCHTHONOUS HUMORAL IMMUNE RESPONSES TO EXOGENOUS AND ENDOGENOUS MURINE

MAMMARY TUMOR VIRUSES IN C-3H JAX AND ICRC MICE.

AU CHIPLUNKAR S V; GANGAL S G; KARANDE K A CS IMMUNOL. DIV. CANCER RES. INST., TATA MEML.

CENT., BOMBAY 400 012, INDIA.

SO INDIAN J EXP BIOL, (1984 (RECD 1985)) 22 (12), 662-665. CODEN: IJEBAA6; ISSN: 0019-5189.

FS BA: OLD

LA English

AB Autochthonous humoral ***antibody*** response directed against murine ***mammary*** ***tumor*** ***virus*** (MuMTV) in

sera of high mammary tumor strains of mice such as C3H(Jax), ICRC and ICRC

breeders and their low tumor incidence sublines C3H (Mect) and ICRCf

carrying only endogenous virus, were estimated by radioimmunoprecipitation

technique using 125I-labeled C3H MuMTV. Sera were obtained from normal

mice of various age groups, parity and lactation stage, which corresponded

to the amounts of MuMTV in the ***milk*** and also from mammary tumor

bearing mice. Highest levels of MuMTV antibodies were observed in mammary

tumor bearing mice carrying both endogenous and exogenous viruses. Mice

carrying only endogenous virus had low amounts of antibodies cross reacting with exogenous MuMTV of C3H (Jax). A sequential

increase in MuMTV

antibodies was seen in normal mice, which preceded the mammary tumor

development.

breast tissues. With 1 exception, 99 carcinomas from 13 organs other than breast and 8 cystosarcomas were all negative.

L19 ANSWER 6 OF 10 EMBASE COPYRIGHT 1999 ELSEVIER SCI. B.V.
 AN 79069978 EMBASE
 DN 1979069978
 TI Naturally occurring humoral immunity to murine mammary tumor virus (MuMTV) and MuMTV GP52 in mice with low mammary tumor incidence. AU Arthur L.O.; Fine D.L.
 CS Viral Oncol. Progr., Frederick Cancer Res. Cent., Frederick, Md. 21701, United States
 SO International Journal of Cancer, (1978) 22/6 (734-740). CODEN: IJGNAW
 CY Switzerland
 DT Journal
 FS 016 Cancer
 026 Immunology, Serology and Transplantation
 047 Virology
 LA English
 SL French
 AB ***Antibodies*** to murine ***mammary***
 tumor
 virus (MuMTV) were found in sera from male and female mice by means of a radiolabelled intact MuMTV precipitation assay. These antibodies were demonstrated both in strains of mice that have a high incidence of mammary tumors and transmit the highly oncogenic MuMTV via the ***milk*** (C3H) and in mice that have been foster-nursed to remove the highly oncogenic ***milk*** -borne MuMTV (C3Hf) and subsequently have a decreased mammary tumor incidence. Antibodies to MuMTV were readily demonstrated in C3H mice at 6 weeks of age, whereas only marginal antibody activity was detected in C3Hf mice of less than 15 weeks of age. Antibody levels increased with age in both strains, but in C3Hf mice the immune response was also accelerated by pregnancy. Several feral and BALB/c NIV mouse sera with high (125I)MuMTV precipitating titers also precipitated the major MuMTV envelope glycoprotein (gp52). Maximum precipitation of the (125I)gp52 was >50% with a 20% endpoint binding titer of 1:40, whereas the same sera precipitated >80% of (125I)MuMTV with a titer of >1:2560. These naturally occurring antibodies were specific for

immunosurveillance role of such natural antiviral antibodies.

L19 ANSWER 5 OF 10 EMBASE COPYRIGHT 1999 ELSEVIER SCI. B.V.DUPLICATE 2
 AN 78304798 EMBASE
 DN 1978304798
 TI Detection in human breast carcinomas of an antigen immunologically related to a group-specific antigen of mouse mammary tumor virus. AU Mesa-Tejada R.; Keydar I.; Ramnarayanan M.; et al.
 CS Inst. Cancer Res., Coll. Physcns Surg., Columbia Univ., New York, N.Y.
 10032, United States
 SO Proceedings of the National Academy of Sciences of the United States of America, (1978) 75/3 (1529-1533). CODEN: PNASA6
 CY United States
 DT Journal
 FS 016 Cancer
 010 Obstetrics and Gynecology
 LA English
 AB An antigen immunologically related to a group-specific antigen (gp52, a 52,000-dalton glycoprotein) of the mouse mammary tumor virus has been identified in paraffin sections of human breast cancers by means of the indirect immunoperoxidase technique. The specificity of the reaction with ***antibody*** against mouse ***mammary***
 tumor
 virus was examined by absorption of the IgG with the following: purified gp52, a number of virus preparations (mouse mammary tumor virus, Rauscher leukemia virus, simian sarcoma virus, baboon endogenous virus, and Mason-Pfizer monkey virus); normal plasma, leukocytes, breast tissue, ***milk***, actin, collagen, and hyaluronic acid, all of human origin; sheep erythrocytes and mucin. Only mouse mammary tumor virus (from C3H or Paris RIII strains and grown in either murine or feline cells) and purified gp52 eliminated the immunohistochemical reaction in the human breast tumors. Positive reactions were seen in 51 of 131 (39%) carcinomas of various histologic types, a minimal estimate in view of the limited number of sections from each tumor that could be examined. Negative reactions were obtained in all 119 benign breast lesions (cystic disease, fibroadenoma, papilloma, gynecomastia) and in all 18 normal

DUPLICATE

L19 ANSWER 4 OF 10 MEDLINE
 AN 81061682 MEDLINE
 DN 81061682
 TI Ubiquity of natural ***antibodies*** to the ***mammary***
 tumor
 virus in mice. AU Bentvelzen P.; Brinkhof J.
 NC NOI CP43328 (NCI)
 SO ARCHIV FUR GESCHWULSTFORSCHUNG, (1980) 50 (3) 193-203.
 Journal code: 746. ISSN: 0003-911X.
 CY GERMANY, EAST: German Democratic Republic
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 198103
 AB Sera of female and male mice from eleven inbred mouse strains collected at either 4, 12, 36 or 60 weeks of age were tested for the presence of natural ***antibodies*** to the murine ***mammary***
 tumor
 virus by means of the Sepharose bead immunofluorescence assay. Antibodies to the virus proved to be ubiquitous, but pronounced strain differences were found in titer and onset of antibody production. These differences were related to neither virus in the ***milk*** nor susceptibility to spontaneous mammary tumour development of a given strain. Immunological specificity of the observed reactions was concluded from a) the failure to block the reaction by absorption with fetal calf serum, mouse ***milk*** or sheep erythrocytes, while absorption with purified virus abolished the reactivity; b) the lack of reactivity of rat sera with the mouse mammary tumour virus in this system; c) the negative response of mouse sera with Sepharose beads coated with ovalbumin; d) the lack of correlation between antibody titers to Rauscher murine leukemia virus and mammary tumour virus in this system; e) the retaining of activity to highly purified viral polypeptides; f) blocking of the reaction by preincubation with rabbit anti-mouse immunoglobulin serum or Protein A from Staphylococcus aureus. Since germfree mice of various strains also have such antibodies, it is concluded that the reactions are not due to horizontal transmission of the virus. From the lack of correlation between antibody titers and tumour incidences, it is concluded that various systems overshadow the potential

gp52 since only MuMTV and purified MuMTV gp52 competed for binding of radiolabelled antigens by limiting dilutions of mouse sera. MuMTV gp52 was found in both C3H and C3Hf mice, predominantly in organs which had secretory functions, such as submaxillary glands and mammary tissue of females and submaxillary, coagulating, and vesicular glands and vas deferens of males. Extracts of other tissues were negative for MuMTV gp52. Neither gp52 nor naturally occurring antibodies for MuMTV were found in BALB/c or C57BL/6 mice.

L19 ANSWER 7 OF 10 CAPLUS COPYRIGHT 1999 ACS
 AN 1979:37410 CAPLUS
 DN 90:37410
 TI Modulation of mouse mammary tumor virus production in the MJY-alpha cell line
 AU Yagi, Mary Jane; Blair, Phyllis B.; Lane, Mary Ann
 CS Sch. Med., Univ. Alabama, Birmingham, Ala., USA
 SO J. Virol. (1978), 28(2), 611-23
 CODEN: JOVIAM; ISSN: 0022-538X
 DT Journal
 LA English
 AB Implantation of mouse mammary tumor virus (MMTV)-producing mammary tumor cell line MJY-alpha into isogenic mice elicited both humoral and T-cell response against MMTV virion antigens. The carcinomas developed from the implanted cells showed a decrease in MMTV synthesis, compared with cells remaining in culture, which was detectable at gnotex 7 days after implantation and for 5 transplant generations. Electron microscopic examn. of thin sections of the tumors revealed that intracytoplasmic A particles, budding particles, and cell-free MMTV B particles were all affected. However, immunofluorescence assays of tumor sections demonstrated the presence of MMTV viral antigens in the cells. Cell cultures initiated from 1st-, 3rd-, and 4th-generation tumors were morphol. identical to the original in vitro cell line, although virus prodn. was barely detectable. Anal. of the cultures by electron microscopy revealed an increase in MMTV virions after 3 in vitro passages. Polypeptide profiles obtained by Na dodecyl sulfate-polyacrylamide gel electrophoresis of virions purified from these cultures were identical to MMTV. Immunodiffusion demonstrated the cross-reactivity between these

virions and MMTV particles obtained from mouse ***milk***
 In vitro treatment of MJY-alpha cell cultures with rabbit anti-MMTV antiserum resulted in a redn. of extracellular MMTV virions, as well as alterations in their Na dodecyl sulfate-polyacrylamide gel electrophoretic polypeptide patterns.
 L19 ANSWER 8 OF 10 MEDLINE
 AN 76137969 MEDLINE
 DN 76137969
 TI Human ***antibodies*** binding to the mouse ***mammary***
 tumor ***virus*** : a nonspecific reaction?.
 AU Newgard K.W.; Cardiff R.D.; Blair P.B.
 SO CANCER RESEARCH, (1976 Feb) 36 (2 pt 2) 765-8.
 Journal code: CNF. ISSN: 0008-5472.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 197607
 AB Specific rabbit antiserum and over 100 human sera were found to precipitate iodinated mouse mammary tumor virus (MTV). The specificity of these reactions was tested in competitive inhibition studies. Three classes of reaction could be distinguished. The Class 1 reaction was the most specific; it could be inhibited only by MTV and was observed exclusively with rabbit anti-MTV. The Class 2 reaction was apparently against mouse cell determinants; it could be inhibited not only by MTV but also by mouse lactating mammary gland and was characteristic of rabbit anti-mouse lactating mammary gland. The Class 3 reaction was the least specific; it could be inhibited not only by MTV and mouse lactating mammary gland but also by dog ***milk***. All of the human sera tested exhibited reactivities.
 L19 ANSWER 9 OF 10 EMBASE COPYRIGHT 1999 ELSEVIER
 SCI B.V.
 AN 77020594 EMBASE
 DN 1977020594
 TI Human ***antibodies*** binding to the mouse ***mammary***
 tumor ***virus*** : a nonspecific reaction?.
 AU Newgard K.W.; Cardiff R.D.; Blair P.B.
 CS Dept. Pathol., Sch. Med., Univ. California, Davis, Calif., United States

SO Cancer Research, (1976) 36/2 (II) (765-768).
 CODEN: CNREA8
 DT Journal
 FS 016 Cancer
 047 Virology
 026 Immunology, Serology and Transplantation
 005 General Pathology and Pathological Anatomy
 009 Surgery
 LA English
 AB Specific rabbit antiserum and over 100 human sera were found to precipitate iodinated mouse mammary tumor virus (MTV). The specificity of these reactions was tested in competitive inhibition studies. Three classes of reaction could be distinguished. The Class 1 reaction was the most specific; it could be inhibited only by MTV and was observed exclusively with rabbit anti MTV. The Class 2 reaction was apparently against mouse cell determinants: it could be inhibited not only by MTV but also by mouse lactating mammary gland and was characteristic of rabbit anti lactating mammary gland. The Class 3 reaction was the least specific; it could be inhibited not only by MTV and mouse lactating mammary gland but also by dog ***milk***. All of the human sera tested exhibited reactivities.
 L19 ANSWER 10 OF 10 EMBASE COPYRIGHT 1999
 ELSEVIER SCI B.V.
 AN 74115184 EMBASE
 DN 1974115184
 TI [Virus induced mammary carcinoma of mice: a genuine model of cancer in man].
 DAS VIRUSINDUZIERTE MAMMAKARZINOM DER MAUS
 - EIN ECHTES MODELL FUR DEN BRUSTKREBS DES MENSCHEN?
 AU Zotter S.; Muller M.
 CS Pathol. Inst., Med. Akad. 'Carl Gustav Carus', Dresden, Germany
 SO Deutsche Gesundheitswesen, (1973) 28/41 (1936-1942).
 CODEN: DEGEA3
 DT Journal
 FS 005 General Pathology and Pathological Anatomy
 047 Virology
 016 Cancer
 LA German
 AB For some years the possible existence of a human mammary cancer inducing virus has been seriously discussed. This assumption is based on the detection of characteristic virus particles resembling the mammary tumor virus of mice, as well as of tumor virus specific enzymes and

nucleic acids in women's ***milk*** and mammary cancer tissue. The results of immunologic studies, e.g. the author's own observation of specific ***antibodies*** directed against the ***mammary*** ***tumor*** ***virus*** in the serum, above all, of women suffering from mammary cancer and from mastopathies, complete these findings. The occurrence of antibodies in healthy women, as well as men and children might be the expression of a wide spreading of the hypothetical human mammary cancer virus. The occurrence of antibodies in infected people might be as analogous to the murine mammary tumor virus. As in mice a mammary tumor virus is invariably associated with formation of antibodies. Providing a confirmation and completion of these findings, also under the aspect of comparative tumor virology, the virus induced mammary carcinoma of mice might well be regarded as a natural animal experimental model for human cancer.

=> s 13

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L20 11 L3

=> dup rem 120

PROCESSING COMPLETED FOR L20
L21 7 DUP REM L20 (4 DUPLICATES REMOVED)

=> d 1 - bib ab

YOU HAVE REQUESTED DATA FROM 7 ANSWERS - CONTINUE? Y(N)?

L21 ANSWER 1 OF 7 CAPLUS COPYRIGHT 1999 ACS
AN 1998-590656 CAPLUS
DN 129-229676
TI Modified antibodies with human milk fat globule specificity for breast cancer diagnosis and therapy
IN Do Couto, Fernando J. R.; Ceriani, Roberto L.; Peterson, Jerry A.
PA Cancer Research Fund of Contra Costa, USA
SO U.S., 76 pp. Cont.-in-part of U.S. Ser. No. 977,696.

CODEN: USXXAM
DT Patent
LA English
FAN/CNT 2
PATENT NO. KIND DATE APPLICATION NO.
DATE

PI US 5804187 A 19980908 US 1993-129930 19930930
US 5792852 A 19980811 US 1992-977696 19921116
WO 9411509 A2 19940526 WO 1993-US11445
19931116
WO 9411509 A3 19940707
W: AT, AU, BB, BG, BR, CA, CH, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, PL, RO, RU, SD, SE, US
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BI, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG

CA 2149529 AA 19940526 CA 1993-2149529 19931116
AU 9463964 A1 19940608 AU 1994-63964 19931116
EP 674710 A1 19951004 EP 1994-903300 19931116
R: DE, ES, FR, GB, IE, IT, NL, SE
JP 09503901 T2 19970422 JP 1993-512520 19931116
PRAI US 1992-977696 19921116
US 1993-129930 19930930
US 1993-134346 19931008
WO 1993-US11445 19931116
AB An analog peptide that comprises the variable regions of the light or heavy chains of an antibody of a first species selectively binding to a carcinoma antigen has 1 to 46 amino acids of the framework regions per chain substituted with amino acids such as those present in equiv. positions in antibodies of a species other than the first species, or fragments thereof comprising 1 to 3 variable region CDRs per chain and optionally flanking regions thereof of 1 to 10 or more amino acids, alone or with an N-terminal fragment of 1 to 10 or more amino acids, combinations or mixts. thereof. The polypeptide may also comprise an effector agent and/or be glycosylated, and is presented as a compn. with a carrier. The analog peptides are used in diagnostic kits for carcinomas and methods for in vivo imaging and treating a primary or metastasized carcinoma, and in vitro diagnosing a carcinoma, ex vivo purging neoplastic cells from a biol. fluid. RNAs and DNAs encode the analog peptide, and a hybrid vector carrying the nucleotides and transfected cells express the peptides and a method produces the analog peptide. An anti-idiotype

polypeptide comprises polyclonal antibodies raised against an anti-carcinoma antibody or the analog peptide of this invention, monoclonal antibodies thereof, Fab, Fab' (Fab')₂, CDR, variable region, or analogs or fragments thereof, combinations thereof with an oligopeptide comprising a TRP primer, tandem repeats thereof, or combination or mixts. thereof. An anti-idiotype hybrid polypeptide with an effector agent and the anti-idiotype polypeptide, an anti-carcinoma vaccine, an anti-carcinoma vaccination kit, a method of vaccinating against carcinoma and a method of lowering the serum concn. of a circulating antibody or polypeptide are provided.

L21 ANSWER 2 OF 7 BIOSIS COPYRIGHT 1999 BIOSIS
AN 1995237460 BIOSIS
DN PREV199598251760
TI Role of human ***milk*** ***antibodies*** against specific invasion ***plasmid*** antigens (Ipas) of Shigella flexneri in protection against invasion of HeLa cells.
AU Gomez, Henry F. (1); Forbes, Cheryl; Medellin, Christopher D.; Pickering, Larry K.; Cleary, Thomas G.
CS (1) Dep. Peds, Univ. Texas Med. Sch., Houston, TX USA
SO Pediatric Research, (1994) Vol. 37, No. 4 PART 2, pp. 175A.
Meeting Info.: 105th Annual Meeting of the American Pediatric Society and the 64th Annual Meeting of the Society for Pediatric Research San Diego, California, USA May 7-11, 1995
ISSN: 0031-3998.
DT Conference
LA English

L21 ANSWER 3 OF 7 CAPLUS COPYRIGHT 1999 ACS
AN 1993-407220 CAPLUS
DN 119-7220
TI Humanized antibodies to human milk fat globules
IN Adair, John Robert; Hamann, Philip R.; Owens, Raymond John; Baker, Terence
Seward, Lyons, Alan Howard; Hinman, Lois M.; Menendez, Ana T.
PA Celtech Ltd., UK
SO Eur. Pat. Appl., 59 pp.
CODEN: EPXXDW
DT Patent
LA English
FAN/CNT 1
PATENT NO. KIND DATE APPLICATION NO.
DATE

PI EP 534742 A1 19930331 EP 1992-308680 19920924
EP 534742 BI 19971119

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU,
MC, NL, PT, SE
CA 1992-2095926 19920924
WO 9306231 A1 19930401 WO 1992-GB1759
19920924
W: AU, CA, CS, FI, HU, JP, KR, NO
AU 9225983 A1 19930427 AU 1992-25983 19920924
B2 19960229
EP 666868 A2 19970702 EP 1997-200482 19920924
EP 781845 A3 19970709
EP 781845 A3 19970709
MC, NL, PT, SE
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU,
AU 160362 E 19971215 AT 1992-308680 19920924
ES 103872 T3 19980101 ES 1992-308680 19920924
IL 103269 A1 19980104 IL 1992-103269 19920924
PRAI GB 1991-20467 19910926
EP 1992-308680 19920924
WO 1992-GB1759 19920924
AB Chimeric and complementarity-detg. region (CDR)-grafted
humanized
antibodies to human milk fat globules are prepd. for use in the
diagnosis
and treatment of breast cancer. The CDRs are derived from the
mouse
IgG1.kappa. monoclonal antibody CTMO1 that recognizes an
antigen found in
high levels in blood of breast cancer patients. The antibody may be
conjugated with antitumor agents for treatment of the disease. The
genes
for the humanized antibodies were constructed by std. methods and
expressed in CHO-L761 cells. Binding and internalization of the
humanized
antibodies by breast carcinoma cell lines was demonstrated.
Conjugation
of the antibodies with calicheamicin .gamma. II was demonstrated.
L21 ANSWER 4 OF 7 EMBASE COPYRIGHT 1999 ELSEVIER
SCI. B. V.
AN 94031560 EMBASE
DN 1994031560
TI ***Antibodies*** to Shigella lipopolysaccharides and invasion
plasmid antigens in colostrum and breast ***milk***
of women
from Puriscal, a rural area of Costa Rica.
AU Achi R.A.; Vives M.; Garcia M.E.; Binh Minh N.; Mata L.;
Lindberg A.A.
CS Department of Clinical Bacteriology, Huddinge Hospital,
Karolinska
Institute,S-141 81 Huddinge, Sweden
SO Serodiagnosis and Immunotherapy in Infectious Disease, (1993)
5/4
(237-244)
ISSN: 0888-0786 CODEN: SIIDE3
CY United Kingdom
DT Journal: Article
FS 004 Microbiology
017 Public Health, Social Medicine and Epidemiology

026	Immunology, Serology and Transplantation	LA English SL AB healthy enzyme immunoassays. Mean relative IgA titres of 790 \pm 790 \pm 640 (ranges = 70-3250), 440 \pm 490 (40-2940), and 280 \pm 230 (10-1000) to Shigella flexneri, Shigella sonnei and Shigella dysenteriae type 1 lipopolysaccharides (LPSs), respectively, were found in colostrum on day 1 post delivery. Titres declined thereafter and only relatively low values were found on days 30, 90 and 180 post partum. Mean IgA anti-invasion plasmid antigens (Ipa) titres of 200 \pm 230, 140 \pm 170 and 120 \pm 190 on days 1-2, 3-4 and 5-8 post delivery were found, respectively. Thereafter the anti-Ipa titres were low. IgM antibody titres were found against the LPS antigens but not to the Ipa. IgG antibody titres were low against both LPS and Ipa, with the exception of four out of 59 (7%) at day 1, with titres to the S. flexneri Y LPS and three out of 59 (5%) at day 1 and one out of 87 (1%) at days 3-4, with positive IgG titres to Ipa. By using a cut off value (mean \pm 2 sD established for a high socioeconomic status group of Costa Rica), 20%, 17% and 5% of colostrum samples, at day 1, had high titres to S. flexneri, S. sonnei and S. dysenteriae, respectively. A good degree of correlation between colostrum IgA anti-S. flexneri Y LPS and the anti-Ipa antibodies, was found. The colostrum antibody titres (day 1) of mothers from Puriscal were intermediate as compared to mothers from the low and the high socioeconomic conditions in the metropolitan area of Costa Rica and tended to be lower than in Vietnamese mothers from endemic areas of shigellosis. Shigella exposure seems to be lower in this rural region than in overcrowded slums of San Jose, Costa Rica.
L21	ANSWER 5 OF 7 MEDLINE	DUPLICATE 1
AN	93078105 MEDLINE	
DN	93078105	
TI	Concentration of ***milk*** secretory ***immunoglobulin*** A	
		against Shigella virulence ***plasmid*** -associated antigens as

a predictor of symptom status in Shigella-infected breast-fed infants.
AU Hayani K C; Guerrero M L; Morrow A L; Gomez H F; Winsor D K; Ruiz-Palacios G M; Cleary T G
CS Department of Pediatrics, University of Texas Medical School, Houston 77030.
NC 55-PO1-HD-13021 (NICHHD)
SO JOURNAL OF PEDIATRICS. (1992 Dec) 121 (6) 852-6.
Journal code: JIJL ISSN: 0022-3476.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals
EM 199303
AB We conducted a prospective, community-based study of healthy breast-fed Mexican infants to determine the protective effects of anti-Shigella secretory IgA antibodies in milk. Milk samples were collected monthly, and stool culture specimens were obtained weekly and at the time of episodes of diarrhea. Nineteen breast-fed infants were found to have Shigella flexneri, Shigella boydii, or Shigella sonnei in stool samples. Ages of the 10 infants with symptomatic infection and the nine with asymptomatic infection did not differ significantly. Milk samples collected up to 12 weeks before infection were evaluated by enzyme-linked immunosorbent assay for secretory IgA antibodies against lipopolysaccharides of S. flexneri, S. boydii serotype 2, S. sonnei, and virulence plasmid-associated antigens. The geometric mean titers of anti-Shigella ***antibodies*** to virulence ***plasmid***-associated antigens in ***milk*** received before infection were eightfold higher in infants who remained well than in those in whom diarrhea developed. The significance of milk secretory IgA directed against lipopolysaccharide was less clear. We conclude that human milk protects infants against symptomatic shigella infection when it contains high concentrations of secretory IgA against virulence plasmid-associated antigens.

L21 ANSWER 6 OF 7 CAPLUS COPYRIGHT 1999 ACS
AN 1993:145584 CAPLUS
DN 118:145584
TI Milk secretory IgA related to Shigella virulence antigens
AU Cleary, Thomas G.; Hyani, Karen; Winsor, Donald K.;

Ruiz-Palacios, Guillermo
CS Med. Sch., Univ. Texas, Houston, TX, USA
SO Adv. Exp. Med. Biol. (1991), 310(Immunol. Milk Neonate), 369-73
CODEN: AEMBAP, ISSN: 0065-2598
DT Journal
LA English
AB Human milk commonly contains antibodies to the major virulence antigens shared by all Shigellae. The levels of these antibodies in milk do not change significantly during lactation either in a high (Mexico) or low Shigella risk population (US). The presence of ***antibodies*** to Shigella virulence ***plasmid*** -coded antigens in the ***milk*** of women from an area where Shigella infection is not common suggests that Shigella-specific, IgA-secreting cells which have been programmed in the distant past are recruited to the breast during pregnancy or lactation.

L21 ANSWER 7 OF 7 MEDLINE DUPLICATE 2
AN 91093893 MEDLINE
DN 91093893
TI Human ***milk*** secretory ***immunoglobulin*** A to Shigella virulence ***plasmid*** -coded antigens.
AU Cleary T G; West M S; Ruiz-Palacios G; Winsor D K; Calva J J; Guerrero M
L, Van R
CS Department of Pediatrics and Microbiology, University of Texas Medical School at Houston 77030.
NC 5-PO1-HD-13021 (NICHD)
SO JOURNAL OF PEDIATRICS, (1991 Jan) 118 (1) 34-8.
Journal code: JI.Z. ISSN: 0022-3476.
CY United States
DT Journal: Article; (JOURNAL ARTICLE)
LA English
FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals
EM 199104
AB Although antibodies to the lipopolysaccharide antigens of Shigella have been demonstrated in human milk, such antibodies do not explain the putative protective effect of breast-feeding against symptomatic Shigella infection. Shigella species do not share related lipopolysaccharides, but they do possess closely related virulence plasmids that code for the proteins essential for cell invasion. We therefore sought to determine the frequency, amount, and duration of excretion of human

milk
antibodies to these shared virulence ***plasmid***-associated antigens in populations of different rates of Shigella infection frequency (Mexico City, high; Houston, low). Such antibodies were present in the milk of virtually all the Mexican women but also were present in a large proportion of milk samples from the women living in Houston. The of these antibodies were highest in colostrum but after 2 weeks of lactation fell to stable levels. The frequency and persistence of these antibodies in the milk of the women from Houston suggest that the memory and drive for secretion of these antibodies is extremely long lived.

=> s 15
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L22 5 L5
=> dup rem l22
PROCESSING COMPLETED FOR L22
L23 5 DUP REM L22 (0 DUPLICATES REMOVED)
=> d 1-bib ab
YOU HAVE REQUESTED DATA FROM 5 ANSWERS -
CONTINUE? Y(N)?

L23 ANSWER 1 OF 5 BIOSIS COPYRIGHT 1999 BIOSIS
AN 1995242446 BIOSIS
DN PREV199598256746
TI ***Recombinant*** ***immunoglobulin*** A expressed in a mouse
mammary gland cell line.
AU Rindisbacher, L. (1); Berdoz, J.; Jeanguenat, N.; Cortesby, B. (1); Kraehenbuehl, J.-P.
CS (1) Inst. Biol. Animale, Univ. Lausanne, Lausanne Switzerland
SO Experientia (Basel), (1995) Vol. 51, No. ABSTR., pp. A77.
Meeting Info.: 27th Annual Meeting of the Swiss Societies for Experimental Biology (USGEB/USSBE) Fribourg, Switzerland March 30-31, 1995
ISSN: 0014-4754.
DT Conference
LA English

L23 ANSWER 2 OF 5 CAPLUS COPYRIGHT 1999 ACS

AN 1991-88772 CAPLUS
DN 114-88772
TI Recombinant ricin A chain-monoclonal antibody conjugates for cancer targeting therapy
IN Frankel, Arthur E.
PA Cetus Corp., USA
SO U.S., 16 pp. Cont.-in-part of U.S. Ser. No. 806,256, abandoned.
CODEN: USXXAM
DT Patent
LA English
FAN CNT 2
PATENT NO. KIND DATE APPLICATION NO.
DATE
PI US 4962188 A 19901009 US 1986-913357 19860930
CA 1287578 A1 19910813 CA 1986-524645 19861205
JP 62209098 A2 19870914 JP 1986-289791 19861206
US 4956453 A 19900911 US 1987-69720 19870706
PRA1 US 1985-806256 19851206
AB The title conjugates comprising unglycosylated recombinant ricin A chain and monoclonal antibodies (MAbs) to e.g. ovarian cancer are manifd. and characterized. The conjugates are maintained at an effective cytotoxic amt. in the circulation of a host animal for substantially longer times than similar conjugates contg. native ricin A chain from, e.g., castor beans. The conjugates of the invention are cleared approx. 9 times slower from the circulation of a host animal than conjugates prepd. from native ricin A chain. Prodn. of the MAbs also is described.

L23 ANSWER 3 OF 5 CAPLUS COPYRIGHT 1999 ACS
AN 1991-40618 CAPLUS
DN 114-40618
TI Vaccination against tumor cells expressing breast cancer epithelial tumor antigen
antigen
AU Hareuveni, Maria; Gautier, Claudie; Kteny, Marie Paule; Wreschner, Daniel; Chabbon, Pierre; Lathe, Richard
CS Lab. Genet. Mol. Eucaryotes, Inst. Chim. Biol., Strasbourg, 67085, Fr.
SO Proc. Natl. Acad. Sci. U. S. A. (1990), 87(23), 9498-502
CODEN: PNASA6; ISSN: 0027-8424
DT Journal
LA English
AB Ninety-one percent of breast tumors aberrantly express an epithelial tumor antigen (ETA) identified by monoclonal antibody H23. Vaccinia virus recombinants expressing tumor antigens have considerable promise in the active immunotherapy of cancer, and the authors have evaluated the

potential of vaccinia recombinants expressing the secreted (S) and cell-assoc. (transmembrane, T) forms of H23 ETA to elicit immunity to tumor cells expressing ETA. Tumorogenic ras-transformed Fischer rat fibroblast lines FR-S and FR-T, expressing the S or T form of H23 ETA, resp., were constructed for use in challenge expts. Expression of H23 ETA in these lines was confirmed by Western blotting and immunofluorescence. When challenged by s.c. seeding of tumor cells, 97% (FR-S) and 91% (FR-T) of syngeneic Fischer rats rapidly developed tumors that failed to regress. Vaccination with recombinant vaccinia virus expressing ETA-T prior to challenge prevented tumor development in 82% of animals seeded with FR-T cells but in only 61% of animals seeded with FR-S. The vaccinia recombinant expressing the S form was a less effective immunogen, and vaccination protected only 29-30% of animals from developing tumors upon challenge with either FR-S or -T cells. The increased immunogenicity of the recombinant expressing ETA-T was reflected in elevated levels of ETA-reactive antibody in vaccinated animals, confirming that antigens expressed from vaccinia virus are less effective immunogens than their membrane-assoc. counterparts.

L23 ANSWER 4 OF 5 CAPLUS COPYRIGHT 1999 ACS
AN 1989:188523 CAPLUS
DN 110:188523
TI Characterization and biodistribution of recombinant and recombinant/chimeric constructs of monoclonal antibody B72.3
AU Colcher, David; Mileic, Diane; Roselli, Mario; Raubitschek, Andrew;
Yarranton, Geoffrey; King, David; Adair, John; Whittle, Nigel; Bodner, Mark; Schlom, Jeffrey
CS Radiat. Oncol. Branch, Natl. Cancer Inst., Bethesda, MD, 20892, USA
SO Cancer Res. (1989), 49(7), 1738-45
CODEN: CNREAB; ISSN: 0008-5472
DT Journal
LA English
AB B72.3 is a murine monoclonal antibody (IgG1) that recognizes a tumor-assoc. glycoprotein, termed TAG-72. B72.3 has been shown, using a variety of methodologies, to have a high degree of selective reactivity for colorectal, ovarian, lung, and breast carcinomas. Radiolabeled B72.3

has been administered both i.v. and i.p. in patients with colorectal ovarian cancer as well as other carcinomas and has been shown to selectively bind to approx. 70-80% of metastatic lesions. Greater than 50% of the patients that have been treated with B72.3 have developed an immunol. response to murine IgG after a single injection. In an attempt to minimize the immune response of these patients to the administered murine monoclonal antibody, a recombinant form of the murine B72.3 has been developed as well as a recombinant/chimeric antibody, using the variable regions of the murine B72.3 and human heavy chain (gamma.4) and light chain (kappa.) const. regions. It is reported here that both the recombinant B72.3 [rB72.3] and the recombinant/chimeric B72.3 [cB72.3(gamma.4)] IgGs maintain the tissue binding and idiotypic specificity of the native murine IgG. The native B72.3, rB72.3, and cB72.3(gamma.4) IgGs were radiolabeled and the biodistribution of these IgGs was studied in athymic mice bearing human colon carcinoma xenografts (LS-174T). Differences were obsd. between the cB72.3(gamma.4) and the native B72.3 in the percent of injected dose/g that localized in the tumor. The somewhat lower abs. amts. of the cB72.3(gamma.4) in the tumor are most likely due to the obsd. more rapid clearance from the blood and body of the mouse as compared to the native B72.3 and rB72.3. All 3 forms [native B72.3, rB72.3, and cB72.3(gamma.4)] of the IgG, however, able to localize the colon tumor with similar radiolocalization indexes [percent of injected dose/g in tumor divided by the percent of injected dose/g in normal tissue].

L23 ANSWER 5 OF 5 CAPLUS COPYRIGHT 1999 ACS
AN 1988:400354 CAPLUS
DN 109:354
TI Response of primary human mammary tumor cell cultures to a monoclonal antibody-recombinant ricin A chain immunotoxin
AU Bjorn, Michael J.; Smith, Helene S.; Dairkee, Shahnaz H.
CS Dep. Protein Chem., Cetus Corp., Emeryville, CA, 94608, USA
SO Cancer Immunol. Immunother. (1988), 26(2), 121-4
CODEN: CIIMDN; ISSN: 0340-7004
DT Journal
LA English
AB Malignant epithelial tumor cells were isolated and cultured from 10 human mammary specimens of cancerous origin. The 260F9 monoclonal antibody

(MAB) bound to frozen sections of all 10 tumors tested and to primary cultured cells from the tumors. Cultured cells from all 10 tumors were sensitive to the clonal inhibitory effects of immunotoxin 260F9 MAB-recombinant ricin A chain. At the immunotoxin concn. of 200 ng/mL (about 1 nM), the inhibition of colony formation was >99% for all 10 tumors.

=> d 2 kwic

L23 ANSWER 2 OF 5 CAPLUS COPYRIGHT 1999 ACS
IT ***Mammary*** gland (neoplasm, monoclonal) ***antibody*** to, ***recombinant*** ricin A chain conjugate with, for targeting therapy)

=> s 17

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L24 4 L7

=> dup rem l24

PROCESSING COMPLETED FOR L24
L25 4 DUP REM L24 (0 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 4 ANSWERS - CONTINUE? Y(N):

L25 ANSWER 1 OF 4 BIOSIS COPYRIGHT 1999 BIOSIS
AN 1995:242446 BIOSIS
DN PREV199598256746
TI ***Recombinant*** ***immunoglobulin*** A expressed in a mouse
mammary gland cell line.
AU Rindisbacher, L. (1); Berdoz, J.; Jeanguenat, N.; Corbhesy, B. (1); Kraehenbuehl, J.-P.
CS (1) Inst. Biol. Animale, Univ. Lausanne, Lausanne Switzerland
SO Experientia (Basel), (1995) Vol. 51, No. ABSTR., pp. A77.
Meeting Info.: 27th Annual Meeting of the Swiss Societies for Experimental Biology (USGEB/USSBE) Fribourg, Switzerland March 30-31, 1995
ISSN: 0014-4754.
DT Conference
LA English

L25 ANSWER 2 OF 4 CAPLUS COPYRIGHT 1999 ACS
AN 1991:88772 CAPLUS
DN 114:88772
TI Recombinant ricin A chain-monoclonal antibody conjugates for cancer targeting therapy
IN Frankel, Arthur E.
PA Cetus Corp., USA
SO U.S. 16 pp. Cont-in-part of U.S. Ser. No. 806,256, abandoned.
CODEN: USXXAM
DT Patent
LA English
FAN,CNT 2
PATENT NO. KIND DATE APPLICATION NO.
DATE

PI US 4962188 A 19901009 US 1986-913357 19860930
CA 1287578 A1 19910813 CA 1986-524645 19861205
JP 62209098 A2 19870914 JP 1986-289791 19861206
US 4956453 A 19900911 US 1987-69720 19870706
PRA1 US 1985-806256 19851206
AB The title conjugates comprising unglycosylated recombinant ricin A chain and monoclonal antibodies (MAbs) to e.g. ovarian cancer are manufactured and characterized. The conjugates are maintained at an effective cytotoxic amt. in the circulation of a host animal for substantially longer times than similar conjugates contg. native ricin A chain from, e.g., castor beans. The conjugates of the invention are cleared approx. 9 times slower from the circulation of a host animal than conjugates prepd. from native ricin A chain. Prodn. of the MAbs also is described.

L25 ANSWER 3 OF 4 CAPLUS COPYRIGHT 1999 ACS
AN 1989:188523 CAPLUS
DN 110:188523
TI Characterization and biodistribution of recombinant and recombinant/chimeric constructs of monoclonal antibody B72.3
AU Colcher, David; Milenic, Diane; Roselli, Mario; Raubitschek, Andrew;
Yarranton, Geoffrey; King, David; Adair, John; Whittle, Nigel; Bodner, Mark; Schlom, Jeffrey
CS Radiat. Oncol. Branch, Natl. Cancer Inst., Bethesda, MD, 20892, USA
SO Cancer Res. (1989), 49(7), 1738-45
CODEN: CNREA8; ISSN: 0008-5472
DT Journal
LA English
AB B72.3 is a murine monoclonal antibody (IgG1) that recognizes a tumor-assocd. glycoprotein, termed TAG-72. B72.3 has been shown, using a variety of methodologies, to have a high degree of selective reactivity for colorectal, ovarian, lung, and breast carcinomas. Radiolabeled B72.3 has been administered both i.v. and i.p. in patients with colorectal and ovarian cancer as well as other carcinomas and has been shown to selectively bind to approx. 70-80% of metastatic lesions. Greater than 50% of the patients that have been treated with B72.3 have developed an immunol. response to murine IgG after a single injection. In an attempt to minimize the immune response of these patients to the administered murine monoclonal antibody, a recombinant form of the murine B72.3 has been developed as well as a recombinant/chimeric antibody, using the variable regions of the murine B72.3 and human heavy chain (gamma.4) and light chain (kappa.) const. regions. It is reported here that both the recombinant B72.3 [rB72.3] and the recombinant/chimeric B72.3 [cB72.3(gamma.4)] iGGs maintain the tissue binding and idiotypic specificity of the native murine IgG. The native B72.3, rB72.3, and cB72.3(gamma.4) IgGs were radiolabeled and the biodistribution of these IgGs was studied in athymic mice bearing human colon carcinoma xenografts (LS-174T). Differences were obsd. between the cB72.3(gamma.4) and the native B72.3 in the percent of injected dose/g that localized in the tumor. The somewhat lower abs. amts. of the cB72.3(gamma.4) in the tumor are most likely due to the obsd. more rapid clearance from the blood and body of the mouse as compared to the native B72.3 and rB72.3. All 3 forms [native B72.3, rB72.3, and cB72.3(gamma.4)] of the IgG, however, localize the colon tumor with similar radiolocalization indexes [percent of injected dose/g in tumor divided by the percent of injected dose/g in normal tissue].

L25 ANSWER 4 OF 4 CAPLUS COPYRIGHT 1999 ACS
AN 1988:400354 CAPLUS
DN 109:354
TI Response of primary human mammary tumor cell cultures to a monoclonal antibody-recombinant ricin A chain immunotoxin
AU Bjorn, Michael J.; Smith, Helene S.; Darke, Shahnaz H.
CS Dep. Protein Chem., Cetus Corp., Emeryville, CA, 94608, USA
SO Cancer Immunol. Immunother. (1988), 26(2), 121-4
CODEN: CIIMDN; ISSN: 0340-7004
DT Journal
LA English
AB Malignant epithelial tumor cells were isolated and cultured from mammary specimens of cancerous origin. The 260F9 monoclonal antibody (MAB) bound to frozen sections of all 10 tumors tested and to primary cultured cells from the tumors. Cultured cells from all 10 tumors were sensitive to the clonal inhibitory effects of immunotoxin 260F9 MAB-recombinant ricin A chain. At the immunotoxin concn. of 200 ng/mL (about 1 nM), the inhibition of colony formation was >99% for all 10 tumors.
=> d 3 kwic
L25 ANSWER 3 OF 4 CAPLUS COPYRIGHT 1999 ACS
IT ***Mammary*** gland (neoplasm, carcinoma, radioiodinated monoclonal ***antibody*** and ***recombinant*** /chimeric constructs metab. by, scintigraphy in relation to)
=> s 19
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L26 0 L9
=> s 111
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L27 1 L11
=> d bib ab
L27 ANSWER 1 OF 1 CAPLUS COPYRIGHT 1999 ACS
AN 1989:188523 CAPLUS
DN 110:188523
TI Characterization and biodistribution of recombinant and recombinant/chimeric constructs of monoclonal antibody B72.3
AU Colcher, David; Milenic, Diane; Roselli, Mario; Raubitschek, Andrew;
Yarranton, Geoffrey; King, David; Adair, John; Whittle, Nigel; Bodner, Mark; Schlom, Jeffrey
CS Radiat. Oncol. Branch, Natl. Cancer Inst., Bethesda, MD, 20892, USA
SO Cancer Res. (1989), 49(7), 1738-45

CODEN: CNREA8; ISSN: 0008-5472
DT Journal
LA English
AB B72.3 is a murine monoclonal antibody (IgG1) that recognizes a tumor-associated glycoprotein, termed TAG-72. B72.3 has been shown, using a variety of methodologies, to have a high degree of selective reactivity for colorectal, ovarian, lung, and breast carcinomas. Radiolabeled B72.3 has been administered both i.v. and i.p. in patients with colorectal and ovarian cancer as well as other carcinomas and has been shown to selectively bind to approx. 70-80% of metastatic lesions. Greater than 50% of the patients that have been treated with B72.3 have developed an immunol. response to murine IgG after a single injection. In an attempt to minimize the immune response of these patients to the administered murine monoclonal antibody, a recombinant form of the murine B72.3 has been developed as well as a recombinant/chimeric antibody, using the variable regions of the murine B72.3 and human heavy chain (gamma.4) and light chain (kappa.) const. regions. It is reported here that both the recombinant B72.3 [rB72.3] and the recombinant/chimeric B72.3 [cB72.3(gamma.4)] IGGs maintain the tissue binding and idiotypic specificity of the native murine IgG. The native B72.3, rB72.3, and cB72.3(gamma.4) IGGs were radiolabeled and the biodistribution of these IGGs was studied in athymic mice bearing human colon carcinoma xenografts (LS-174T). Differences were obsd. between the cB72.3(gamma.4) and the native B72.3 in the percent of injected dose/g that localized in the tumor. The somewhat lower abs. ants. of the cB72.3(gamma.4) in the tumor are most likely due to the obsd. more rapid clearance from the blood and body of the mouse as compared to the native B72.3 and rB72.3. All 3 forms [native B72.3, rB72.3, and cB72.3(gamma.4)] of the IgG, however, able to localize the colon tumor with similar radiolocalization indexes [percent of injected dose/g in tumor divided by the percent of injected dose/g in normal tissue].
=> d kwic
L27 ANSWER 1 OF 1 CAPLUS COPYRIGHT 1999 ACS
IT ***Mammary*** gland (neoplasm, carcinoma, radioiodinated monoclonal

antibody
recombinant and ***recombinant*** /chimeric
constructs metab. by, scintigraphy in relation to)
=> s 111
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L28 1 L11
=> d
L28 ANSWER 1 OF 1 CAPLUS COPYRIGHT 1999 ACS
AN 1989:188523 CAPLUS
DN 110:188523
TI Characterization and biodistribution of recombinant and recombinant/chimeric constructs of monoclonal antibody B72.3
AU Colcher, David; Milenic, Diane; Roselli, Mario; Raubitschek, Andrew;
Yarranton, Geoffrey; King, David; Adair, John; Whittle, Nigel; Bodmer, Mark; Schlom, Jeffrey
CS Radiat. Oncol. Branch, Natl. Cancer Inst., Bethesda, MD, 20892, USA
SO Cancer Res. (1989), 49(7), 1738-45
CODEN: CNREA8; ISSN: 0008-5472
DT Journal
LA English
=> s 112
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
3 FILES SEARCHED...
'AB' IS NOT A VALID FIELD CODE
L29 9 L12
=> dup rem 129
PROCESSING COMPLETED FOR L29
L30 4 DUP REM L29 (5 DUPLICATES REMOVED)
=> d 1- bib ab
YOU HAVE REQUESTED DATA FROM 4 ANSWERS -
CONTINUE? Y(N)?
L30 ANSWER 1 OF 4 BIOSIS COPYRIGHT 1999 BIOSIS
AN 1998:409805 BIOSIS
DN PREV199800409805
TI Cloning, expression, and characterization of recombinant Fab antibodies

against dioxin.
AU Lee, Nanju (1); Holtzapple, Carol K.; Stanker, Larry H.
CS (1) Food Anim. Protection Res. Lab., Agricultural Res. Service, U.S. Dep. Agriculture, 2881 F and B Road, College Station, TX 77845-9594 USA
SO Journal of Agricultural and Food Chemistry, (Aug., 1998) Vol. 46, No. 8, pp. 3381-3388.
ISSN: 0021-8561.
DT Article
LA English
AB Using two hybridoma cell lines (DD1 and DD3) secreting anti-dioxin monoclonal antibodies as a source for messenger RNA and cDNA, light and heavy chain gene fragments of Fab domains were amplified by the polymerase chain reaction (PCR). The amplified gene fragments were cloned into the pFabUSDA1 ***vector*** for expression of ***recombinant*** Fab ***antibodies*** in Escherichia coli. Expression of the soluble and ***functional*** recombinant Fab antibodies (designated rFab1-1 and rFab3-3) was confirmed by an indirect immunoassay using dioxin conjugated to rabbit serum albumin. On the basis of these rFabs, two competitive inhibition immunoassays using 2,3,7,8-tetrachlorodibenzo-p-dioxin (2,3,7,8-TCDD) as a competitor were developed. The concentration of 2,3,7,8-TCDD required to inhibit color development by 50% (IC50) determined from the dose response curves for rFab1-1 and rFab3-3 were 10.4 +/- 2.4 and 12.2 +/- 6.0 ng/mL, respectively. The binding properties of both rFab antibodies for other chemically related compounds were relatively similar to those of their respective monoclonal antibodies and enzymatically derived Fab fragments.
L30 ANSWER 2 OF 4 MEDLINE DUPLICATE 1
AN 1998455662 MEDLINE
DN 98455662
TI Isolation and recombinant expression of an MHV-JHM neutralising monoclonal antibody.
AU Kolb A F; Lechnermaier M; Heister A; Toksoy A; Siddell S G
CS Institute of Virology and Immunology, University of Wurzburg, Germany.
SO ADVANCES IN EXPERIMENTAL MEDICINE AND BIOLOGY, (1998) 440 657-64.
Journal code: 2LU. ISSN: 0065-2598.
CY United States

DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199903
EW 19990303
AB The monoclonal antibody A1 (mab A1) efficiently neutralises the infection of susceptible cells by the murine hepatitis virus MHV-JHM in vitro and in vivo (Wege et al., 1984). The variable regions of mab A1 were amplified from mRNA of the respective hybridoma cell line by RT-PCR and integrated into different eukaryotic expression vectors. The biological ***function*** of the ***recombinant*** ***antibody*** ***constructs*** was verified by virus neutralisation assays. Whereas a complete recombinant antibody (mab A1 rec.) expressed in transfected murine myeloma cells inhibited the MHV-JHM infection as well as the parental antibody, a single-chain Fv derived from mab A1 did not show any neutralising activity.

L30 ANSWER 3 OF 4 MEDLINE DUPLICATE 2
AN 97041563 MEDLINE
DN 97041563
TI Lung cancer-reacting human recombinant antibody AE6F4: potential usefulness in the sputum cytodiagnosis.
AU Shoji M; Kawamoto S; Seki K; Teruya K; Setoguchi Y; Mochizuki K; Kato M; Hashizume S; Hanagiri T; Yoshimatsu T; Nakamishi K; Yasumoto K; Nagashima A; Nakahashi H; Suzuki T; Imai T; Shirahata S; Nomoto K; Murakami H
CS Morinaga Institute of Biological Science, Yokohama, Japan.
SO HUMAN ANTIBODIES AND HYBRIDOMAS, (1996) 7 (1) 27-36.
Journal code: A6A. ISSN: 0956-960X.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199706
EW 19970601
AB Human monoclonal antibody (hMAb) AE6F4 has been shown to be potentially useful for immunocytological detection of lung cancer cells in sputum. By recombinant DNA technology, IgM type hMAb AE6F4 was switched to IgG. The IgG mimic ***recombinant*** AE6F4 ***antibody*** expression ***plasmid*** was ***assembled*** using the ***antibody*** heavy chain gene, which ligated the gene encoding VH and CHI(mu)

domains of hMAb AE6F4 heavy chain to the gene encoding CH2(gamma 1) and CH3(gamma 1) domains of human IgG heavy chain, and the antibody light chain gene of hMAb AE6F4. The recombinant antibody expressed by baby hamster kidney (BHK)-21 cells showed molecular size equivalence to IgG, and human mu-gamma hybrid heavy and kappa light chains. The immunological specificity of the recombinant antibody was the same as that of hMAb AE6F4 by immunoblotting analysis to the 14-3-3 protein, the putative antigen of hMAb AE6F4, and by immunohistochemical and immunocytological analyses using tissue sections and sputa of lung cancer patients. The BHK-21 cells produced the recombinant antibody persistently and the productivity was greater than 20 times that by human-human hybridoma producing hMAb AE6F4.

L30 ANSWER 4 OF 4 CAPLUS COPYRIGHT 1999 ACS
AN 1992:631697 CAPLUS
DN 117:231697
TI Cloning, bacterial expression and crystallization of Fv antibody fragments
AU Eisele, Jean Luc; Boulot, Ginette; Chitarra, Veronique; Riottot, Marie
A.: Bhat, T
Narayana; Spinelli, Silvia; Poljak, Roberto J.
CS Dep. Immunol., Inst. Pasteur, Paris, F-75724, Fr.
SO J. Cryst. Growth (1992), 122(1-4), 337-43
CODEN: JCRGAE; ISSN: 0022-0248
DT Journal; General Review
LA English
AB A review, with 16 refs., of the author's work on the variable Fv fragments of ***antibodies***, cloned in ***recombinant*** ***plasmids***, and expressed in bacteria as ***functional*** proteins having immunochem. properties similar or identical with those of the corresponding parts of the parent eukaryotic antibodies.

=> s113

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L31 0 L13
=> s115

domains of hMAb AE6F4 heavy chain to the gene encoding CH2(gamma 1) and CH3(gamma 1) domains of human IgG heavy chain, and the antibody light chain gene of hMAb AE6F4. The recombinant antibody expressed by baby hamster kidney (BHK)-21 cells showed molecular size equivalence to IgG, and human mu-gamma hybrid heavy and kappa light chains. The immunological specificity of the recombinant antibody was the same as that of hMAb AE6F4 by immunoblotting analysis to the 14-3-3 protein, the putative antigen of hMAb AE6F4, and by immunohistochemical and immunocytological analyses using tissue sections and sputa of lung cancer patients. The BHK-21 cells produced the recombinant antibody persistently and the productivity was greater than 20 times that by human-human hybridoma producing hMAb AE6F4.

L30 ANSWER 4 OF 4 CAPLUS COPYRIGHT 1999 ACS
AN 1992:631697 CAPLUS
DN 117:231697
TI Cloning, bacterial expression and crystallization of Fv antibody fragments
AU Eisele, Jean Luc; Boulot, Ginette; Chitarra, Veronique; Riottot, Marie
A.: Bhat, T
Narayana; Spinelli, Silvia; Poljak, Roberto J.
CS Dep. Immunol., Inst. Pasteur, Paris, F-75724, Fr.
SO J. Cryst. Growth (1992), 122(1-4), 337-43
CODEN: JCRGAE; ISSN: 0022-0248
DT Journal; General Review
LA English
AB A review, with 16 refs., of the author's work on the variable Fv fragments of ***antibodies***, cloned in ***recombinant*** ***plasmids***, and expressed in bacteria as ***functional*** proteins having immunochem. properties similar or identical with those of the corresponding parts of the parent eukaryotic antibodies.

=> s113

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L31 0 L13
=> s115

'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
'AB' IS NOT A VALID FIELD CODE
L32 0 L15
=> d his

(FILE 'HOME' ENTERED AT 10:20:31 ON 10 AUG 1999)

FILE 'MEDLINE' ENTERED AT 10:20:36 ON 10 AUG 1999
L1 58323 S IMMUNOGLOB? OR ANTIBOD?/AB,BI
L2 843 S LI(5A)(CONSTRUCT# OR PLASMID# OR VECTOR#)/AB,BI
L3 2 S L2(5A)(MILK)/AB,BI
L4 357 S LI(5A)(MAMMARY)
L5 0 S L4(10A)(RECOMBINANT)
L6 357 S LI(5A)(MAMMARY)/AB,BI
L7 0 S L6(5A)(RECOMBINANT)/AB,BI
L8 33 S L6(5A)(SPECIFIC)/AB,BI
L9 0 S L8 AND PROMOTER#/AB,BI
L10 57 S L2(5A)(RECOMBIN?)/AB,BI
L11 0 S LI(05A)(MILK OR MAMMARY)/AB,BI
L12 2 S LI(10A)(FUNCTION? OR ASSEMBL?)/AB,BI
L13 0 S L2(5A)(MAMMARY)(W)(TUMOR OR TUMOUR)/AB,BI
L14 74 S LI(5A)(MAMMARY)(W)(TUMOR OR TUMOUR)/AB,BI
L15 0 S LI(45A)(PROMOTER#)/AB,BI
L16 42 S LI(45A)(VIRUS)/AB,BI
L17 2 S L16 AND MILK/AB,BI

FILE 'MEDLINE, EMBASE, BIOSIS, INPADOC, CAPLUS' ENTERED AT 10:28:48 ON 10 AUG 1999
L18 13 S LI7
L19 10 DUP REM L18 (3 DUPLICATES REMOVED)
L20 11 S L3
L21 7 DUP REM L20 (4 DUPLICATES REMOVED)
L22 5 S L5
L23 5 DUP REM L22 (0 DUPLICATES REMOVED)
L24 4 S L7
L25 4 DUP REM L24 (0 DUPLICATES REMOVED)
L26 0 S L9
L27 1 S LI1
L28 1 S LI1
L29 9 S LI2
L30 4 DUP REM L29 (5 DUPLICATES REMOVED)
L31 0 S LI3
L32 0 S LI5
=>

---Logging off of STN---

*

 FILE 'USPAT' ENTERED AT 09:28:32 ON 10 AUG 1999

 * U.S. PATENT TEXT FILE *
 *
 * THE WEEKLY PATENT TEXT AND IMAGE DATA IS
 CURRENT *
 * THROUGH AUGUST 10,1999 *
 *

 => s mammary tumor
 4414 MAMMARY
 22585 TUMOR
 L1 1307 MAMMARY TUMOR
 (MAMMARY(W)TUMOR)
 => s l1(5a)(promoter)
 27962 PROMOTER
 L2 202 L1(5A)(PROMOTER)
 => s l2(10a)(antibod? or immunoglobulin#)
 36150 ANTIBOD?
 11405 IMMUNOGLOBULIN#
 L3 1 L2(10A)(ANTIBOD? OR IMMUNOGLOBULIN#)
 => d
 1. 5,733,768, Mar. 31, 1998, Amyloid precursor protein protease; Eric
 P.
 Dixon, et al., 435/226 [IMAGE AVAILABLE]
 => s mammary(5a)(promoter#)
 4414 MAMMARY
 36145 PROMOTER#
 L4 299 MAMMARY(5A)(PROMOTER#)
 => s l4(10a)(antibod? or immunoglob? or chain#)
 36150 ANTIBOD?
 11609 IMMUNOGLOB?
 343081 CHAIN#
 L5 3 L4(10A)(ANTIBOD? OR IMMUNOGLOB? OR
 CHAIN#)
 => d l- cit ab
 1. 5,919,650, Jul. 6, 1999, Method for inactivation of protein function;

Mariano Barbacid, et al., 435/69.1, 320.1, 330 [IMAGE
 AVAILABLE]
 US PAT NO: 5,919,650 [IMAGE AVAILABLE] L5: 1 of 3
 ABSTRACT:
 Method for inactivating the function produced by a protein using an
 intracellularly expressed antibody or fragment thereof.
 2. 5,880,327, Mar. 9, 1999, Transgenic mammals expressing human
 coagulation factor VIII; Henryk Lubon, et al., 800/7; 435/455; 800/4,
 14,
 15, 16, 17, 18, 21, 24, 25 [IMAGE AVAILABLE]
 US PAT NO: 5,880,327 [IMAGE AVAILABLE] L5: 2 of 3
 ABSTRACT:
 A non-human transgenic mammalian animal, as described above,
 contains an
 exogenous double stranded DNA sequence stably integrated into the
 genome
 of the animal, which comprises cis-acting regulatory units operably
 linked to a DNA sequence encoding human Factor VIII protein and a
 signal
 peptide, where the cis-acting regulatory units are active in mammary
 gland cells and the signal peptide is active in directing newly
 expressed
 Factor VIII into the milk of the animal. The promoter may be a milk
 protein promoter such as for whey acidic protein, casein, lactalbumin,
 or
 beta-lactoglobulin promoter. The transgenic mammals are preferably
 farm
 animals, for example, cows, goats, sheep, rabbits and pigs. Concurrent
 expression of a gene for human von Willebrand's Factor into milk may
 be
 used to stabilize newly-secreted Factor VIII.
 3. 5,733,768, Mar. 31, 1998, Amyloid precursor protein protease; Eric
 P.
 Dixon, et al., 435/226 [IMAGE AVAILABLE]
 US PAT NO: 5,733,768 [IMAGE AVAILABLE] L5: 3 of 3
 ABSTRACT:
 This invention provides an APP-cleaving protein and related nucleic
 acid
 compounds. The invention also provides methods, materials and
 assays. The
 compounds of this invention will further the characterization of
 neurological diseases such as Alzheimer's disease and Down's
 syndrome.
 => s mouse mammary
 46615 MOUSE
 4414 MAMMARY
 L6 888 MOUSE MAMMARY
 (MOUSE(W)MAMMARY)
 => s l6(5a)(promoter#)
 36145 PROMOTER#
 L7 210 L6(5A)(PROMOTER#)
 => s l7(5a)(immunoglob? or antibod? or chain#)
 11609 IMMUNOGLOB?
 36150 ANTIBOD?
 343081 CHAIN#
 L8 1 L7(5A)(IMMUNOGLOB? OR ANTIBOD? OR CHAIN#)
 => d
 1. 5,733,768, Mar. 31, 1998, Amyloid precursor protein protease; Eric
 P.
 Dixon, et al., 435/226 [IMAGE AVAILABLE]
 => d kwic
 US PAT NO: 5,733,768 [IMAGE AVAILABLE] L8: 1 of 1
 SUMMARY:
 BSUM(64)
 The . . . Promoters which may be used, for example, are the
 thymidine
 kinase promoter, the metalloionin promoter, the heat shock
 promoter,
 the **mouse**mammary** tumor virus **promoter** or various
 other vital
 and **immunoglobulin** promoters.
 => s tumor virus promoter#
 22585 TUMOR
 24029 VIRUS
 36145 PROMOTER#
 L9 74 TUMOR VIRUS PROMOTER#
 (TUMOR(W)VIRUS(W)PROMOTER#)
 => s l9(5a)(mammary)
 4414 MAMMARY
 L10 74 L9(5A)(MAMMARY)
 => s l10(10a)(immunoglob? or antibod? or chain#)
 11609 IMMUNOGLOB?
 36150 ANTIBOD?
 343081 CHAIN#
 L11 1 L10(10A)(IMMUNOGLOB? OR ANTIBOD? OR

CHAIN#)

=> d

1. 5,733,768, Mar. 31, 1998, Amyloid precursor protein protease; Eric P. Dixon, et al., 435/226 [IMAGE AVAILABLE]

=> file epaab

FILE 'EPOABS' ENTERED AT 09:36:47 ON 10 AUG 1999

* EUROPEAN PATENT ABSTRACTS *

=> s 111

2533 TUMOR

3729 VIRUS

3960 PROMOTER#

0 TUMOR VIRUS PROMOTER#

(TUMOR(W)VIRUS(W)PROMOTER#)

254 MAMMARY

894 IMMUNOGLOB?

9129 ANTIBOD?

39372 CHAIN#

L12 0 L10(10A)(IMMUNOGLOB? OR ANTIBOD? OR CHAIN#)

=> s (metalloth?)(3a)(promoter#)

67 METALLOTH?

3960 PROMOTER#

L13 13 (METALLOTH?)(3A)(PROMOTER#)

=> s 113(10a)(mammary)

254 MAMMARY

L14 0 L13(10A)(MAMMARY)

=> file uspat

FILE 'USPAT' ENTERED AT 09:37:55 ON 10 AUG 1999

* U.S. PATENT TEXT FILE *

*

* THE WEEKLY PATENT TEXT AND IMAGE DATA IS CURRENT *

* THROUGH AUGUST 10,1999 *

*

=> s 114

2276 METALLOTH?

36145 PROMOTER#

4414 MAMMARY

L15 52 L13(10A)(MAMMARY)

=> s 115(10a)(immunoglob? or antibod? or chain#)

11609 IMMUNOGLOB?

36150 ANTIBOD?

343081 CHAIN#

L16 1 L15(10A)(IMMUNOGLOB? OR ANTIBOD? OR CHAIN#)

=> d

1. 5,733,768, Mar. 31, 1998, Amyloid precursor protein protease; Eric P. Dixon, et al., 435/226 [IMAGE AVAILABLE]

=> d kwic

US PAT NO: 5,733,768 [IMAGE AVAILABLE] L16: 1 of 1

SUMMARY:

BSUM(64)

The . . . necessary, the appropriate regulatory elements using well known techniques. Promoters which may be used, for example, are the thymidine kinase **promoter**, the **metallothionin** **promoter**, the heat shock promoter, the mouse **mammary** tumor virus promoter or various other vital and **immunoglobulin** promoters.

=> s immunoglob? or antibod?

11609 IMMUNOGLOB?

36150 ANTIBOD?

L17 37141 IMMUNOGLOB? OR ANTIBOD?

=> s 117(5)(linked)

MISSING OPERATOR 'L17/5'

=> s 117(5a)(linked)

149754 LINKED

L18 3157 L17(5A)(LINKED)

=> s 118(5a)(promoter#)

36145 PROMOTER#

L19 16 L18(5A)(PROMOTER#)

=> s 119(5a)(mammary)

4414 MAMMARY

L20 0 L19(5A)(MAMMARY)

=> d 119 1- cit ab

1. 5,922,545, Jul. 13, 1999, In vitro peptide and antibody display libraries; Larry C. Mattheakis, et al., 435/6, 5, 7, 1; 436/518 [IMAGE AVAILABLE]

US PAT NO: 5,922,545 [IMAGE AVAILABLE] L19: 1 of 16

ABSTRACT:

Improved methods and novel compositions for identifying peptides and single-chain antibodies that bind to predetermined receptors or epitopes. Such peptides and antibodies are identified by improved and novel methods for affinity screening of polysomes displaying nascent peptides.

2. 5,919,452, Jul. 6, 1999, Methods of treating TNF.alpha.-mediated disease using chimeric anti-TNF antibodies; Junning Le, et al., 424/133.1, 145.1, 158.1, 530/387.3, 388.23, 389.2 [IMAGE AVAILABLE]

US PAT NO: 5,919,452 [IMAGE AVAILABLE] L19: 2 of 16

ABSTRACT:

Treatment of tumor necrosis factor, TNF, mediated pathologies is provided by administering anti-TNF compounds, such as anti-TNF antibodies and anti-TNF peptides, which compounds are specific for tumor necrosis factor-alpha. (TNF.alpha.) or tumor necrosis factor-beta. (TNF.beta.) and which are useful for in vivo therapy or diagnosis of TNF.alpha.-mediated pathologies and conditions, wherein the anti-TNF compound is selected from the group consisting of at least one of an immunoglobulin variable region, a fragment of a TNF receptor and an anti-TNF peptide, such as a structural analog of a anti-TNF antibody fragment or a TNF receptor fragment.

3. 5,871,901, Feb. 16, 1999, Assay for inhibitors of DP-1 and other DP proteins; Nicholas Berrie La Thangue, 435/4, 15, 21, 29, 194, 375; 530/358, 388.24, 389.2 [IMAGE AVAILABLE]

US PAT NO: 5,871,901 [IMAGE AVAILABLE] L19: 3 of 16

<p>ABSTRACT:</p> <p>The protein DP-1, part of the DP-1/E2F-1 transcription factor complex, as well as DP-2 and DP-3 has its phosphorylation level regulated during cell cycle progression. This finding allows assays to be based on changes in phosphorylation of DP proteins, in particular for agents which may affect the phosphorylation state of DP. DP-1 has been found to have a greater affinity to DNA when in a hypophosphorylated state.</p> <p>Antibodies that recognize phosphorylation sites on DP-1 are also disclosed.</p> <p>4. 5,851,829, Dec. 22, 1998, Method of intracellular binding of target molecules; Wayne A. Marasco, et al., 435/328; 424/577, 578, 435/325, 330, 333, 339, 339.1, 366, 372, 419 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,851,829 [IMAGE AVAILABLE] L19: 4 of 16</p> <p>ABSTRACT:</p> <p>The present invention relates to a method by which one can target an undesired target molecule or target antigen, preferably a protein. The method comprises the intracellular expression of an antibody capable of binding to the target. A DNA sequence is delivered to a cell, the DNA sequence contains a sufficient number of nucleotides coding for the portion of an antibody capable of binding to the target operably linked to a promoter that will permit expression of the antibody in the cell(s) of interest. The antibody is then expressed intracellularly and binds to the target, thereby disrupting the target from its normal actions.</p> <p>5. 5,849,992, Dec. 15, 1998, Transgenic production of antibodies in milk; Harry Meade, et al., 800/14, 7, 15, 16, 17, 18 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,849,992 [IMAGE AVAILABLE] L19: 5 of 16</p> <p>ABSTRACT:</p> <p>A method for the production of monoclonal antibodies in mammals milk through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.</p> <p>6. 5,834,597, Nov. 10, 1998, Mutated nonactivating IgG2 domains and anti CD3 antibodies incorporating the same; J. Yun Tso, et al., 530/387.3 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,834,597 [IMAGE AVAILABLE] L19: 6 of 16</p>	<p>ABSTRACT:</p> <p>The invention provides mutated IgG2 constant regions and anti-CD3 antibodies incorporating the same. Such antibodies specifically bind to the CD3 antigen on T-cells but induce reduced mitogenic response compared with otherwise identical antibodies bearing natural IgG2 constant regions. The antibodies can be used for treating disorders requiring immune suppression with fewer side effects than result from treatment with prior anti-CD3 antibodies.</p> <p>7. 5,827,690, Oct. 27, 1998, Transgenic production of antibodies in milk; Harry Meade, et al., 800/7; 530/867 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,827,690 [IMAGE AVAILABLE] L19: 7 of 16</p> <p>ABSTRACT:</p> <p>A method for the production of monoclonal antibodies in mammals milk, through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.</p> <p>8. 5,800,815, Sep. 1, 1998, Antibodies to P-selectin and their uses; Robert W. Chestnut, et al., 424/153.1, 133.1, 143.1, 173.1; 435/7.24, 70.21, 326, 328, 343, 346; 530/387.3, 388.2, 388.7, 389.6; 536/23.53 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,800,815 [IMAGE AVAILABLE] L19: 8 of 16</p> <p>ABSTRACT:</p> <p>The present invention relates to compositions and methods for treating inflammation and other pathological conditions using novel blocking P-selectin antibodies that inhibit adhesion of leukocytes to activated platelets and/or to activated vascular endothelium in vivo. Both murine and humanized antibodies are provided.</p> <p>9. 5,753,225, May 19, 1998, Antibodies that mimic actions of neurotrophins; Douglas O. Clary, et al., 424/130.1, 141.1, 143.1, 156.1; 530/387.1, 388.1, 388.22 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,753,225 [IMAGE AVAILABLE] L19: 9 of 16</p> <p>ABSTRACT:</p> <p>The use and production of immunoglobulins which activate trk receptors and imitate effects of neurotrophins are provided. Immunoglobulins which block trk receptor activation and methods of use are also provided.</p> <p>10. 5,714,352, Feb. 3, 1998, Directed switch-mediated DNA recombination; Aya Jakobovits, 435/462, 320.1, 328, 372.3 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,714,352 [IMAGE AVAILABLE] L19: 10 of 16</p>	<p>ABSTRACT:</p> <p>Switch regions derived from an immunoglobulin (Ig) gene are used to direct recombination between a targeting construct containing a promoter, a switch region (S sub.1), and 2) a target locus minimally containing a promoter, a switch region (S sub.2), and a target sequence.</p> <p>11. 5,698,195, Dec. 16, 1997, Methods of treating rheumatoid arthritis using chimeric anti-TNF antibodies; Junming Le, et al., 424/133.1, 141.1, 142.1, 145.1; 514/825; 530/351, 387.3, 388.1, 388.23 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,698,195 [IMAGE AVAILABLE] L19: 11 of 16</p> <p>ABSTRACT:</p> <p>Anti-TNF antibodies, fragments and regions thereof which are specific for human tumor necrosis factor-α (TNF-α) and are useful in vivo for diagnosis and therapy of a number of TNF-α-mediated pathologies and conditions, including rheumatoid arthritis as well as polynucleotides coding for murine and chimeric antibodies, methods of producing the antibody, methods of use of the anti-TNF antibody, or fragment, region or derivative thereof, in immunoassays and immunotherapeutic approaches are provided.</p> <p>12. 5,656,272, Aug. 12, 1997, Methods of treating TNF-α-mediated Crohn's disease using chimeric anti-TNF antibodies; Junming Le, et al., 424/133.1, 139.1, 145.1, 435/69.1, 69.6, 69.7, 530/387.3, 388.23 [IMAGE AVAILABLE]</p> <p>US PAT NO: 5,656,272 [IMAGE AVAILABLE] L19: 12 of 16</p> <p>ABSTRACT:</p> <p>Anti-TNF antibodies, fragments and regions thereof which are specific for human tumor necrosis factor-α (TNF-α) and are useful in vivo for diagnosis and therapy of a number of TNF-α-mediated pathologies and conditions, including Crohn's disease, as well as polynucleotides coding for murine and chimeric antibodies, methods of producing the antibody, methods of use of the anti-TNF antibody, or fragment, region or derivative thereof, in immunoassays and immunotherapeutic</p>
--	---	--

approaches are provided.

L3. 5,635,603, Jun. 3, 1997, Preparation and use of immunoconjugates; Hans J. Hansen, et al., 530/391.5; 424/172.1; 435/69.6 [IMAGE AVAILABLE]

US PAT NO: 5,635,603 [IMAGE AVAILABLE] L19: 13 of 16

ABSTRACT:

The present invention relates to immunoconjugates comprising an antibody fragment which is covalently bound to a diagnostic or therapeutic principle through a carbohydrate moiety in the light chain variable region of the antibody fragment. The invention also relates to immunoconjugates comprising an antibody moiety that is an intact antibody containing a glycosylation site in the light chain variable domain which has been introduced into the antibody by mutating the nucleotide sequence encoding the light chain. The resultant immunoconjugates retain the immunoreactivity of the antibody fragment or intact antibody, and target the diagnostic or therapeutic principle to a target tissue where the diagnostic or therapeutic effect is realized. Thus, the invention contemplates the use of such immunoconjugates for diagnosis and immunotherapy. The invention further relates to methods for preparing such immunoconjugates.

14. 5,529,774, Jun. 25, 1996, In vivo transfer of the HSV-TK gene implanted retroviral producer cells; David Barba, et al., 424/93.21, 93.2, 93.6; 514/44 [IMAGE AVAILABLE]

US PAT NO: 5,529,774 [IMAGE AVAILABLE] L19: 14 of 16

ABSTRACT:

The present invention is directed to methods of transferring genes to brain tumor cells in order to kill the cells. In general, the method of the present invention comprises: (1) introducing a retrovirus containing a selectable marker and at least one gene required for its replication into producer cells such that integration of the proviral DNA corresponding to the retrovirus into the genome of the producer cell results in the generation of a modified retrovirus wherein at least one of the genes required for replication of the retrovirus is replaced by the therapeutic gene or genes; (2) selecting producer cells in which the modified retrovirus is incorporated as part of the genome of the producer cells; (3) grafting the producer cells in proximity to the dividing tumor cell in order to infect the tumor cell with the modified retrovirus, thereby transferring the therapeutic gene or genes to the tumor cell; and

(4) killing the cells by administering a substance that is metabolized by the therapeutic gene transferred to the tumor cells into a metabolite that kills the cells. Suitable retroviral vectors and methods for generating them, producer cells, and grafting methods are described.

15. 5,474,771, Dec. 12, 1995, Murine monoclonal antibody (5c8) recognizes a human glycoprotein on the surface of T-lymphocytes, compositions containing same; Seth Lederman, et al., 424/133.1, 130.1, 144.1, 153.1, 154.1; 435/70.21, 343.2; 530/388.7, 388.73, 388.75 [IMAGE AVAILABLE]

US PAT NO: 5,474,771 [IMAGE AVAILABLE] L19: 15 of 16

ABSTRACT:

This invention provides a monoclonal antibody which specifically recognizes and forms a complex with a protein located on the surface of activated T cells and thereby inhibits T cell activation of B cells. This invention also provides the monoclonal antibody 5c8 (ATCC Accession No. HB 10916).

This invention provides a human CD4^{sup} - T cell leukemia cell line designated DI.1 (ATCC Accession No. CRL 10915) capable of constitutively providing contact-dependent helper function to B cells. This invention also provides an isolated protein from the surface of activated T cells, wherein the protein is necessary for T cell activation of B cells. This invention further provides an isolated, soluble protein from the surface of activated T cells, wherein the protein is necessary for T cell activation of B cells.

16. 5,443,953, Aug. 22, 1995, Preparation and use of immunoconjugates; Hans J. Hansen, et al., 424/1.49, 1.53, 9.341, 178.1, 179.1, 180.1, 181.1, 182.1, 183.1; 435/7.1, 7.2, 7.23, 69.6; 530/387.3, 391.3, 391.5, 391.7, 391.9 [IMAGE AVAILABLE]

US PAT NO: 5,443,953 [IMAGE AVAILABLE] L19: 16 of 16

ABSTRACT:

The present invention relates to immunoconjugates comprising an antibody fragment which is covalently bound to a diagnostic or therapeutic principle through a carbohydrate moiety in the light chain variable region of the antibody fragment. The invention also relates to immunoconjugates comprising an antibody moiety that is an intact antibody containing a glycosylation site in the light chain variable domain which has been introduced into the antibody by mutating the nucleotide sequence encoding the light chain. The resultant immunoconjugates retain the immunoreactivity of the antibody fragment or intact antibody, and target

the diagnostic or therapeutic principle to a target tissue where the diagnostic or therapeutic effect is realized. Thus, the invention contemplates the use of such immunoconjugates for diagnosis and immunotherapy. The invention further relates to methods for preparing such immunoconjugates.

=> d his

(FILE 'USPAT' ENTERED AT 09:28:32 ON 10 AUG 1999)

L1 1307 S MAMMARY TUMOR
L2 202 S L1(5A)(PROMOTER)
L3 1 S L2(10A)(ANTIBOD? OR IMMUNOGLOBULIN#)
L4 299 S MAMMARY(5A)(PROMOTER#)
L5 3 S L4(10A)(ANTIBOD? OR IMMUNOGLOB? OR CHAIN#)
L6 888 S MOUSE MAMMARY
L7 210 S L6(5A)(PROMOTER#)
L8 1 S L7(5A)(IMMUNOGLOB? OR ANTIBOD? OR CHAIN#)
L9 74 S TUMOR VIRUS PROMOTER#
L10 74 S L9(5A)(MAMMARY)
L11 1 S L10(10A)(IMMUNOGLOB? OR ANTIBOD? OR CHAIN#)

FILE 'EPOABS' ENTERED AT 09:36:47 ON 10 AUG 1999

L12 0 S L11
L13 13 S (METALLOTH?)(3A)(PROMOTER#)
L14 0 S L13(10A)(MAMMARY)

FILE 'USPAT' ENTERED AT 09:37:55 ON 10 AUG 1999

L15 52 S L14
L16 1 S L15(10A)(IMMUNOGLOB? OR ANTIBOD? OR CHAIN#)
L17 37141 S IMMUNOGLOB? OR ANTIBOD?
L18 3157 S L17(3A)(LINKED)
L19 16 S L18(5A)(PROMOTER#)
L20 0 S L19(5A)(MAMMARY)

=> s 117(5a)(recombinant)

L21 21367 RECOMBINANT
L22 2752 L17(5A)(RECOMBINANT)

=> s 117(3a)(recombinant)

L23 21367 RECOMBINANT
L24 2094 L17(3A)(RECOMBINANT)

=> s 122(5a)(construct# or plasmid# or vector#)

96626 CONSTRUCT#
16628 PLASMID#
77567 VECTOR#
L25 84 L22(5A)(CONSTRUCT# OR PLASMID# OR VECTOR#)

=> s I23(5a)(promoter#)

36145 PROMOTER#

L24 0 L23(5A)(PROMOTER#)

=> s I23 and mammary

4414 MAMMARY

L25 9 L23 AND MAMMARY

=> d 1- cit ab

1. 5,912,160, Jun. 15, 1999, Gab1, Grb2 binding protein, and compositions for making and methods of using the same; Albert J. Wong, et al., 435/252.3, 69.1, 320.1; 530/350; 536/23.5, 24.3 [IMAGE AVAILABLE]

US PAT NO: 5,912,160 [IMAGE AVAILABLE] L25: 1 of 9

ABSTRACT:

A substantially pure protein, Gab1, that binds to Grb2 is disclosed. Isolated nucleic acid molecules that encode Gab1 is disclosed. Pharmaceutical compositions comprising a pharmaceutically acceptable carrier in combination with nucleic acid molecules are disclosed. Fragments of nucleic acid molecules that encode Gab1 having at least 10 nucleotides and oligonucleotide molecule comprising a nucleotide sequence complementary to a nucleotide sequence of at least 10 nucleotides are disclosed. Recombinant expression vectors that comprise the nucleic acid molecule that encode Gab1, and host cells that comprise such **recombinant** vectors** are disclosed. **Antibodies** that bind to an epitope on Gab1 are disclosed. Methods of identifying inhibitors, activators and substrates of Gab1 are disclosed. Antisense compounds and methods of using the same are disclosed.

2. 5,880,268, Mar. 9, 1999, Modulators of the interaction between ICAM-R and .alpha. sub.d /CD18; W. Michael Gallatin, et al., 530/387.3, 387.9, 388.1, 388.22 [IMAGE AVAILABLE]

US PAT NO: 5,880,268 [IMAGE AVAILABLE] L25: 2 of 9

ABSTRACT:

DNA sequences encoding a novel human intercellular adhesion molecule polypeptide (designated "ICAM-R") and variants thereof are disclosed along with methods and materials for production of the same by recombinant procedures. Binding molecules specific for ICAM-R and

variants thereof are also disclosed as useful in both the isolation of ICAM-R from natural cellular sources and the modulation of ligand/receptor binding biological activities of ICAM-R. Specifically, antibody substances which modulate the interaction between ICAM-R and ad/CD18 are provided.

3. 5,869,262, Feb. 9, 1999, Method for monitoring an inflammatory disease state by detecting circulating ICAM-R; W. Michael Gallatin, et al., 435/7.1, 7.92, 7.94, 7.95; 436/811 [IMAGE AVAILABLE]

US PAT NO: 5,869,262 [IMAGE AVAILABLE] L25: 3 of 9

ABSTRACT:

Methods for monitoring the progression of systemic lupus erythematosus (SLE) in a patient by detecting elevated levels of circulating ICAM-R wherein progression is indicated in an SLE patient whose circulating ICAM-R levels are increased as compared to normal individuals or individuals with in active SLE. Methods for the detection of an inflammatory disease state selected from the group consisting of rheumatoid arthritis, SLE, and Guillain-Barre syndrome and multiple sclerosis in a patient by detecting elevated levels of circulating ICAM-R wherein the presence of the inflammatory disease state is indicated in a patient whose circulating ICAM-R levels are increased as compared to normal healthy individuals. ICAM-R is also known as ICAM-3 and CDw50 in the art.

4. 5,837,822, Nov. 17, 1998, Humanized antibodies specific for ICAM related protein; W. Michael Gallatin, et al., 530/387.3, 388.1, 388.22 [IMAGE AVAILABLE]

US PAT NO: 5,837,822 [IMAGE AVAILABLE] L25: 4 of 9

ABSTRACT:

DNA sequences encoding a novel human intercellular adhesion molecule polypeptide (designated "ICAM-R") and variants thereof are disclosed along with methods and materials for production of the same by recombinant procedures. Binding molecules specific for ICAM-R and variants thereof are also disclosed as useful in both the isolation of ICAM-R from natural cellular sources and the modulation of ligand/receptor binding biological activities of ICAM-R. More specifically, humanized antibodies specific for ICAM-R proteins are disclosed.

5. 5,811,517, Sep. 22, 1998, ICAM-related protein variants; W. Michael Gallatin, et al., 530/350; 435/69.1, 69.7, 252.3, 320.1, 325; 536/23.1, 23.4 [IMAGE AVAILABLE]

US PAT NO: 5,811,517 [IMAGE AVAILABLE] L25: 5 of 9

ABSTRACT:

DNA sequences encoding a novel human intercellular adhesion molecule polypeptide (designated "ICAM-R") and variants thereof are disclosed along with methods and materials for production of the same by recombinant procedures. Binding molecules specific for ICAM-R and variants thereof are also disclosed as useful in both the isolation of ICAM-R from natural cellular sources and the modulation of ligand/receptor binding biological activities of ICAM-R.

6. 5,773,218, Jun. 30, 1998, Method to identify compounds which modulate ICAM-related protein interactions; W. Michael Gallatin, et al., 435/6 [IMAGE AVAILABLE]

US PAT NO: 5,773,218 [IMAGE AVAILABLE] L25: 6 of 9

ABSTRACT:

DNA sequences encoding a novel human intercellular adhesion molecule polypeptide (designated "ICAM-R") and variants thereof are disclosed along with methods and materials for production of the same by recombinant procedures. Binding molecules specific for ICAM-R and variants thereof are also disclosed as useful in both the isolation of ICAM-R from natural cellular sources and the modulation of ligand/receptor binding biological activities of ICAM-R.

7. 5,672,500, Sep. 30, 1997, Mch2, an apoptotic cysteine protease, and compositions for making and methods of using the same; Gerald Litwack, et al., 435/252.3, 320.1; 530/350; 536/23.2 [IMAGE AVAILABLE]

US PAT NO: 5,672,500 [IMAGE AVAILABLE] L25: 7 of 9

ABSTRACT:

A substantially pure protein that is a member of the apoptotic Ced-3/Ice cysteine protease gene family, Mch2.alpha., and an inactive isoform of it, Mch2.beta., are disclosed. Isolated nucleic acid molecules that encode Mch2.alpha. and Mch2.beta., respectively, are disclosed. Pharmaceutical compositions comprising a pharmaceutically acceptable carrier in combination with the protein or the nucleic acid molecules are disclosed. Fragments of nucleic acid molecules that encode Mch2.alpha. and Mch2.beta. having at least 10 nucleotides and oligonucleotide molecule comprising a nucleotide sequence complimentary to a nucleotide sequence of at least 10 nucleotides are disclosed. Recombinant

expression vectors that comprise the nucleic acid molecule that encode Mch2.alpha. or Mch2.beta., and host cells that comprise such **recombinant** **vectors** are disclosed. **Antibodies** that bind to an epitope on Mch2.alpha. and/or Mch2.beta. are disclosed. Methods of identifying inhibitors, activators and substrates of Mch2.alpha. are disclosed. Antisense compounds and methods of using the same are disclosed.				SUMMARY: BSUM(112) The . . . either sequentially or simultaneously, or by using a vector construct comprising both the L-chain and H-chain genes, for example a **recombinant** single-chain **antibody** gene **construct** as indicated hereinbefore.		L2 202 S L1(5A)(PROMOTER) L3 1 S L2(10A)(ANTIBOD? OR IMMUNOGLOBULIN#) L4 299 S MAMMARY(5A)(PROMOTER#) L5 3 S L4(10A)(ANTIBOD? OR IMMUNOGLOB? OR CHAIN#) L6 888 S MOUSE MAMMARY L7 210 S L6(5A)(PROMOTER#) L8 1 S L7(5A)(IMMUNOGLOB? OR ANTIBOD? OR CHAIN#) L9 74 S TUMOR VIRUS PROMOTER# L10 74 S L9(5A)(MAMMARY) L11 1 S L10(10A)(IMMUNOGLOB? OR ANTIBOD? OR CHAIN#)	
8. 5,631,133, May 20, 1997, Transition in transcriptional activation by intracellular hormone receptors at the tumor stage of dermal fibrosarcoma development; Douglas Hanahan, et al., 435/6, 69.4 [IMAGE AVAILABLE]				SUMMARY: BSUM(113) Preferred are host cells transformed with a **recombinant** single-chain **antibody** gene **construct** comprising DNA coding for the heavy chain variable domain of an anti-c-erbB-2 antibody, DNA coding for a spacer group, DNA. . . chain variable domain of an anti-c-erbB-2 antibody and DNA coding for an effector molecule, in particular transfected with the preferred **recombinant** single-chain **antibody** gene **construct** as indicated hereinbefore. Further examples of host cells of the invention are cells transferred with similar recombinant plasmids which contain.		FILE 'EPOABS' ENTERED AT 09:36:47 ON 10 AUG 1999 L12 0 S L11 L13 13 S (METALLOTH?)(3A)(PROMOTER#) L14 0 S L13(10A)(MAMMARY) FILE 'USPAT' ENTERED AT 09:37:55 ON 10 AUG 1999 L15 52 S L14 L16 1 S L15(10A)(IMMUNOGLOB? OR ANTIBOD? OR CHAIN#) L17 37141 S IMMUNOGLOB? OR ANTIBOD? L18 3157 S L17(5A)(LINKED) L19 16 S L18(5A)(PROMOTER#) L20 0 S L19(5A)(MAMMARY) L21 2752 S L17(5A)(RECOMBINANT) L22 2094 S L17(3A)(RECOMBINANT) L23 84 S L22(5A)(CONSTRUCT# OR PLASMID# OR VECTOR#) L24 0 S L23(5A)(PROMOTER#) L25 9 S L23 AND MAMMARY	
US PAT NO: 5,631,133 [IMAGE AVAILABLE] L25: 8 of 9						==> s antibody? or immunoglobul? 36150 ANTIBOD? 11437 IMMUNOGLOBUL? L26 37116 ANTIBOD? OR IMMUNOGLOBUL? ==> s l26(10a)mlk 29073 MILK L27 721 L26(10A)MILK ==> s l26(5a)mlk 29073 MILK L28 393 L26(5A)MILK ==> s l28(10a)(recombinant) 21367 RECOMBINANT L29 8 L28(10A)(RECOMBINANT) ==> d l - cit ab	
ABSTRACT: Intracellular hormone receptors are discovered to undergo posttranslational regulation. Assays to assess cancer progression and to permit discovery of a new class of biologically active compounds are provided. Related kits are also provided.				1.3.1 . . . immunofluorescent staining of mouse cells expressing high levels of the human c-erbB-2 protein. To isolate these cells the HC11 mouse **mammary** epithelial cell line (Ball at al., EMBO J. 7: 2089, 1988) is transfected according to conventional, previously described methods (Graham. . .			
9. 5,571,894, Nov. 5, 1996, Recombinant antibodies specific for a growth factor receptor; Winfried S. Wells, et al., 530/387.3; 435/69.1; 530/350; 536/23.4 [IMAGE AVAILABLE]				DETDESC: DETD(7) 15.1 Immunotoxin treatment of cell lines: Human breast and ovarian tumor cell lines SK-BR3, MDAMB-231, MDA-MB-453, HTB77, the mouse **mammary** epithelial cell line HC11, and HC11 cells transfected with the human c-erbB-2 cDNA are plated on 48 well tissue culture. . . ==> d his			
US PAT NO: 5,571,894 [IMAGE AVAILABLE] L25: 9 of 9						(FILE 'USPAT' ENTERED AT 09:28:32 ON 10 AUG 1999) L1 1307 S MAMMARY TUMOR	
ABSTRACT: The invention concerns recombinant antibodies directed to the extracellular domain of the human growth factor receptor c-erbB-2 comprising a light chain variable domain and a heavy chain variable domain of a monoclonal antibody, monoclonal antibodies directed to c-erbB-2 themselves, a method of manufacturing those recombinant and monoclonal antibodies, hybridoma cells secreting those monoclonal antibodies, a method of manufacturing those hybridoma cells, DNAs encoding the heavy and light chain variable domains and the recombinant antibody, a method of manufacturing that DNA, hybrid vectors suitable for the expression of that DNA, host cells transformed with that DNA, and processes of using those recombinant and monoclonal antibodies in the diagnosis and treatment of tumors.							
==> d 9 kwic							
US PAT NO: 5,571,894 [IMAGE AVAILABLE] L25: 9 of 9							

preferred aspect of the invention, the adenosine deaminase is conjugated

conjugated to one or more strands of polyethylene glycol to prolonged activity in vivo.

7. 5,691,135, Nov. 25, 1997, Immunoglobulin superantigen binding to gp120 from HIV; Jonathan Braun, et al., 435/5, 7.1, 7.2, 7.24, 7.92, 974 [IMAGE AVAILABLE]

US PAT NO: 5,691,135 [IMAGE AVAILABLE]

ABSTRACT: VH3 and VH4 type immunoglobulins display superantigen-type binding affinity for the HIV gp120 envelope glycoprotein. VH3 and VH4 type antibody molecules, including IgG and IgM, are shown to suppress infection *in vivo* and *in vitro*. Determining the level of such antibody molecules is correlated to the advancement of HIV disease state.

8. 4, 873,316, Oct. 10, 1989, Isolation of exogenous recombinant proteins from the milk of transgenic mammals: Harry Meade, et al., 80077; 435/69.1, 69.2, 69.4, 69.5, 69.6, 69.8, 530/360, 361, 416, 417, 418, 832, 833; 536/23.1, 23.4, 23.5; 800/18 [IMAGE AVAILABLE]

US PAT NO: 4,873,316 [IMAGE AVAILABLE]

ABSTRACT: This invention relates to the production of recombinant proteins in mammals' milk. Particularly, this invention relates to an expression system comprising the mammal's casein promoter which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.

\Rightarrow d his

L1	1307 S MAMMARY TUMOR
L2	202 S L1(5A)(PROMOTER)
L3	1 S L2(10A)(ANTIBOD)? OR IMMUNOGLOBULIN#)
L4	299 S MAMMARY(5A)(PROMOTER#)
L5	3 S L4(10A)(ANTIBOD)? OR IMMUNOGLOB? OR
CHAIN#)	
L6	888 S MOUSE MAMMARY
L7	210 S L6(5A)(PROMOTER#)
L8	1 S L7(5A)(IMMUNOGLOB? OR ANTIBOD? OR

...molecules encoding the said polypeptides, and to subproducts comprising the said DNA molecules. The invention further relates to processes for producing the said RSSI variants and for producing transgenic plants containing the said variants.

Furthermore the invention relates to such transgenic animals as well as to infant formulas comprising milk from such transgenic animals. The invention also relates to pharmaceutical compositions comprising the said polypeptides; and the use of the said polypeptides and DNA molecules for the manufacture of medicaments.

5. 5,750,172, May 12, 1998, Transgenic non human mammal milk;
Harry
Meade, et al., 426/580, 435/69.1, 69.4, 69.51, 69.52, 69.6, 183, 215;
800/7 [IMAGE AVAILABLE]

US PAT NO: 5,750,172 [IMAGE AVAILABLE]

ABSTRACT: This invention relates to the production of recombinant proteins, such as coagulation factors VIII and IX, tissue plasminogen activator (TPA), urokinase, growth hormone, insulin, interferons, interleukins, peptide hormones and immunoglobulins, in mammals' milk. Particularly, this invention relates to an expression system which when transgenically incorporated into a mammal permits the female species of that mammal to produce the desired recombinant protein in or along with its milk. This invention also relates to the transgenic mammal that produces the desired recombinant product in its milk.

6. 5,728,560, Mar. 17, 1998, Method of treating CD4+ T cell lymphopenia in immuno-compromised patients; Robert G. L. Shorr, et al., 435/103, 188, 227; 514/4, 21, 46 [IMAGE AVAILABLE]

US PAT NO: 5,728,560 [IMAGE AVAILABLE]

ABSTRACT: The present invention is directed to methods of treating CD4+ T cell lymphopenia in HIV-infected patients. The methods include administering an effective amount of adenosine deaminase or related enzymatic material to a patient in need thereof. In preferred aspects of the invention, the method is employed in conjunction with HIV-infected patients having CD4+ T cell levels of less than about 200/ μ ml. Effective amounts of the enzyme range from about 5 to about 50 IU/kg/week. In one particularly

1. 5,849,992, Dec. 15, 1998, Transgenic production of antibodies in milk; Harry Meade, et al., 800/14, 7, 15, 16, 17, 18 [IMAGE AVAILABLE]

US PAT NO: 5,849,992 [IMAGE AVAILABLE]

ABSTRACT:
A method for the production of monoclonal antibodies in mammal's milk through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.

2. 5,827,690, Oct. 27, 1998, Transgenic production of antibodies in milk: Harry Meade et al. 800/7-530/867 [IMAGE AVAILABLE]

US PAT NO: 5,827,690 [IMAGE AVAILABLE]

ABSTRACT: A method for the production of monoclonal antibodies in mammal's milk, through the creation of transgenic animals that selectively express foreign antibody genes in mammary epithelial cells.

3. 5,827,683, Oct. 27, 1998, Nucleic acids encoding BSSL variants;
Lars
Gustav Blackberg, et al., 435/69.1, 69.7, 70.1, 70.3, 71.1, 200, 243,
270.1 325-536/23.1 273.2 23.5 [IMAGE AVAILABLE]

US PAT NO: 5,827,683 [IMAGE AVAILABLE]

ABSTRACT: The invention discloses nucleic acids encoding variant Bile Salt Stimulated Lipase (BSSL; EC 3.1.1.1). The encoded variant BSSL enzymes maintain catalytic activity but contain fewer glycosylation sites that full-length BSSL. This reduced glycosylation facilitates purification and characterization of recombinant BSSL proteins.

4. 5,763,739, Jun. 9, 1998, Transgenic non-human mammals producing BSSL variants; Lars Gustav Blackberg, et al., 800/18; 435/69.1; 800/7, 14, 15, 16 17 21 25 [IMAGE AVAILABLE]

US PAT NO: 5,763,739 [IMAGE AVAILABLE]

ABSTRACT: The present invention relates to novel polypeptides which are variants of Bile Salt-Stimulated Lipase (BSSL; EC 3.1.1.1). It also relates to DNA

CHAIN#)
L9 74 S TUMOR VIRUS PROMOTER#
L10 74 S L9(5A)(MAMMARY)
L11 1 S L10(10A)(IMMUNOGLOB? OR ANTIBOD? OR
CHAIN#)
FILE 'EPOABS' ENTERED AT 09:36:47 ON 10 AUG 1999
L12 0 S L11
L13 13 S (METALLOTH?)(3A)(PROMOTER#)
L14 0 S L13(10A)(MAMMARY)
FILE 'USPAT' ENTERED AT 09:37:55 ON 10 AUG 1999
L15 52 S L14
L16 1 S L15(10A)(IMMUNOGLOB? OR ANTIBOD? OR
CHAIN#)
L17 37141 S IMMUNOGLOB? OR ANTIBOD?
L18 3157 S L17(5A)(LINKED)
L19 16 S L18(5A)(PROMOTER#)
L20 0 S L19(5A)(MAMMARY)
L21 2752 S L17(5A)(RECOMBINANT)
L22 2094 S L17(3A)(RECOMBINANT)
L23 84 S L22(5A)(CONSTRUCT# OR PLASMID# OR
VECTOR#)
L24 0 S L23(5A)(PROMOTER#)
L25 9 S L23 AND MAMMARY
L26 37116 S ANTIBOD? OR IMMUNOGLOBUL?
L27 721 S L26(10A)MILK
L28 393 S L26(5A)MILK
L29 8 S L28(10A)(RECOMBINANT)

=> log y

U.S. Patent & Trademark Office LOGOFF AT 09:50:37 ON 10 AUG
1999

FILE 'USPAT' ENTERED AT 16:02:55 ON 06 AUG 1999

* U.S. PATENT TEXT FILE *
*
* THE WEEKLY PATENT TEXT AND IMAGE DATA IS
CURRENT *
* THROUGH AUGUST 3, 1999 *
* *
* *****
=> s immunoglobulin#(10a)(construct#)
11365 IMMUNOGLOBULIN#
96404 CONSTRUCT#
L1 139 IMMUNOGLOBULIN#(10A)(CONSTRUCT#)
=> s immunoglobulin#(5a)(construct#)
11365 IMMUNOGLOBULIN#
96404 CONSTRUCT#
L2 90 IMMUNOGLOBULIN#(5A)(CONSTRUCT#)
=> s l2(10a)(light chain)
679503 LIGHT
318197 CHAIN
2289 LIGHT CHAIN
(LIGHT(W)CHAIN)
L3 5 L2(10A)(LIGHT CHAIN)
=> s l3(10a)(heavy chain)
226973 HEAVY
318197 CHAIN
2515 HEAVY CHAIN
(HEAVY(W)CHAIN)
L4 4 L3(10A)(HEAVY CHAIN)
=> d l- cit ab

1. 5,891,717, Apr. 6, 1999, Methods and compositions for inhibiting hexokinase; Christopher B. Newgard, et al., 435/325, 69 I, 69 7, 194, 320 I, 455, 456, 458, 463; 536/23.2, 23.4 [IMAGE AVAILABLE]
US PAT NO: 5,891,717 [IMAGE AVAILABLE] L4: 1 of 4
ABSTRACT:
Disclosed are compositions and methods for inhibiting hexokinase enzymes in mammalian cells. Specifically provided are proteins that stimulate the production of trehalose-6-phosphate and their respective genes;

hexokinase-specific ribozymes and genes encoding such constructs; and agents that competitively reduce hexokinase activity, e.g., by displacing hexokinase from mitochondria, and their respective genes. The latter group of agents includes inactive hexokinases and fragments thereof that retain mitochondrial binding functions and hexokinase-glucokinase chimeras that further substitute glucokinase activity for hexokinase activity. Mammalian cells including such hexokinase inhibitors, methods of making such cells and various in vitro and in vivo methods of using cells with reduced hexokinase activity are also described herein.
2. 5,854,067, Dec. 29, 1998, Hexokinase inhibitors; Christopher B. Newgard, et al., 435/366, 4, 6, 91.1, 91.31, 183, 320.1, 325; 536/23.1, 24.31, 24.5 [IMAGE AVAILABLE]
US PAT NO: 5,854,067 [IMAGE AVAILABLE] L4: 2 of 4
ABSTRACT:
Disclosed are compositions and methods for inhibiting hexokinase enzymes in mammalian cells. Specifically provided are proteins that stimulate the production of trehalose-6-phosphate and their respective genes; hexokinase-specific ribozymes and genes encoding such constructs; and agents that competitively reduce hexokinase activity, e.g., by displacing hexokinase from mitochondria, and their respective genes. The latter group of agents includes inactive hexokinases and fragments thereof that retain mitochondrial binding functions and hexokinase-glucokinase chimeras that further substitute glucokinase activity for hexokinase activity. Mammalian cells including such hexokinase inhibitors, methods of making such cells and various in vitro and in vivo methods of using cells with reduced hexokinase activity are also described herein.
3. 5,786,213, Jul. 28, 1998, Inhibition of endogenous gastrin expression for treatment of colorectal cancer; Pomila Singh, et al., 435/320.1; 424/93.21; 435/69.1, 325; 514/2, 44; 536/23.1, 24.3 [IMAGE AVAILABLE]
US PAT NO: 5,786,213 [IMAGE AVAILABLE] L4: 3 of 4
ABSTRACT:
The present invention discloses is for the treatment of colon cancer. The expression of gastrin by colon cancers is inhibited by the use of antisense gastrin expression. Methods are disclosed for the preparation of expression constructs and the use of such constructs to inhibit colon cancer growth.
4. 5,610,034, Mar. 11, 1997, Immunoglobulin production by

trichoderma;
Eni Nyyssönen, et al., 435/69.6, 69.8, 254.6, 320.1, 484; 536/23.53, 24.1 [IMAGE AVAILABLE]
US PAT NO: 5,610,034 [IMAGE AVAILABLE] L4: 4 of 4
ABSTRACT:
Methods for the production of recombinant immunoglobulins in a Trichoderma host are described.
For
=> s l2(p)(assembly?)
807526 ASSEMBL?
L5 6 L2(P)(ASSEMBL?)
=> d l- cit ab

1. 5,831,036, Nov. 3, 1998, Soluble fragments of human intercellular adhesion molecule-1; Timothy A. Springer, et al., 530/395; 424/185.1; 435/69.3; 530/300, 350 [IMAGE AVAILABLE]
US PAT NO: 5,831,036 [IMAGE AVAILABLE] L5: 1 of 6
ABSTRACT:
The present invention relates to intercellular adhesion molecules (ICAM-1) which are involved in the process through which lymphocytes recognize and migrate to sites of inflammation as well as attach to cellular substrates during inflammation. The invention is directed toward such molecules, screening assays for identifying such molecules and antibodies capable of binding such molecules. The invention also includes uses for adhesion molecules and for the antibodies that are capable of binding them.
2. 5,639,947, Jun. 17, 1997, Compositions containing glycopolypeptide multimers and methods of making same in plants; Andrew C. Hiant, et al., 800/267; 435/69.6; 530/387.1, 387.3; 536/23.53; 800/288, 298 [IMAGE AVAILABLE]
US PAT NO: 5,639,947 [IMAGE AVAILABLE] L5: 2 of 6
ABSTRACT:
The present invention contemplates a transgenic plant having somatic and germ cells containing at least two mammalian genes coding for polypeptides capable of autogenously associating with each other to form a biologically active multimer. In addition, the invention describes a method for producing a glycopolypeptide multimer by introducing first

and second mammalian genes encoding the constituent parts of the multimer into first and second respective members of a plant species, generating a progeny from the first and second plant species members, and isolating the glycopolypeptide multimer from the progeny plant.

3. 5,612,216, Mar. 18, 1997, Nucleotide sequence encoding intercellular adhesion molecule-1 and fragments thereof; Timothy A. Springer, et al., 435/252.3, 69.1, 320.1, 530/395, 536/23.5 [IMAGE AVAILABLE]

US PAT NO: 5,612,216 [IMAGE AVAILABLE] L5: 3 of 6

ABSTRACT:
The present invention relates to intercellular adhesion molecules (ICAM-1) which are involved in the process through which lymphocytes recognize and migrate to sites of inflammation as well as attach to cellular substrates during inflammation. The invention is directed toward such molecules, screening assays for identifying such molecules and antibodies capable of binding such molecules. The invention also includes uses for adhesion molecules and for the antibodies that are capable of binding them.

4. 5,475,091, Dec. 12, 1995, R6-5-D6, an antibody which binds intercellular adhesion molecule-1; Timothy A. Springer, et al., 530/388.22, 388.85, 389.2 [IMAGE AVAILABLE]

US PAT NO: 5,475,091 [IMAGE AVAILABLE] L5: 4 of 6

ABSTRACT:
The present invention relates to intercellular adhesion molecules (ICAM-1) which are involved in the process through which lymphocytes recognize and migrate to sites of inflammation as well as attach to cellular substrates during inflammation. The invention is directed toward such molecules, screening assays for identifying such molecules and antibodies capable of binding such molecules. The invention also includes uses for adhesion molecules and for the antibodies that are capable of binding them.

5. 5,284,931, Feb. 8, 1994, Intercellular adhesion molecules, and their binding ligands; Timothy A. Springer, et al., 424/139.1, 152.1, 153.1, 154.1, 172.1, 173.1, 514/8, 530/388.22, 395, 808, 868 [IMAGE AVAILABLE]

US PAT NO: 5,284,931 [IMAGE AVAILABLE] L5: 5 of 6

ABSTRACT:
Pharmaceutical compositions comprising antibodies to intercellular adhesion molecule-1 (ICAM-1 or CD54) are useful in methods of decreasing the severity of inflammation associated with the adhesion of leukocytes to cells bearing ICAM-1. Treatment with anti-ICAM-1 antibodies reduced the severity of inflammation associated with acute organ or tissue rejection and prolonged allograft survival time. Such compositions may optionally contain other immunosuppressive agents.

6. 5,202,422, Apr. 13, 1993, Compositions containing plant-produced glycopolypeptide multimers, multimeric proteins and method of their use; Andrew C. Hiatt, et al., 424/132.1, 133.1, 150.1, 804, 435/69.6, 70.21, 188.5, 252.3, 320.1, 530/387.1, 387.3, 388.1, 388.4, 861, 800/288 [IMAGE AVAILABLE]

US PAT NO: 5,202,422 [IMAGE AVAILABLE] L5: 6 of 6

ABSTRACT:
The present invention contemplates glycopolypeptide multimers having a polypeptide that contain an immunoglobulin amino acid residue sequence and an oligosaccharide that comprises a core pentasaccharide and N-acetylglucosamine-containing outer branches, such that the multimer is free from sialic acid. The production of passive immunity in an animal by administering a sialic acid free glycopolypeptide multimer is also contemplated. In addition, the invention describes a method for producing a glycopolypeptide multimer by introducing first and second mammalian genes encoding the constituent parts of the multimer into first and second respective members of a plant species, generating a progeny from the first and second plant species members, and isolating the glycopolypeptide multimer from the progeny plant.

=> s (immunoglobulin# or antibody#)(10a)(sequence#)

11365 IMMUNOGLOBULIN#
36023 ANTIBODY#
366097 SEQUENCE#

L6 5358 (IMMUNOGLOBULIN# OR ANTIBODY#)(10a)(SEQUENCE#)

=> s l6 and (assemb? or function?)

807563 ASSEMB?
1160058 FUNCTION?

L7 4899 L6 AND (ASSEMB? OR FUNCTION?)

adhesion molecule-1 (ICAM-1 or CD54) are useful in methods of decreasing the severity of inflammation associated with the adhesion of leukocytes to cells bearing ICAM-1. Treatment with anti-ICAM-1 antibodies reduced the severity of inflammation associated with acute organ or tissue rejection and prolonged allograft survival time. Such compositions may optionally contain other immunosuppressive agents.

280. 4,935,496, Jun. 19, 1990, Mouse-human chimaeric immunoglobulin heavy chain specific for the call antigen; Akira Kudo, et al., 530/387.3, 388.15, 388.73, 388.75, 808, 809, 828, 866, 867 [IMAGE AVAILABLE]

US PAT NO: 4,935,496 [IMAGE AVAILABLE] L8: 280 of 292

ABSTRACT:
A mouse-human chimaeric immunoglobulin heavy chain comprising (a) the amino acid sequence of a mouse immunoglobulin heavy chain variable region and (b) the amino acid sequence of a human immunoglobulin heavy chain constant region and reacting specifically with human common acute lymphocytic leukemia antigen and a chimaeric DNA fragment which encodes the amino acid sequence of the above mouse-human chimaeric immunoglobulin heavy chain.

281. 4,920,213, Apr. 24, 1990, Method and compositions useful in preventing equine influenza; Beverly Dale, et al., 536/23.72; 435/69.1, 69.3, 200, 201, 235.1, 320.1, 536/23.2, 23.7 [IMAGE AVAILABLE]

US PAT NO: 4,920,213 [IMAGE AVAILABLE] L8: 281 of 292

ABSTRACT:
Recombinant vaccines for immunizing horses against equine influenza virus (EIV) are disclosed. The DNA sequences encoding the hemagglutinin (HA) and neuraminidase (NA) glycoproteins from the two strains of EIV currently infective in horses are used to construct vaccinia carried vaccines, to design synthetic peptides for primer and booster administration, and to permit recombinant synthesis of HA and/or NA protein based vaccines. These DNA sequences also provide probes useful for preparing similar vaccines from fresh isolates of new strains generated by genetic drift.

282. 4,906,564, Mar. 6, 1990, Antigenic determinants recognized by antibodies obtained using a pathogenic agent or a derivative thereof that presents a restricted set of antigens; Jeffery A. Lyon, et al., 435/7.22,

- enhancer element. Methods of producing non-immunoglobulin protein and DNA molecules are also provided.
285. 4,834,976, May 30, 1989, Monoclonal antibodies to pseudomonas aeruginosa flagella; Mae J. Rosok, et al., 424/142.1, 150.1; 435/7.3, 340, 804, 875; 436/512, 513, 519, 548, 811; 530/388.15, 388.4 [IMAGE AVAILABLE]
- US PAT NO: 4,834,976 [IMAGE AVAILABLE] L8: 285 of 292
- ABSTRACT:
Cell lines have been produced that secrete monoclonal antibodies capable of binding to the flagellar proteins of selected Pseudomonas aeruginosa strains. Some of these antibodies have been found to be protective against lethal challenges of P. aeruginosa. Pharmaceutical compositions containing these antibodies, which can be in combination with other monoclonal antibodies, blood plasma fractions and antimicrobial agents, and the prophylactic and therapeutic use of such compositions in the management of infections, are included.
- Prior to filing this application, the continuous transformed cell lines PaF4 IVE8, FA6 IIG5, 20H11, and 21B8, described herein, were deposited in the America Type Culture Collection and given the designations HB9129, HB9130, CRL 9300, and CRL 9301, respectively.
286. 4,806,312, Feb. 21, 1989, Multizone analytical element having detectable signal concentrating zone; Alfred C. Greenquist, 422/56, 57, 58; 435/7.7, 7.72, 7.92, 805, 968; 436/807, 810, 815 [IMAGE AVAILABLE]
- US PAT NO: 4,806,312 [IMAGE AVAILABLE] L8: 286 of 292
- ABSTRACT:
A multizone test device for the determination of analyte from a liquid test medium upon contact with the liquid test medium and a labeled reagent comprising a chemical group having a detectable chemical property. The test device preferably comprises multilayers including a reagent layer incorporated with an immobilized reagent and a detection layer incorporated with an immobilized form of an interactive detection reagent for the labeled reagent. The immobilized reagent in the reagent layer and the labeled reagent comprise specific binding partners which will bind to each other dependent upon the amount of analyte present. The reagent layer migrates into the detection layer and becomes bound to and immobilized by the immobilized binding substance therein. As a result, reverse migration of the labeled reagent into the reagent layer is prevented and the detectable physical property provided by the label of the labeled reagent is localized in the detection layer for the precise measurement thereof and correlation to the amount of analyte in the test medium.
288. 4,803,156, Feb. 7, 1989, Peptide-beta-lactamase conjugates for enzyme-linked immunoassays; Alexander R. Neurath, et al., 435/5, 7.92, 18, 19; 436/820, 828, 930/142, 200, 221, 222, 223, 260, 310, DIG.820 [IMAGE AVAILABLE]
- US PAT NO: 4,803,156 [IMAGE AVAILABLE] L8: 288 of 292
- ABSTRACT:
A reagent for an ELISA determination of an antibody, the reagent comprising a peptide covalently linked to beta-lactamase. The reagent can be used in the following method to detect antibodies in a sample
- 5, 29, 530/350, 388.6, 412, 413 [IMAGE AVAILABLE]
- US PAT NO: 4,906,564 [IMAGE AVAILABLE] L8: 282 of 292
- ABSTRACT:
A method provides peptides that are antigenic determinants identified by antibodies obtained using intact pathogenic agents that present a restricted set of antigens to surveillance by the immune system.
283. 4,906,562, Mar. 6, 1990, Monoclonal antibodies and antigen for human non-small cell lung carcinomas; Ingegerd Hellstrom, et al., 435/7.23, 188; 436/514, 537, 542, 547, 548; 530/387.5, 388.8, 806, 808, 828, 866 [IMAGE AVAILABLE]
- US PAT NO: 4,906,562 [IMAGE AVAILABLE] L8: 283 of 292
- ABSTRACT:
The present invention is concerned with novel monoclonal antibodies which define a glycolipid antigen associated with human non-small cell lung carcinomas ("NSCLC") and certain other human carcinomas. The antibodies bind to normal human cells to a much lesser degree than to tumor cells. The antibodies find use in diagnostic methods such as the detection of malignant cells associated with NSCLC and in therapeutic methods. Also disclosed in a novel glycolipid antigen. The invention also comprises a method for determining the presence of a malignant condition in lung tissue and other human tissue. The method involves examining the human tissue for the presence of a glycolipid antigen having the terminal carbohydrate sequence: GalNAc.beta.1.fwdarw.4Gal.beta.1.fwdarw.3GalNAc.beta.1.fwdarw.4Gal.beta.1.fwdarw.R.
284. 4,889,802, Dec. 26, 1989, Enhanced production of recombinant proteins in myeloma cells; Tristram G. Parslow, et al., 435/69.1, 69.4, 69.5, 69.51, 69.52, 69.6, 320.1, 355; 536/23.4, 23.5, 24.1 [IMAGE AVAILABLE]
- US PAT NO: 4,889,802 [IMAGE AVAILABLE] L8: 284 of 292
- ABSTRACT:
A mammalian myeloma cell comprising a double-stranded DNA molecule in its genome containing a coding sequence encoding a non-immunoglobulin protein, a non-immunoglobulin promoter sequence adjacent to the 5' terminus of said coding sequence, and the 8-base pair nucleotide sequence 5'-ATTTCAT-3' located 5' to the transcription initiation site of said promoter sequence. The DNA molecule may optionally contain an
- the immobilized interactive detection reagent therein which results in the localized generation of a detectable reaction product which preferably is also immobilized in the detection zone. As a result, reverse migration of the labeled reagent, and preferably the detectable reaction product from the detection layer is prevented and the detectable chemical property provided by the label of the labeled reagent is localized in the detection layer for the precise measurement thereof and correlation to the amount of analyte in the test medium.
287. 4,806,311, Feb. 21, 1989, Multizone analytical element having labeled reagent concentration zone; Alfred C. Greenquist, 422/56, 57, 58; 435/7.4, 7.5, 7.8, 805, 968; 436/807, 810, 815 [IMAGE AVAILABLE]
- US PAT NO: 4,806,311 [IMAGE AVAILABLE] L8: 287 of 292
- ABSTRACT:
A multizone test device for the determination of analyte from a liquid test medium upon contact with the liquid test medium and a labeled reagent comprising a chemical group having a detectable physical property. The test device preferably comprises multilayers including a reagent layer incorporated with an immobilized reagent and a detection layer incorporated with an immobilized form of a binding substance for the labeled reagent. The immobilized reagent and the labeled reagent comprise specific binding partners which will bind to each other dependent upon the amount of analyte present. Labeled reagent which does not become bound to the immobilized reagent in the reagent layer migrates into the detection layer and becomes bound to and immobilized by the immobilized binding substance therein. As a result, reverse migration of the labeled reagent into the reagent layer is prevented and the detectable physical property provided by the label of the labeled reagent is localized in the detection layer for the precise measurement thereof and correlation to the amount of analyte in the test medium.
288. 4,803,156, Feb. 7, 1989, Peptide-beta-lactamase conjugates for enzyme-linked immunoassays; Alexander R. Neurath, et al., 435/5, 7.92, 18, 19; 436/820, 828, 930/142, 200, 221, 222, 223, 260, 310, DIG.820 [IMAGE AVAILABLE]
- US PAT NO: 4,803,156 [IMAGE AVAILABLE] L8: 288 of 292
- ABSTRACT:
A reagent for an ELISA determination of an antibody, the reagent comprising a peptide covalently linked to beta-lactamase. The reagent can be used in the following method to detect antibodies in a sample

- which involves
- contacting the sample with protein A linked to a solid support,
 - incubating the sample-protein A linked to the solid support,
 - washing the incubated sample-protein A linked to the solid support,
 - contacting the washed sample-protein A with the reagent,
 - incubating the sample-protein A and reagent,
 - washing the incubated sample-protein A-reagent, and
 - determining the enzymatic activity of the resultant mass.
289. 4,631,191, Dec. 23, 1986, Methods and compositions useful in preventing equine influenza; Beverly Dale, et al., 424/186.1, 209.1; 530/324, 325, 326, 806, 811; 536/23.72; 930/220, 240 [IMAGE AVAILABLE]
- US PAT NO: 4,631,191 [IMAGE AVAILABLE] L8: 289 of 292
- ABSTRACT:**
Recombinant vaccines for immunizing horses against equine influenza virus (EIV) are disclosed. The DNA sequences encoding the hemagglutinin (HA) and neuraminidase (NA) glycoproteins from the two strains of EIV currently infective in horses are used to construct vaccinia carried vaccines, to design synthetic peptides for primer and booster administration, and to permit recombinant synthesis of HA and/or NA protein based vaccines. These DNA sequences also provide probes useful for preparing similar vaccines from fresh isolates of new strains generated by genetic drift.
290. 4,625,015, Nov. 25, 1986, Broad spectrum influenza antisera; Nicola Green, et al., 530/324; 424/139.1, 159.1, 186.1, 210.1; 530/328, 387.9, 388.3, 389.4, 403; 930/220, DIG.801, DIG.820 [IMAGE AVAILABLE]
- US PAT NO: 4,625,015 [IMAGE AVAILABLE] L8: 290 of 292
- ABSTRACT:**
Antisera against synthetic peptides which neutralize influenza viruses of differing hemagglutinin subtypes, provide protection against infection by influenza virus and methods of preparing the same are disclosed.
291. 4,489,710, Dec. 25, 1984, Composition and method for transplantation therapy; Lynn E. Spitzer, 128/898; 424/140.1, 154.1, 183.1, 809; 530/388.75; 391.7, 866 [IMAGE AVAILABLE]
- US PAT NO: 4,489,710 [IMAGE AVAILABLE] L8: 291 of 292
- ABSTRACT:**
An improved transplantation therapy and method is provided which comprises specifically killing cells known to be problematic in the transplantation process. Novel compositions of the present invention are conjugates prepared by generating antibodies specific to surface receptors of the unwanted cells, preparing Fab or F(ab') sub.2 fragments from the antibodies, and coupling the fragments to A chains of lectins or other cytotoxic agents to render the conjugates thus formed strongly cytotoxic to the cells to which the antibody was directed. The conjugates are used in vitro to eliminate unwanted cells prior to bone marrow transplantation.
292. 4,410,634, Oct. 18, 1983, Method of passively adsorbing immuno-reactive haptens to solid phases; Harold R. Cooper, et al., 436/500; 435/7.93, 961, 966; 436/532, 543, 804, 815, 823 [IMAGE AVAILABLE]
- US PAT NO: 4,410,634 [IMAGE AVAILABLE] L8: 292 of 292
- ABSTRACT:**
The method comprises covalently binding an immuno-reactive to a selected macromolecular carrier and then contacting the resulting hapten-carrier conjugate at a selected concentration in a liquid phase with a selected solid phase until a desired amount of the hapten-carrier conjugate is adsorbed to the surface of the solid phase. Unbound hapten-carrier conjugate is then separated from the solid phase, and the solid phase containing the bound hapten-carrier conjugate is recovered for use in quantitative immunoassays and the like. The solid phase can be, for example, surfaces of a test tube or microtiter well or the like. The method is simple and inexpensive and permits hapten assays of improved sensitivity.
- => s 18(p)(vector# or construct# or plasmid#)
- 77317 VECTOR#
96404 CONSTRUCT#
16546 PLASMID#
- L9 40 L8(P)(VECTOR# OR CONSTRUCT# OR PLASMID#)
- => d 1- cit ab
1. 5,919,650, Jul. 6, 1999, Method for inactivation of protein function; Mariano Barbacid, et al., 435/69.1, 320.1, 330 [IMAGE AVAILABLE]
- US PAT NO: 5,919,650 [IMAGE AVAILABLE] L9: 1 of 40
- ABSTRACT:**
Method for inactivating the function produced by a protein using an intracellularly expressed antibody or fragment thereof.
2. 5,912,133, Jun. 15, 1999, Method for isolating stem cells expressing flk-1 receptors; Ihor R. Lemischka, 435/7.21, 971; 530/388.7, 389.6 [IMAGE AVAILABLE]
- US PAT NO: 5,912,133 [IMAGE AVAILABLE] L9: 2 of 40
- ABSTRACT:**
Isolated mammalian nucleic acid molecules encoding receptor protein tyrosine kinases expressed in primitive hematopoietic cells and not expressed in mature hematopoietic cells are provided. Also included are the receptors encoded by such nucleic acid molecules; the nucleic acid molecules encoding receptor protein tyrosine kinases having the sequences shown in FIG. 1a (murine flk-2), FIG. 1b (human flk-2) and FIG. 2 (murine flk-1); the receptor protein tyrosine kinases having the amino acid sequences shown in FIG. 1a, FIG. 1b and FIG. 2; ligands for the receptors; nucleic acid sequences that encode the ligands; and methods of stimulating the proliferation and/or differentiation of primitive mammalian hematopoietic stem cells comprising contacting the stem cells with a ligand that binds to a receptor protein tyrosine kinase expressed in primitive mammalian hematopoietic cells and not expressed in mature hematopoietic cells.
3. 5,885,573, Mar. 23, 1999, Methods and materials for modulation of the immunosuppressive activity and toxicity of monoclonal antibodies; Jeffrey A. Bluestone, et al., 424/133.1, 144.1; 530/387.3 [IMAGE AVAILABLE]
- US PAT NO: 5,885,573 [IMAGE AVAILABLE] L9: 3 of 40
- ABSTRACT:**
The binding specificity of the murine OKT3 has been transferred into a human antibody framework in order to reduce its immunogenicity. This "humanized" anti-CD3 mAb (gOKT3-5) was previously shown to retain, in vitro, all the properties of native OKT3, including T cell activation which has been correlated, in vivo, with the severe side-effects observed in transplant recipients after the first administration of the mAb. Disclosed is a single amino acid mutation from a leucine to a glutamic acid at position 235 in the Fc receptor (FcR) binding segment of the gOKT3-5 mAb to produce Glu-235 mAb. Also disclosed is an amino acid mutation from the contiguous phenylalanine at position 234 to a

leucine (Leu-234).	heteromeric receptors; William D. Huse, 435/69.7, 69.1, 252.3, 320.1; 536/23.4 [IMAGE AVAILABLE]	**sequence** comprising all or a **functional** part of the DNA sequence between the EcoRI site 3.8 kb downstream of the Xho I site in the rearranged mouse .lambda..sub.1 gene and the SnaBI site 10 kb downstream of this Xho I site. The functional mouse immunoglobulin .lambda..sub.1 enhancer may comprise all or a functional part of i) the 1.3 kb first HindIII to HindII DNA fragment downstream of the EcoRI site 3.8 kb downstream of the Xho I site in the rearranged mouse .lambda..sub.1 gene, ii) the 3.3 kb HindII to HindII DNA fragment downstream of the EcoRI site 3.8 kb downstream of the Xho I site in the rearranged mouse .lambda..sub.1 gene and spanning the SnaBI site 10 kb downstream of this Xho I site.
4. 5,877,397, Mar. 2, 1999, Transgenic non-human animals capable of producing heterologous antibodies of various isotypes; Nils Lonberg, et al., 800/18; 536/23.1, 23.5, 23.53; 800/6 [IMAGE AVAILABLE]	US PAT NO: 5,871,974 [IMAGE AVAILABLE] L9: 6 of 40	8. 5,855,887, Jan. 5, 1999, Blockade of lymphocyte down-regulation associated with CTLA-4 signaling; James Patrick Allison, et al., 424/144.1, 133.1, 139.1, 143.1; 435/7.24 [IMAGE AVAILABLE]
US PAT NO: 5,877,397 [IMAGE AVAILABLE] L9: 4 of 40	ABSTRACT: A composition of matter comprising a plurality of prokaryotic cells containing diverse combinations of first and second DNA sequences encoding first and second polypeptides which form a heteromeric receptor exhibiting binding activity toward a preselected molecule, those heteromeric receptors being expressed on the surface of filamentous bacteriophage.	US PAT NO: 5,855,887 [IMAGE AVAILABLE] L9: 8 of 40
ABSTRACT: The invention relates to transgenic non-human animals capable of producing heterologous antibodies and transgenic non-human animals having inactivated endogenous immunoglobulin genes. In one aspect of the invention, endogenous immunoglobulin genes are suppressed by antisense polynucleotides and/or by antiserum directed against endogenous immunoglobulins. Heterologous antibodies are encoded by immunoglobulin genes not normally found in the genome of that species of non-human animal. In one aspect of the invention, one or more transgenes containing sequences of unrearranged heterologous human immunoglobulin heavy chains are introduced into a non-human animal thereby forming a transgenic animal capable of **functionally** rearranging transgenic **immunoglobulin** sequences** and producing a repertoire of **antibodies** of various isotypes encoded by human immunoglobulin genes.	US PAT NO: 5,859,309 [IMAGE AVAILABLE] L9: 7 of 40	ABSTRACT: T cell activation in response to antigen is increased by the administration of binding agents that block CTLA-4 signaling. When CTLA-4 signaling is thus blocked, the T cell response to antigen is released from inhibition. Such an enhanced response is useful for the treatment of tumors, chronic viral infections, and as an adjuvant during immunization.
US PAT NO: 5,877,397 [IMAGE AVAILABLE] L9: 4 of 40	ABSTRACT: Such heterologous human antibodies are produced in B-cells which are thereafter immortalized, e.g., by fusing with an immortalizing cell line such as a myeloma or by manipulating such B-cells by other techniques to perpetuate a cell line capable of producing a monoclonal heterologous antibody. The invention also relates to heavy and light chain immunoglobulin transgenes for making such transgenic non-human animals as well as methods and **vectors** for disrupting endogenous immunoglobulin loci in the transgenic animal.	9. 5,851,525, Dec. 22, 1998, Recombinant IL-5 antagonists useful in treatment of IL-5 mediated disorders; Robert S. Ames, Jr., et al., 424/145.1, 152.1, 158.1, 172.1; 530/387.1, 387.3, 388.23 [IMAGE AVAILABLE]
5. 5,874,264, Feb. 23, 1999, Gibbon ape leukemia virus receptor; Bryan Mark O'Hara, 435/6, 69.1, 320.1; 530/350; 536/23.5 [IMAGE AVAILABLE]	US PAT NO: 5,874,264 [IMAGE AVAILABLE] L9: 5 of 40	ABSTRACT: Chimeric, humanized and other IL-5 mAbs, derived from high affinity neutralizing mAbs, pharmaceutical compositions containing same, methods of treatment and diagnostics are provided.
ABSTRACT: The present invention relates to novel purified gibbon ape leukemia receptor proteins and purified DNA sequences encoding these receptor proteins.	US PAT NO: 5,840,540 [IMAGE AVAILABLE] L9: 10 of 40	10. 5,840,540, Nov. 24, 1998, Nucleic acids encoding presenilin II; Peter H. St. George-Hyslop, et al., 435/69.1, 252.3, 320.1, 325; 530/350; 536/23.1, 24.3 [IMAGE AVAILABLE]

- ABSTRACT:**
The present invention describes the identification, isolation and cloning of two human presenilin genes, PS-1 and PS-2, mutations in which lead to Familial Alzheimer's Disease. Also identified are presenilin homologous genes in mice, C. elegans and D. melanogaster. Transcripts and products of these genes are useful in detecting and diagnosing Alzheimer's disease, developing therapeutics for treatment of Alzheimer's disease, as well as the isolation and manufacture of the protein and the constructions of transgenic animals expressing the mutant genes.
11. 5,840,300, Nov. 24, 1998, Methods and compositions comprising single chain recombinant antibodies; William V. Williams, et al., 424/135.1, 148.1; 530/324, 325, 326, 388.35; 536/23.1 [IMAGE AVAILABLE]
- US PAT NO: 5,840,300 [IMAGE AVAILABLE] L9: 11 of 40
- ABSTRACT:**
Methods and compositions for the generation of single chain antibody fragments.
12. 5,817,308, Oct. 6, 1998, Tolerogenic fusion proteins of immunoglobulins and methods for inducing and maintaining tolerance; David W. Scott, et al., 424/93.21, 130.1, 133.1, 184.1, 185.1; 435/91.31, 320.1, 325, 326, 328, 514/44; 530/387.3; 536/22.1, 23.1 [IMAGE AVAILABLE]
- US PAT NO: 5,817,308 [IMAGE AVAILABLE] L9: 12 of 40
- ABSTRACT:**
The invention provides methods and compositions for inducing and maintaining tolerance to epitopes or antigens containing the epitopes. The compositions include expression cassettes and **vectors** including DNA **sequences** coding for a fusion **immunoglobulin** operably linked to transcriptional and translational control regions **functional** in a hemopoietic or lymphoid cell. The fusion immunoglobulin includes at least one heterologous tolerogenic epitope at the N-terminus variable region of the immunoglobulin. Cells stably transformed with the expression **vector** are formed and used to produce fusion immunoglobulin. The invention also provides methods for screening for novel tolerogenic epitopes and for inducing and maintaining tolerance. The methods of the invention are useful in the diagnosis and treatment of autoimmune or allergic immune responses.
13. 5,814,318, Sep. 29, 1998, Transgenic non-human animals for producing heterologous antibodies; Nils Lonberg, et al., 424/184.1; 435/69.6; 530/387.1; 536/23.1, 23.53; 800/6 [IMAGE AVAILABLE]
- US PAT NO: 5,814,318 [IMAGE AVAILABLE] L9: 13 of 40
- ABSTRACT:**
The invention relates to transgenic non-human animals capable of producing heterologous antibodies and transgenic non-human animals having inactivated endogenous immunoglobulin genes. In one aspect of the invention, endogenous immunoglobulin genes are suppressed by antisense polynucleotides and/or by antiserum directed against endogenous immunoglobulins. Heterologous antibodies are encoded by immunoglobulin genes not normally found in the genome of that species of non-human animal. In one aspect of the invention, one or more transgenes containing sequences of unrearranged heterologous human immunoglobulin heavy chains are introduced into a non-human animal thereby forming a transgenic animal capable of **functionally** rearranging transgenic **immunoglobulin** **sequences** and producing a repertoire of **antibodies** of various isotypes encoded by human immunoglobulin genes. Such heterologous human antibodies are produced in B-cells which are thereafter immortalized, e.g., by fusing with an immortalizing cell line such as a myeloma or by manipulating such B-cells by other techniques to perpetuate a cell line capable of producing a monoclonal heterologous antibody. The invention also relates to heavy and light chain immunoglobulin transgenes for making such transgenic non-human animals as well as methods and **vectors** for disrupting endogenous immunoglobulin loci in the transgenic animal.
14. 5,811,097, Sep. 22, 1998, Blockade of T lymphocyte down-regulation associated with CTLA-4 signaling; James Patrick Allison, et al., 424/144.1, 130.1, 133.1, 135.1, 141.1, 143.1, 152.1, 154.1, 810; 514/2, 12, 885; 530/387.1, 387.3, 388.1, 388.2, 388.7 [IMAGE AVAILABLE]
- US PAT NO: 5,811,097 [IMAGE AVAILABLE] L9: 14 of 40
- ABSTRACT:**
T cell activation in response to antigen is increased by the administration of binding agents that block CTLA-4 signaling. When CTLA-4 signaling is thus blocked, the T cell response to antigen is released from inhibition. Such an enhanced response is useful for the treatment
- of tumors, chronic viral infections, and as an adjuvant during immunization.
15. 5,789,650, Aug. 4, 1998, Transgenic non-human animals for producing heterologous antibodies; Nils Lonberg, et al., 800/18; 530/387.1 [IMAGE AVAILABLE]
- US PAT NO: 5,789,650 [IMAGE AVAILABLE] L9: 15 of 40
- ABSTRACT:**
The invention relates to transgenic non-human animals capable of producing heterologous antibodies and transgenic non-human animals having inactivated endogenous immunoglobulin genes. In one aspect of the invention, endogenous immunoglobulin genes are suppressed by antisense polynucleotides and/or by antiserum directed against endogenous immunoglobulins. Heterologous antibodies are encoded by immunoglobulin genes not normally found in the genome of that species of non-human animal. In one aspect of the invention, one or more transgenes containing sequences of unrearranged heterologous human immunoglobulin heavy chains are introduced into a non-human animal thereby forming a transgenic animal capable of **functionally** rearranging transgenic **immunoglobulin** **sequences** and producing a repertoire of **antibodies** of various isotypes encoded by human immunoglobulin genes. Such heterologous human antibodies are produced in B-cells which are thereafter immortalized, e.g., by fusing with an immortalizing cell line such as a myeloma or by manipulating such B-cells by other techniques to perpetuate a cell line capable of producing a monoclonal heterologous antibody. The invention also relates to heavy and light chain immunoglobulin transgenes for making such transgenic non-human animals as well as methods and **vectors** for disrupting endogenous immunoglobulin loci in the transgenic animal.
16. 5,783,420, Jul. 21, 1998, Method and compositions for controlling gene expression; Eric H. Davidson, 435/69.1, 320.1; 536/23.4, 23.53, 24.1 [IMAGE AVAILABLE]
- US PAT NO: 5,783,420 [IMAGE AVAILABLE] L9: 16 of 40
- ABSTRACT:**
The present invention is directed to methods and compositions useful for altering the transcriptional expression of genes in eukaryotic cells. The

invention employs novel antibody derivative molecules which
 function
 to recognize and bind to specific cis-regulatory DNA **sequence**
 elements of a eukaryotic gene. When two **antibody** derivative
 molecules
 are bound to adjacent cis-regulatory DNA sequence elements of a
 gene,
 those molecules may interact to form an antibody binding site which is
 capable of recognizing and binding to a transcription factor protein for
 the target gene, thereby affecting the functionality of that
 transcription factor protein and, in turn, the transcriptional activity
 of the gene. Also provided herein are isolated nucleic acids encoding
 the
 novel antibody derivative molecules of the present invention and
 expression **vectors** comprising those nucleic acids.

17. 5,783,184, Jul. 21, 1998, Method for treatment and diagnosis of
 IL-5
 mediated disorders; Edward Robert Appelbaum, et al., 424/130.1,
 133.1,
 141.1, 145.1; 435/7.1; 530/388.1, 388.23 [IMAGE AVAILABLE]

US PAT NO: 5,783,184 [IMAGE AVAILABLE] L9: 17 of
 40

ABSTRACT:
 The present invention relates to treatment and diagnosis of conditions
 mediated by IL-5 and excess eosinophil production, and more
 specifically
 to mAbs and other altered antibodies such as Fabs, chimeric, human
 humanized antibodies that do not block binding of human IL-5 to the
 alpha-chain of the human IL-5 receptor.

18. 5,776,677, Jul. 7, 1998, Methods of detecting cystic fibrosis gene
 by nucleic acid hybridization; Lap-Chee Tsui, et al., 435/6, 91.2;
 536/23.2, 24.3, 24.33 [IMAGE AVAILABLE]

US PAT NO: 5,776,677 [IMAGE AVAILABLE] L9: 18 of
 40

ABSTRACT:
 The cystic fibrosis gene and its gene product are described for both the
 normal and mutant forms. The genetic and protein information is used
 in
 developing DNA diagnosis, protein diagnosis, carrier and patient
 screening, drug and gene therapy, cloning of the gene and manufacture
 of
 the protein, and development of cystic fibrosis affected animals.

19. 5,770,449, Jun. 23, 1998, Vector for integration site independent
 gene expression in mammalian host cells which permit
 immunoglobulin gene
 expression; Sarah Jane Eccles, et al., 435/375, 69.1, 70.3, 70.4, 320.1,
 455; 514/44 [IMAGE AVAILABLE]

US PAT NO: 5,770,449 [IMAGE AVAILABLE] L9: 19 of

40

ABSTRACT:
 A **vector** for the integration of a gene into the genetic material of
 a
 mammalian host cell such that the gene may be expressed by the host
 cell.
 The **vector** comprises a promoter and the gene and in an
 immunoglobulin
 dominant control region derived from the mouse lambda.
 immunoglobulin
 gene locus capable of eliciting host cell-type restricted, integration
 site independent, copy number dependent expression of said gene. The
 DNaseI super hypersensitive site exemplified are i) about 2.35 kb
 upstream of the CAP site of the rearranged lambda.sub.1 gene, ii)
 about
 2.5 kb upstream of the genomic V lambda.sub.2 segment or iii) about
 30
 kb downstream of the rearranged lambda.sub.1 gene. Mammalian
 host cells
 transformed with the **vector** are disclosed as are transgenic
 mammals
 transformed with the **vector** and a method of producing a
 polypeptide
 comprising culturing a transformed mammalian cell. A method of gene
 therapy comprising the steps of i) removing stem cells from the body
 of a
 mammal, ii) optionally killing stem cells remaining in the body, iii)
 transforming the removed stem cells with the **vector** containing a
 gene
 deficient or absent in the body, and iv) replacing the transformed stem
 cells in the body is also disclosed. Also disclosed is **functional**
 mouse **immunoglobulin** lambda.sub.1 enhancer consisting of a
 DNA
 sequence comprising all or a **functional** part of the DNA
 sequence
 between the EcoRI site 3.8 kb downstream of the Xho I site in the
 rearranged mouse lambda.sub.1 gene and the SnaBI site 10 kb
 downstream
 of this Xho I site. The functional mouse immunoglobulin
 lambda.sub.1
 enhancer may comprise all or a functional part of i) the 1.3 kb first
 HindIII to HindIII DNA fragment downstream of the EcoRI site 3.8
 kb
 downstream of the Xho I site in the rearranged mouse lambda.sub.1
 gene,
 ii) the 3.3 kb HindIII to HindIII DNA fragment downstream of the
 EcoRI
 site 3.8 kb downstream of the Xho I site in the rearranged mouse
 lambda.sub.1 gene and spanning the SnaBI site 10 kb downstream of
 this
 Xho I site.

20. 5,747,651, May 5, 1998, Antibodies against tyrosine kinase
 receptor
 flk-1; Ihor R. Lemischka, 530/387.9, 388.22, 388.7, 389.1, 389.6
 [IMAGE

AVAILABLE]

US PAT NO: 5,747,651 [IMAGE AVAILABLE] L9: 20 of
 40

ABSTRACT:
 Isolated mammalian nucleic acid molecules encoding receptor protein
 tyrosine kinases expressed in primitive hematopoietic cells and not
 expressed in mature hematopoietic cells are provided. Also included
 are
 the receptors encoded by such nucleic acid molecules; the nucleic acid
 molecules encoding receptor protein tyrosine kinases having the
 sequences shown in FIG. 1a (murine flk-2), FIG. 1b (human flk-2) and FIG. 2
 (murine
 flk-1); the receptor protein tyrosine kinases having the amino acid
 sequences shown in FIG. 1a, FIG. 1b and FIG. 2; ligands for the
 receptors; nucleic acid sequences that encode the ligands; and methods
 of
 stimulating the proliferation and/or differentiation of primitive
 mammalian hematopoietic stem cells comprising contacting the stem
 cells
 with a ligand that binds to a receptor protein tyrosine kinase expressed
 in primitive mammalian hematopoietic cells and not expressed in
 mature
 hematopoietic cells.

21. 5,712,379, Jan. 27, 1998, Method and compositions for controlling
 gene expression; Eric H. Davidson, 536/23.4; 435/69.7; 536/23.53
 [IMAGE
 AVAILABLE]

US PAT NO: 5,712,379 [IMAGE AVAILABLE] L9: 21 of
 40

ABSTRACT:
 The present invention is directed to methods and compositions useful
 for
 altering the transcriptional expression of genes in eukaryotic cells. The
 invention employs novel antibody derivative molecules which
 function
 to recognize and bind to specific cis-regulatory DNA **sequence**
 elements of a eukaryotic gene. When two **antibody** derivative
 molecules
 are bound to adjacent cis-regulatory DNA sequence elements of a
 gene,
 those molecules may interact to form an antibody binding site which is
 capable of recognizing and binding to a transcription factor protein for
 the target gene, thereby affecting the functionality of that
 transcription factor protein and, in turn, the transcriptional activity
 of the gene. Also provided herein are isolated nucleic acids encoding
 the
 novel antibody derivative molecules of the present invention and
 expression **vectors** comprising those nucleic acids.

22. 5,698,426, Dec. 16, 1997, Surface expression libraries of
 heteromeric receptors; William D. Huse, 435/91.41, 69.1, 69.7, 320.1,

- 475; 530/387.1 [IMAGE AVAILABLE]
- US PAT NO: 5,698,426 [IMAGE AVAILABLE] L9: 22 of 40
- ABSTRACT:**
A composition of matter comprising a plurality of prokaryotic cells containing diverse combinations of first and second DNA sequences encoding first and second polypeptides which form a heteromeric receptor exhibiting binding activity toward a preselected molecule, said heteromeric receptors being expressed on the surface of filamentous bacteriophage.
23. 5,693,323, Dec. 2, 1997, Recombinant IL-5 antagonists useful in treatment of IL-5 mediated disorders; Robert S. Ames, Jr., et al., 424/145.1; 435/328, 335; 530/387.3, 388.23 [IMAGE AVAILABLE]
- US PAT NO: 5,693,323 [IMAGE AVAILABLE] L9: 23 of 40
- ABSTRACT:**
Chimeric, humanized and other IL-5 mAbs, derived from high affinity neutralizing mAbs, pharmaceutical compositions containing same, of treatment and diagnostics are provided.
24. 5,683,892, Nov. 4, 1997, DNA encoding recombinant IL-5 antagonists useful in treatment of IL-5 mediated disorders; Robert S. Ames, Jr., et al., 435/69.1, 69.3, 70.21, 252.3, 320.1, 328; 536/723.53 [IMAGE AVAILABLE]
- US PAT NO: 5,683,892 [IMAGE AVAILABLE] L9: 24 of 40
- ABSTRACT:**
DNA encoding chimeric, humanized and other IL-5 mAbs, derived from high affinity neutralizing mAbs, pharmaceutical compositions containing same, methods of treatment and diagnostics are provided.
25. 5,681,942, Oct. 28, 1997, Fanconi Anemia Type C gene; Manuel Buchwald, et al., 536/23.5, 24.2, 24.31, 24.33 [IMAGE AVAILABLE]
- US PAT NO: 5,681,942 [IMAGE AVAILABLE] L9: 25 of 40
- ABSTRACT:**
Fanconi Anemia is a human genetic disease, the precise cause of which is, to date, unknown. This invention provides an isolated human cDNA molecule which is able to specifically complement, in one type of Fanconi Anemia,
- (type C) the characteristic defect exhibited by cells derived from patients with Fanconi Anemia. The genomic gene from which this cDNA is derived is also provided as is the sequence of the protein encoded by this gene. Mutations in this gene are proposed to underlie Fanconi Anemia
- Type C. Diagnostic and therapeutic applications which derive from this work are described. The murine homolog of the human cDNA is also provided.
26. 5,661,016, Aug. 26, 1997, Transgenic non-human animals capable of producing heterologous antibodies of various isotypes; Nils Lönberg, et al., 435/452; 424/184.1; 435/91.1; 530/387.1; 536/23.1, 23.53 [IMAGE AVAILABLE]
- US PAT NO: 5,661,016 [IMAGE AVAILABLE] L9: 26 of 40
- ABSTRACT:**
The invention relates to transgenic non-human animals capable of producing heterologous antibodies and transgenic non-human animals having inactivated endogenous immunoglobulin genes. In one aspect of the invention, endogenous immunoglobulin genes are suppressed by antisense polynucleotides and/or by antiserum directed against endogenous immunoglobulins. Heterologous antibodies are encoded by immunoglobulin genes not normally found in the genome of that species of non-human animal. In one aspect of the invention, one or more transgenes containing sequences of unrearranged heterologous human immunoglobulin heavy chains are introduced into a non-human animal thereby forming a transgenic animal capable of **functionally** rearranging transgenic **immunoglobulin** sequences** and producing a repertoire of **antibodies** of various isotypes encoded by human immunoglobulin genes.
- Such heterologous human antibodies are produced in B-cells which are thereafter immortalized, e.g., by fusing with an immortalizing cell line such as a myeloma or by manipulating such B-cells by other techniques to perpetuate a cell line capable of producing a monoclonal heterologous antibody. The invention also relates to heavy and light chain immunoglobulin transgenes for making such transgenic non-human animals as well as methods and **vectors** for disrupting endogenous immunoglobulin loci in the transgenic animal.
27. 5,627,052, May 6, 1997, Methods for the production of proteins with a desired function; John W. Schrader, 435/69.6, 465; 530/387.1,
- 388.24 [IMAGE AVAILABLE]
- US PAT NO: 5,627,052 [IMAGE AVAILABLE] L9: 27 of 40
- ABSTRACT:**
The present invention provides a method for producing proteins with a desired function, generally comprising the steps of (a) providing a population of antibody-forming cells suspected of containing at least one cell capable of producing an antibody exhibiting a desired function; (b) suspending the population of antibody-forming cells in a medium, the medium having an indicator system incorporated therein, the indicator system also being capable of indicating the presence and location of a cell which forms antibodies exhibiting the desired function; (c) identifying a cell forming an antibody exhibiting the desired function; (d) isolating the identified antibody-forming cell from the medium; (e) determining the amino acid sequence of the variable region or a portion thereof which confers the desired function of the antibody produced by the isolated antibody-forming cell; and (f) synthesizing a protein with a desired function, the protein containing the amino acid sequence of the variable region or portion thereof which confers the desired function.
28. 5,601,988, Feb. 11, 1997, Immunocapture assay for cancer procoagulant antibody complex in biological samples; Stuart G. Gordon, 435/723.7, 792, 7.94, 975; 436/63, 64, 507, 813 [IMAGE AVAILABLE]
- US PAT NO: 5,601,988 [IMAGE AVAILABLE] L9: 28 of 40
- ABSTRACT:**
This invention provides a specific immunocapture ELISA for the quantitation of cancer procoagulant antibody complex (CPAC) in biological samples. In particular, this invention provides methods and techniques for specifically selecting and quantitatively measuring CPAC from a sample material using anti-CP antibodies followed by labeled anti-immunoglobulin antibodies. The amount of captured CPAC is then determined by measuring the amount of label in the captured CPAC.
29. 5,556,744, Sep. 17, 1996, Methods and compositions for diagnosing and treating certain HIV infected patients; David B. Weiner, et al., 435/5, 7.1, 974, 975; 530/324, 325, 326, 327, 328, 826 [IMAGE AVAILABLE]
- US PAT NO: 5,556,744 [IMAGE AVAILABLE] L9: 29 of 40
- ABSTRACT:**
The present invention provides a panel of HIV peptides useful in diagnosing whether or not a patient is of vertical HIV transmission

status, methods for diagnosing same, methods for identifying epitopes and peptides associated with non-transmission status, and pharmaceutical and vaccine compositions containing same.

30. 5,548,065, Aug. 20, 1996, Tyrosine kinase receptor human flk-2-specific antibodies; Ihor R. Lemischka, 530/388.22, 387.9, 388.23, 388.7, 389.2, 389.6 [IMAGE AVAILABLE]

US PAT NO: 5,548,065 [IMAGE AVAILABLE] L9: 30 of 40

ABSTRACT:
Isolated mammalian nucleic acid molecules encoding receptor protein tyrosine kinases expressed in primitive hematopoietic cells and not expressed in mature hematopoietic cells are provided. Also included are the receptors encoded by such nucleic acid molecules; the nucleic acid molecules encoding receptor protein tyrosine kinases having the sequences shown in FIG. 1a (murine flk-2), FIG. 1b (human flk-2) and FIG. 2 (murine flk-1); the receptor protein tyrosine kinases having the amino acid sequences shown in FIG. 1a, FIG. 1b and FIG. 2; ligands for the receptors; nucleic acid sequences that encode the ligands, and methods of stimulating the proliferation and/or differentiation of primitive mammalian hematopoietic stem cells comprising contacting the stem cells with a ligand that binds to a receptor protein tyrosine kinase expressed in primitive mammalian hematopoietic cells and not expressed in mature hematopoietic cells.

31. 5,545,806, Aug. 13, 1996, Ransgenic non-human animals for producing heterologous antibodies; Nils Lonberg, et al., 800/6; 424/184.1; 435/69.6, 320.1; 536/23.1, 23.5, 23.53; 800/18 [IMAGE AVAILABLE]

US PAT NO: 5,545,806 [IMAGE AVAILABLE] L9: 31 of 40

ABSTRACT:
The invention relates to transgenic non-human animals capable of producing heterologous antibodies and transgenic non-human animals having inactivated endogenous immunoglobulin genes. In one aspect of the invention, endogenous immunoglobulin genes are suppressed by antisense polynucleotides and/or by antiserum directed against endogenous immunoglobulins. Heterologous antibodies are encoded by immunoglobulin genes not normally found in the genome of that species of non-human animal. In one aspect of the invention, one or more transgenes

containing sequences of unrearranged heterologous human immunoglobulin heavy chains are introduced into a non-human animal thereby forming a transgenic animal capable of functionally rearranging transgenic immunoglobulin sequences and producing a repertoire of antibodies of various isotypes encoded by human immunoglobulin genes.

Such heterologous human antibodies are produced in B-cells which are thereafter immortalized, e.g., by fusing with an immortalizing cell line such as a myeloma or by manipulating such B-cells by other techniques to perpetuate a cell line capable of producing a monoclonal heterologous antibody. The invention also relates to heavy and light chain immunoglobulin transgenes for making such transgenic non-human animals as well as methods and vectors for disrupting endogenous immunoglobulin loci in the transgenic animal.

32. 5,429,746, Jul. 4, 1995, Antibody purification; Paula J. Shadle, et al., 210/635, 656; 530/390.5, 413, 417 [IMAGE AVAILABLE]

US PAT NO: 5,429,746 [IMAGE AVAILABLE] L9: 32 of 40

ABSTRACT:
This invention relates to the application of hydrophobic interaction chromatography combination chromatography to the purification of antibody molecule proteins.

33. 5,424,398, Jun. 13, 1995, Peptides and nucleic acid sequences related to the Epstein Barr virus; Jaap M. Middeldorp, et al., 530/350, 387.1 [IMAGE AVAILABLE]

US PAT NO: 5,424,398 [IMAGE AVAILABLE] L9: 33 of 40

ABSTRACT:
The present invention relates to peptides immunochemically reactive with antibodies to the Epstein-Barr virus (EBV), comprising at least part of the VCA-p18 or VCA-p40 protein, encoded within the EBV open reading frames BFRF3 and BpRF1 respectively, or a functional variant thereof.

The invention further relates to nucleic acid sequences encoding these peptides, monoclonal antibodies against these peptides, cell lines capable of producing monoclonal antibodies and anti-idiotypic antibodies. The invention also relates to recombinant vectors comprising a nucleic acid sequence according to the invention and host cells transformed or transfected with these vectors and reagents. The invention is further concerned with immunological reagents and methods for the detection of EBV or anti-EBV antibodies and a

method for the amplification and detection of Epstein Barr viral nucleic acid.

34. 5,414,076, May 9, 1995, DNA encoding gibbon ape leukemia virus receptor; Bryan M. O'Hara, 536/23.5; 530/324, 325, 326, 327, 328, 329, 350 [IMAGE AVAILABLE]

US PAT NO: 5,414,076 [IMAGE AVAILABLE] L9: 34 of 40

ABSTRACT:
The present invention relates to the gibbon ape leukemia virus (GALV) receptor protein and gene, as well as methods for regulating viral entry into cells.

35. 5,413,907, May 9, 1995, Diagnosis for malignant hyperthermia; Ronald G. Worton, et al., 435/6; 536/23.5, 24.31 [IMAGE AVAILABLE]

US PAT NO: 5,413,907 [IMAGE AVAILABLE] L9: 35 of 40

ABSTRACT:
A method for isolating a cDNA specific for the human ryanodine receptor is disclosed. The gene is associated with malignant hyperthermia, a hypermetabolic syndrome triggered primarily by inhalation anesthetics. The cDNA can be cloned and expressed in a recombinant plasmid or phage. The cDNA, or fragments thereof, is used as diagnostic probes for individuals at risk for malignant hyperthermia using restriction fragment length polymorphism analysis. The cDNA is that sequenced in FIG. 2 of this specification.

36. 5,367,057, Nov. 22, 1994, Tyrosine kinase receptor flk-2 and fragments thereof; Ihor R. Lemischka, 530/350, 403 [IMAGE AVAILABLE]

US PAT NO: 5,367,057 [IMAGE AVAILABLE] L9: 36 of 40

ABSTRACT:
Isolated mammalian nucleic acid molecules encoding receptor protein tyrosine kinases expressed in primitive hematopoietic cells and not expressed in mature hematopoietic cells are provided. Also included are the receptors encoded by such nucleic acid molecules; the nucleic acid molecules encoding receptor protein tyrosine kinases having the sequences shown in FIG. 1 (murine flk-2), FIG. 2 (human flk-2) and FIG. 3 (murine

flk-1); the receptor protein tyrosine kinases having the amino acid sequences shown in FIG. 1 (murine flk-2); FIG. 2 (human flk-2) and FIG.

3; ligands for the receptors; nucleic acid sequences that encode the ligands; and methods of stimulating the proliferation and/or differentiation of primitive mammalian hematopoietic stem cells comprising contacting the stem cells with a ligand that binds to a receptor protein tyrosine kinase expressed in primitive mammalian hematopoietic cells and not expressed in mature hematopoietic cells.

37. 5,358,649, Oct. 25, 1994, Diagnosis for porcine malignant hyperthermia; David H. MacLennan, et al., 435/6, 91.2; 536/24.31, 24.33 [IMAGE AVAILABLE]

US PAT NO: 5,358,649 [IMAGE AVAILABLE] L9: 37 of 40

ABSTRACT:

A purified DNA molecule comprises a DNA sequence of approximately 15.1 kb coding for normal or mutant RYR1 protein having a molecular weight of approximately 564,740 daltons. The DNA molecule has an endonuclease restriction map of FIG. 1 and a sequence of FIG. 2.

38. 5,270,458, Dec. 14, 1993, Nucleic acids encoding fragments of hematopoietic stem cell receptor flk-2; Ihor R. Lemischka, 536/23.5; 435/69.1, 320.1; 530/350, 403 [IMAGE AVAILABLE]

US PAT NO: 5,270,458 [IMAGE AVAILABLE] L9: 38 of 40

ABSTRACT:

Isolated mammalian nucleic acid molecules encoding receptor protein tyrosine kinases expressed in primitive hematopoietic cells and not expressed in mature hematopoietic cells are provided. Also included are the receptors encoded by such nucleic acid molecules; the nucleic acid molecules encoding receptor protein tyrosine kinases having the sequences shown in FIG. 1a (murine flk-2), FIG. 1b (human flk-2) and FIG. 2 (murine flk-1); the receptor protein tyrosine kinases having the amino acid sequences shown in FIG. 1a, FIG. 1b and FIG. 2; ligands for the receptors; nucleic acid sequences that encode the ligands; and methods of stimulating the proliferation and/or differentiation of primitive mammalian hematopoietic stem cells comprising contacting the stem cells with a ligand that binds to a receptor protein tyrosine kinase expressed in primitive mammalian hematopoietic cells and not expressed in mature hematopoietic cells.

39. 5,151,361, Sep. 29, 1992, Host cells expressing gibbon ape leukemia virus receptor; Bryan M. O'Hara, 435/354, 69.1, 254.2; 530/350 [IMAGE AVAILABLE]

US PAT NO: 5,151,361 [IMAGE AVAILABLE] L9: 39 of 40

ABSTRACT:

The present invention relates to novel purified gibbon ape leukemia receptor proteins and purified DNA sequences encoding these receptor proteins. The invention also relates to a method for identifying receptor proteins using the isolated DNA sequence as a probe, and a method for regulating viral entry into cells by manipulation of the GALV receptor.

40. 4,975,369, Dec. 4, 1990, Recombinant and chimeric KSI/4 antibodies directed against a human adenocarcinoma antigen; Lisa S. Beavers, et al., 435/69.1, 320.1, 464, 465; 530/387.3, 388.15, 388.85, 867; 536/23.53, 23.72 [IMAGE AVAILABLE]

US PAT NO: 4,975,369 [IMAGE AVAILABLE] L9: 40 of 40

ABSTRACT:

The present invention comprises novel recombinant DNA compounds which encode monoclonal antibody KSI/4 and chimeric derivatives of monoclonal antibody KSI/4. Eukaryotic expression vectors have been constructed that comprise novel KSI/4-encoding DNA and drive expression of KSI/4 when transformed into an appropriate host cell. The novel expression vectors can be used to create modified and chimeric derivatives of KSI/4. The recombinant-produced KSI/4, KSI/4 derivatives and KSI/4 chimeras are useful for the diagnosis, prognosis and treatment of disease states including adenocarcinoma.

U.S. Patent & Trademark Office LOGOFF AT 16:31:19 ON 06 AUG 1999